arXiv:1201.4082v1 [math.DS] 19 Jan 2012

The parameter capture map for V_3

Mary Rees

Department of Mathematical Sciences, University of Liverpool, Mathematics Building, Peach St., Liverpool L69 7ZL, U.K.
E-mail address: `maryrees@liv.ac.uk`

2000 *Mathematics Subject Classification.* 37F10,37B10

Abstract. This is a study of the Wittner capture construction for critically finite quadratic rational maps for which one critical point is periodic, and the second critical point is in the backward orbit of the first. This construction gives a way of describing rational maps up to topological conjugacy. It is known that representations as Wittner captures are not unique. We show that, in a certain parameter space which we call V_3, the set of maps with exactly 2^r representations as a Wittner capture is of density bounded from 0 for each $r \geq 0$, and for each fixed preperiod of the second critical point.

Contents

Chapter 1. Introduction 1
- 1.1. The setting 1
- 1.2. The parameter space, its maps and hyperbolic components 1
- 1.3. Counting hyperbolic components 3
- 1.4. Thurston equivalence 4
- 1.5. Capture paths and capture maps 4
- 1.6. Laminations and Lamination maps 6
- 1.7. Counting capture maps and components 7
- 1.10. So what? 10
- 1.11. The parameter capture map is not typical 11
- 1.12. Contrast with the subhyperbolic case 12
- 1.13. Organisation of the paper 12

Chapter 2. The resident's view of the main theorems 15
- 2.1. A notation for Thurston equivalence 15
- 2.2. The boundary of the universal cover 16
- 2.3. Action of mapping class groups 16
- 2.4. The Resident's View. 16
- 2.5. Key features of the fundamental domain 19
- 2.6. Key features of the fundamental domain paths 20
- 2.7. Transferring the fundamental domain under ρ 21
- 2.9. Remarks 24

Chapter 3. Symbolic dynamics, paths in $R_{m,0}$, and homeomorphisms 25
- 3.1. Symbolic Dynamics 25
- 3.2. Paths in $R_{m,0}$ 26
- 3.3. Properties of the sequences w_i, w_i', $w_i'(.,x)$ 27
- 3.4. Definition of w_1 28
- 3.5. Definition of $w_2(w)$ if $D(w) \subset D(BC)$ 29
- 3.6. Definition of w_1' 29
- 3.7. Definition of $w_1'(.,x)$ 30
- 3.8. 31
- 3.9. Basic exchangeable pairs 34
- 3.10. Exchangeable pairs for BC 38

3.11. Exchangeable pairs for (ζ, η) 39
3.12. The homeomorphisms $\psi_{m,q}$. 40
3.13. $\psi_{m,q}$ 49
3.14. First stage in the proof of Theorem 2.8 49

Chapter 4. Quadruples of paths 53
4.1. Notation 53
4.2. Principal arcs and levels 53
4.3. Principal arcs for $[\psi]$ and $R_{m,0}$ 55
4.4. The ζ-sequence $\zeta_i(x)$ 56
4.5. Quadruples 59
4.6. Quadruples of types C and AC 59
4.7. Intersections between sets E_{η_1,η_3} and $s^{-n}E_{\omega_1,\omega_3}$ 61
4.8. A basic result about the quadruple homeomorphisms 64
4.9. The first four elements of the β_i sequence 68
4.10. A general property of β quadruples 70
4.11. Satellite arcs 74
4.12. Final proof of Theorem 2.8 74

Chapter 5. Specific examples 79
5.1. Some old and new notation 79
5.3. Lemmas about the ζ sequence for $x_{r,\alpha}$ 83
5.4. The β sequence for $x_{r,\alpha}$ 92

Chapter 6. The main theorems 99
6.3. Reductions 100
6.5. Why 6.4 implies 6.1 and 6.2 103
6.6. The choice of U for Theorem 6.1 106
6.7. The choice of U for Theorem 6.2 108
6.8. Paths in $\mathcal{Z}(3/7+, -, 0)$ 112
6.9. Idea of the proof of 6.4 121
6.10. Proof of 6.4 123

Bibliography 129

CHAPTER 1

Introduction

1.1. The setting

This paper is concerned with the parameter space V_3 of quadratic rational maps:

$$h_a : z \mapsto \frac{(z-a)(z-1)}{z^2}, \quad a \in \mathbb{C}, \ a \neq 0.$$

It is a sequel to [**22**] and addresses some questions at the end of that paper. Roughly speaking, this is a series of questions about how many critically finite rational maps in V_3 can be represented by a construction called *Wittner capture* [**28**] and in how many ways. More precisely, we shall consider Wittner captures by the *aeroplane polynomial.* More formally, the questions concern the *parameter capture map for* V_3 *and the aeroplane polynomial.* The domain of the parameter capture map is a union of finite sets parametrised by m, and for each such restricted finite set domain, there is a natural restriction of the range to another finite set. We denote the restricted parameter capture map by Φ_m. The domain and range of Φ_m both have size of the order of 2^m. We are interested in the image size, and point inverse image sizes of Φ_m, for varying m.

This is, therefore, a consideration of a problem in a very specific setting. But it is also a particular case of a very basic question in dynamics: how to find simple representations of dynamical systems in a parameter space, and, at the same time, to determine the number of duplicate representations. In complex dynamics, mating, and the more easily analysed Wittner captures, are among the more popular representations. This paper is perhaps the first serious analysis of the number of duplicate representations arising.

1.2. The parameter space, its maps and hyperbolic components

The map h_a has critical points 0 and $c_2(a) = \frac{2a}{a+1}$, with corresponding critical values ∞ and $v_2(a) = -\frac{(a-1)^2}{4a}$. The point 0 is of period 3 under h_a, with orbit $0 \mapsto \infty \mapsto 1 \mapsto 0$. Every quadratic rational map with a critical point c_1 of period 3 is represented in V_3 up to conjugation by a Möbius transformation. The representative is unique if the Möbius transformation is chosen to map c_1 to 0. There are three polynomials in h_a up to Möbius conjugacy: the *rabbit*, the *antirabbit* and the *aeroplane*, corresponding to parameter values $a = a_0$ with $\mathrm{Im}(a_0) > 0$ [**22**], $\overline{a_0}$ and a_1, which is real and < 0. Therefore, with abuse of notation, we shall sometimes refer to h_a, for $a = a_0$, $\overline{a_0}$ or a_1, as "polynomials". There are also two *type II* critically finite maps in V_3 given by parameter values $a = \pm 1$. We see that $c_2(1) = 1$ and $c_2(-1) = \infty$, so that, for both these parameter values, the critical points lie in the same period three orbit.

Each of these maps h_a, for $a = a_0$, $\overline{a_0}$, a_1 and ± 1, is a *hyperbolic* rational map — an iterate of the map is expanding on the Julia set in the spherical metric — since the simple equivalent condition for hyperbolicity is trivially satisfied. It is clear that each critical point is in the Fatou set, since each critical point is in a superattracting cycle. Each of these maps h_a therefore lies in an open set H_a of rational hyperbolic maps in V_3, which we call the hyperbolic component, such that each maps $h_{a'}$ in h_a is conjugate to h_a on a neighbourhood of the Julia set $J(h_{a'})$ of $h_{a'}$.

A branched covering of the Riemann sphere is *critically finite* if the forward orbit of any critical point is finite. Any hyperbolic component contains at most one critically finite map – always exactly one in V_3 – and so the sets H_a for $a = a_0$, $\overline{a_0}$, a_1 and ± 1 are disjoint. Identifying h_a with a, we can regard H_a as an open subset of the complex plane. It is simply connected in each case. In fact, the uniformising map in each case is a natural parametrisation of the dynamical variation within the hyperbolic component ([**18**], for example). Also, H_a is symmetric about the real axis whenever a is real. The closures of the hyperbolic components $H_{\pm 1}$ meet in three points: 0 (which is in $\mathbb{C}$, but excluded from V_3) and points which we shall call x (with $\mathrm{Im}(x) > 0$) and $\overline{x}$, which are uniquely determined by the property that h_x has a parabolic fixed point with multiplier $e^{2\pi i/3}$. (Of course, it follows immediately that $h_{\overline{x}}$ has a parabolic fixed point with multiplier $e^{-2\pi i/3}$.) The closure of H_{a_1} also includes x and $\overline{x}$. These two points are accessible from $H_{\pm 1}$ [**18**] and H_{a_1}, and 0 is accessible from $H_{\pm 1}$ along the real axis. Since H_{a_1} is unbounded — it includes all points in the negative real axis in $(-\infty, a_1]$, for example — it lies in the unbounded component of the complement of $H_1 \cup H_{-1} \cup H_{a_1} \cup \{x, \overline{x}\}$. We call this unbounded complementary component $V_3(a_1, +)$ [**22**], because it meets the positive real axis. Its closure meets $\overline{H_1} \setminus \{x, \overline{x}\}$ and $\overline{H_{a_1}} \setminus \{x, \overline{x}\}$, but not $\overline{H_{-1}} \setminus \{x, \overline{x}\}$. There are three other complementary components. One is bounded by $H_{-1} \cup H_{a_1} \cup \{x, \overline{x}\}$ and does not meet $\overline{H_1} \setminus \{x, \overline{x}\}$. This complementary component is called $V_3(a_1, -)$ (and meets the negative real axis). The other two complementary components are bounded by $H_{a_0} \cup H_{\overline{a_0}} \cup \{x, \overline{x}\}$. One of these, $V_3(a_0)$, is in the upper half-plane, and contains H_{a_0}. The other, $V_3(\overline{a_0})$, is in the lower half-plane, and contains $H_{\overline{a_0}}$. This is shown in Figure 1.



FIGURE 1. The parameter space V_3

1.3. Counting hyperbolic components

Each of the sets $V_3(a_0)$, $V_3(\overline{a_0})$, $V_3(a_1,-)$ and $V_3(a_1,+)$ contains infinitely many hyperbolic components of each of types III and IV, each with a critically finite centre. For maps in a *type III* hyperbolic component (in the space of quadratic rational maps) there is a single periodic cycle of Fatou components, containing exactly one of the critical values, and the other critical value is in the full orbit of this periodic cycle. For hyperbolic components intersecting V_3, the cycle is of period 3, and the second critical value is in a Fatou component of preperiod $m > 0$. For the critically finite centre h_a, the second critical value $v_2(a)$ has preperiod m. For *type IV* hyperbolic components, there are two distinct cycles of periodic Fatou components, each containing a critical point and value. As in [**22**], we are concerned in this paper only with type III hyperbolic components, and, in particular, with their centres. For each fixed m, the set of type III centres for preperiod m is finite, and has cardinality

$$\frac{23}{21}2^{m+1} + O(1).$$

Moreover, we can identify the number of type III components of preperiod m in each of the sets $V_3(a_0)$, $V_3(\overline{a_0})$, $V_3(a_1,-)$ and $V_3(a_1,+)$. We call denote the set of critically finite centres of type III components by $P_{3,m}$, and the intersections with each of the above sets of V_3 by, respectively, $P_{3,m}(a_0)$, $P_{3,m}(\overline{a_0})$, $P_{3,m}(a_1,+)$ and $P_{3,m}(a_1,-)$. As a consequence of the main result Theorem 2.10 of [**22**], supplemented by the counting in section 3.4 of [**22**], the numbers in each of these sets are, respectively,

$$\frac{3}{7}2^{m+1} + O(1), \quad \frac{3}{7}2^{m+1} + O(1), \quad \frac{1}{7}2^{m+1} + O(1), \quad \frac{2}{21}2^{m+1} + O(1).$$

In fact, we only need the statement of the "easy" parts 1 and 2 of Theorem 2.10 of [**22**], together with the counting, to deduce this. Counting results of this type, although not this particular one, are proved in [**12**].

1.4. Thurston equivalence

For a branched covering f of $\overline{\mathbb{C}}$, we define

$$X(f) = \{f^n(c) \mid c \text{ critical}, \, n > 0\}.$$

We say that f is *critically finite* if $X(f)$ is finite. The convention (not universal) in this paper is to number the critical values. Adopting this convention, two critically finite maps f_0 and f_1 with numbered critical values are *(Thurston) equivalent* if there is a homotopy f_t from f_0 to f_1 such that $\#(X(f_t))$ is constant in t, so that the finite set $X(f_t)$ varies isotopically with t, and the isotopy between $X(f_0)$ and $X(f_1)$ preserves the numbering of critical values. If $\#(X(f_t)) \geq 3$, this is equivalent to the existence of homeomorphisms φ and $\psi : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ with φ and ψ isotopic via an isotopy mapping $X(f_0)$ to $X(f_1)$, and preserving the numbering of critical values, and such that

$$\varphi \circ f_0 \circ \psi^{-1} = f_1.$$

We shall write $\simeq$ for the equivalence relation of Thurston equivalence, and will sometimes write simply "equivalence" or "equivalent" when it is clear that Thurston equivalence is meant.

Thurston's theorem [**5, 21**] gives a necessary and sufficient condition for a critically finite rational map f to be equivalent to a rational map g, which is usually unique up to Möbius conjugation. This is certainly true if the forward orbit of every critical point contains a periodic critical point, which is the only case which concerns us here. (In fact, if it is not true, then $\#(X(f)) = 4$, every critical point is strictly preperiodic, and maps forward to an orbit of period one or two.)

1.5. Capture paths and capture maps

A capture path is a particular type of path in the dynamical plane of a critically periodic quadratic polynomial, or any rational map which is Möbius conjugate to a critically periodic quadratic polynomial, that is, one for which the finite critical point is periodic. In particular, we can define capture paths for the maps h_a for $a = a_0$, $\overline{a_0}$ and a_1. We shall also use capture paths for branched coverings which are Thurston equivalent to quadratic polynomials.

So now let $a = a_0$, $\overline{a_0}$ or a_1. Recalling that 0 is of period 3 under h_a (in fact this is the case for any $h_a \in V_3$), we define

$$Z_m(h_a) = h_a^{-m}(\{0, 1, \infty\}) = h_a^{-m}(\{0, h_a(0), h_a^2(0)\})$$

and

$$Z(h_a) = \cup_{m \geq 0} Z_m(h_a) = \cup_{i \geq 0} h_a^{-i}(\{0, 1, \infty\}).$$



FIGURE 2. σ_β

A *capture path for* h_a is a path $\beta : [0,1] \to \overline{\mathbb{C}}$ from the fixed critical point $\beta(0) = c_2(a) = \frac{2a}{a+1}$ to a point $\beta(1) = x \in Z(h_a)$. To be a capture path, the path has to cross the Julia set $J(h_a)$ just once, into the Fatou component containing x, and this Fatou component must be adjacent to the ray of entry. Two paths β_1 and β_1 are *equivalent* if they are homotopic in $\overline{\mathbb{C}} \setminus \{h_a^n(x) \mid n \geq 0\}$ via a homotopy fixing endpoints. For $a = a_0$ or $\overline{a_0}$, a capture path is uniquely determined up to equivalence, by its endpoint. This is because the Julia set of h_a in these cases is homeomorphic to the Julia set of the rabbit or antirabbit polynomial respectively, and the forward orbit of x is in just one component of the complement in the filled Julia set, of the closure of the immediate basin of attraction. For $a = a_1$, the forward orbit of x sometimes intersects two components of the complement, in the filled Julia set, of the closure of the immediate basin of attraction of x. But the set of x of preperiod m for which there are two such components, is of density tending to 0 as m tends to infinity. We shall be more precise about this below, in 1.7

Any capture path is an arc up to equivalence and from now on we assume that capture paths are arcs. If $\beta : [0,1] \to \overline{\mathbb{C}}$ is an arc then we can define a homeomorphism σ_β as follows. Take a suitably small disc neighbourhood U of β. Define σ_β to be the identity outside U, and to map $\beta(0)$ to $\beta(1)$. See Figure 2. It is possible to define σ_β for any continuous path β, by writing β as a union of arcs, up to homotopy. We shall need to employ this later when β is a closed loop.

A capture path β is *type II* if the second endpoint is periodic, and *type III* if the second endpoint is preperiodic. If β is a type III capture path for h_a, for $a = a_0$, $\overline{a_0}$ or a_1, then $\sigma_\beta \circ h_a$ is a critically finite branched covering. If U is sufficently small to be disjoint from $\{h^n(x) \mid n > 0\}$ then $\sigma_\beta \circ h_a$ is uniquely determined up to Thurston equivalence by the equivalence (homotopy) class of β. Hence, if β is a capture path, $\sigma_\beta \circ h_a$ is determined up to equivalence by the endpoint $\beta(1)$, except for a set of endpoints of density tending to 0 as the preperiod tends to infinity. If β is a capture path then we call $\sigma_\beta \circ h_a$ a *capture* or a *Wittner capture.* This definition of capture was used in [**19, 20, 21, 22, 23**], following the introduction of captures by Ben Wittner in his thesis [**28**], but some authors use "capture" more generally for what we call "type III". We refer to a hyperbolic component whose centre is a capture in our stricter sense, up to Thurston equivalence, as a *capture component*.

If β is a type II capture path for h_a then we can define a critically finite branched covering using β in a slightly different way (see [**19, 20, 21, 22**]). Let ζ be the uniquely determined path such that $h_a \circ \zeta = \beta$ and such that $\zeta(1)$ is periodic under h_a. Then $\sigma_{\zeta}^{-1} \circ \sigma_\beta \circ h_a$ is a critically finite branched covering of type II, which is, again, uniquely determined up to Thurston equivalence by the equivalence class of β.

1.6. Laminations and Lamination maps

Invariant laminations were introduced by Thurston [**26**] to describe the dynamics of polynomials with locally connected Julia sets. The *leaves* of a lamination L are straight line segments in $\{z : |z| \leq 1\}$. Invariance of a lamination means that if there is a leaf with endpoints z_1 and z_2, then there are also a leaf with endpoints z_1^2 and z_2^2, a leaf with endpoints $-z_1$ and $-z_2$, and a leaf with endpoints w_1 and w_2, where $w_1^2 = z_1$ and $w_2^2 = z_2$. A leaf with endpoints $e^{2\pi i a_1}$ and $e^{2\pi i a_2}$, for $0 \leq a_1 < a_2 < 1$, is then said to have *length* $\min(a_2 - a_1, a_1 + 1 - a_2)$. *Gaps* of the lamination are components of $\{z : |z| < 1\} \setminus (\cup L)$. If the longest leaf of L has length $< \frac{1}{2}$ then there are exactly two with the same image, which is called the *minor leaf*. A lamination L is *clean* if finite-sided gaps of L are never adjacent. Minor leaves of clean laminations are either equal or have disjoint interiors. We are only interested, here, in minor leaves which have endpoints which are periodic under $z \mapsto z^2$. If one endpoint of a minor leaf of a clean lamination is periodic, then the other is too, and of the same period. Any two such minor leaves have distinct endpoints. There is a natural partial ordering on minor leaves. We say that $\mu < \mu'$ if μ separates μ' from 0 in the unit disc. For any minor leaf μ there is a unique minimal minor leaf $\nu = \nu(\mu)$ such that $\nu \leq \mu'$ whenever $\mu' \leq \mu$.

Each point $e^{2\pi i t_1} \neq 1$, with t_1 an odd denominator rational, is an endpoint of the minor leaf μ_{t_1} of a unique clean lamination, which we call L_{t_1} – and also L_{t_2}, if $e^{2\pi i t_2}$ is the other endpoint of μ_{t_1} (in which case $\mu_{t_1} = \mu_{t_2}$). We can define a *lamination map* $s_{t_1} = s_{t_2}$ which maps L_{t_1} to L_{t_1}, and such that $s_{t_1}(z) = z^2$ for $|z| \geq 1$, and gaps are mapped to gaps. The gap of L_{t_1} containing 0 has infinitely many sides and is periodic under s_{t_1}, of period n, and is mapped with degree two onto its image by s, but the rest of the periodic cycle maps homeomorphically. We can therefore choose s_{t_1} so that 0 is a degree two critical point, of period n, and hence is a degree two critically periodic branched covering. Then s_{t_1} is Thurston equivalent to a unique critically finite quadratic polynomial, and conversely.

For technical reasons, in [**22**], it turned out to be easier to work with the lamination map which is Thurston equivalent to h_a, rather than with h_a itself. This will be true here, too. The corresponding lamination map is s_q for $q = \frac{1}{7}$ or $\frac{6}{7}$ or $\frac{3}{7}$ respectively. Capture paths transfer to this setting. A *capture path for* s_q is a path from the fixed critical point ∞ of s_q to a point x in the full orbit $Z(s_q) = \cup_{n>0} s_q^{-n}(\{0, s_q(0), s_q^2(0)\})$ of the critical point 0, which crosses the unit circle just once, into the infinite-sided gap of L_q which contains x. The associated type III and type II capture maps are defined in exact analogy to those in 1.5.

1.7. Counting capture maps and components

The following information is given in [**22**], with a little more detail in places, but for ease of reference, it is given here. A theorem of Tan Lei [**25**], of which a more general version is proved in [**19, 20**], gives a necessary and sufficient criterion for a capture $\sigma_\gamma \circ s_q$ (for any odd denominator rational q) to be Thurston equivalent to a rational map. If $q = \frac{r}{7}$ for $1 \leq r \leq 6$, then this rational map is automatically in V_3. Actually, Tan Lei's theorem deals with matings rather than captures, but the result for captures can be proved by the same methods. The criterion is easily described using minor leaves, and therefore we use the captures $\sigma_\gamma \circ s_q$. The critically finite branched covering $\sigma_\gamma \circ s_q$ is Thurston equivalent to a rational map if and only if $\gamma(1)$ is in the larger region of the unit disc bounded by the minimal minor leaf $\nu(\mu_q)$ with $\nu(\mu_q) \leq \mu_q$. The minimal minor leaves $\nu(\mu_q)$ for $q = \frac{1}{7}$, $\frac{6}{7}$ and $\frac{3}{7}$ are $\mu_{1/7}$ (which has endpoints $e^{2\pi i(1/7)}$ and $e^{2\pi i(2/7)}$), $\mu_{6/7}$ (which has endpoints $e^{2\pi i(5/7)}$ and $e^{2\pi i(6/7)}$) and $\mu_{1/3}$ (which has endpoints $e^{\pm 2\pi i(1/3)}$) respectively.

We write $\mathcal{Z}(q)$ for the set of equivalence classes of capture paths γ for s_q such that $\sigma_\gamma \circ s_q$ is equivalent to a rational map, and $\mathcal{Z}_m(q)$ for the subset for which the second endpoint of the path is of preperiod m. The number $\#(\mathcal{Z}_m(q))$ is closely related to the the number of points of $Z_m(s_q) = s_q^{-m}(\{s_q^i(0) \mid i \geq 0\})$ in the larger region of the unit disc bounded by $\nu(\mu_q)$. The numbers of such points for preperiod m are, respectively,

$$\frac{3}{7}2^{m+1} + O(1), \quad \frac{3}{7}2^{m+1} + O(1), \quad \frac{1}{3}2^{m+1} + O(1).$$

For $q = \frac{1}{7}$ and $\frac{6}{7}$, it is not hard to show that capture paths γ_1 and γ_2 for s_q are equivalent if and only if the captures $\sigma_{\gamma_1} \circ s_q$ and $\sigma_{\gamma_2} \circ s_q$ are Thurston equivalent. Also, it was shown in [**22**] that the equivalent rational maps to captures $\sigma_\gamma \circ s_{1/7}$ are precisely the centres of hyperbolic components in $V_3(a_0)$, and similarly for captures $\sigma_\gamma\circ s_{6/7}$ and centres of hyperbolic components in $V_3(\overline{a_0})$. (It was pointed out in [**22**] that this can be shown by elementary methods.)

For $q=\frac{3}{7}$, the number of equivalence classes of capture paths is slightly more than the number of endpoints, because , for some choices of endpoint x, there are two equivalence classes of capture paths with endpoint x. But even for $q=\frac{3}{7}$, there is just one equivalence class of capture path with endpoint x, unless the closure of the gap of $L_{3/7}$ which contains x intersects both $\{e^{2\pi it}:t\in(\frac{2}{7},\frac{1}{3})\}$ and $\{e^{2\pi it}:t\in(\frac{2}{3},\frac{6}{7})\}$. In that case, there are exactly two equivalence classes of capture paths for $s_{3/7}$ ending at x: those which intersect $\{e^{2\pi it}\mid t\in(\frac{2}{7},\frac{1}{3}\}$ and those which intersect $\{e^{2\pi it}:t\in(\frac{2}{3},\frac{6}{7})\}$. There may then be either one or two Thurston equivalence classes of captures $\sigma_\gamma\circ s_{3/7}$ for γ with endpoint at x. Examples of both cases are given in [**22**]. But at any rate, the number of maps in V_3 which are equivalent to capture maps $\sigma_\gamma\circ s_{3/7}$ is

$$\frac{1}{3}2^{m+1}(1+o(1)).$$

However, there are Thurston equivalences between some captures $\sigma_\gamma\circ s_{3/7}$ and captures $\sigma_\zeta\circ s_q$ for $q=\frac{1}{7}$ or $\frac{6}{7}$. These arise from the so-called *Wittner flip construction.* If γ crosses the unit circle at $e^{2\pi it}$ for $t\in(\frac{1}{7},\frac{2}{7})$, then $\sigma_\gamma\circ s_{3/7}$ is equivalent to a capture $\sigma_\zeta\circ s_{6/7}$, and similarly, if $t\in(\frac{5}{7},\frac{2}{7})$ then $\sigma_\gamma\circ s_{3/7}$ is equivalent to a capture $\sigma_\zeta\circ s_{1/7}$. This very interesting construction is not used in the results of this paper, so we will not discuss it further.

Capture paths γ_1 and γ_2 which cross the unit circle in $\{e^{2\pi it}:t\in(-\frac{1}{7},\frac{1}{7})\}$ are equivalent if and only if the captures $\sigma_{\gamma_1}\circ s_q$ and $\sigma_{\gamma_2}\circ s_p$ are Thurston equivalent. It was shown in [**22**] that the rational maps equivalent to these captures are precisely all the centres of the hyperbolic components in $V_3(a_1,-)$. Ideas of Timorin (private communication) indicate that there is a simple direct way of proving this.

Thus, using Thurston equivalence, there are natural one-to-one correspondences between the hyperbolic components in the regions $V_3(a_0)$, $V_3(\overline{a_0})$ and $V_3(a_1,-)$ and equivalence classes of captures by s_q for $q=\frac{1}{7}$, $\frac{6}{7}$ and $\frac{3}{7}$ respectively, where the capture paths are taken to have endpoints in $\cup_{n>0}s_q^{-n}(0)$, and in, respectively:

- the larger region of the unit disc bounded by the leaf $\mu_{1/7}$;
- the larger region of the unit disc bounded by the leaf $\mu_{6/7}$;
- the smaller region of the unit disc bounded by the leaf with endpoints $e^{\pm 2\pi i(1/7)}$.

But none of this is true for $V_3(a_1,+)$. The only captures which can be equivalent to centres of hyperbolic components in this region are those of the form $\sigma_\gamma \circ s_{3/7}$ where γ crosses the unit circle at $e^{2\pi it}$ for $t \in (\frac{2}{7}, \frac{1}{3}) \cup (\frac{2}{3}, \frac{6}{7})$, since as we have seen all the other captures are equivalent to centres in the other regions $v_3(a_0)$ or $V_3(\overline{a_0})$ or $V_3(a_1, -)$. The number of preperiod m endpoints of capture paths in this region is

$$\frac{1}{21} \cdot 2^{m+1} + O(1).$$

It was shown in [**23**] that for any integer $n > 0$ there are n inequivalent capture paths – in fact, with distinct endpoints – all crossing the unit circle at points in $\{e^{2\pi it} : t \in (\frac{2}{7}, \frac{17}{56})\}$. There are plenty of capture components in $V_3(a_1, +)$, but not every type III hyperbolic component in $V_3(a_1, +)$ is a capture component, since as we have seen in 1.3 there are at least twice as many type III hyperbolic components as can be represented. In fact because of the multiple equivalences it is somewhat less. In fact, the main results of this paper imply that the number of type III hyperbolic components in $V_3(a_1, +)$ which are represented by captures is between $c_1 2^{m+1}$ and $c_2 2^{m+1}$ where $0 < c_1 < c_2 < \frac{1}{21}$.

We now introduce some more notation. We write $\mathcal{Z}_m(3/7, +)$ for the subset of $\mathcal{Z}_m(3/7)$ such that the intersection of the paths with the unit circle is in

$$\left\{e^{2\pi it} : t \in \left(\frac{2}{7}, \frac{1}{3}\right) \cup \left(\frac{2}{3}, \frac{5}{7}\right)\right\},$$

and $\mathcal{Z}_m(3/7, -)$ for the subset of $\mathcal{Z}_m(\frac{3}{7})$ such that the intersection with the unit circle is in

$$\left\{e^{2\pi it} : t \in \left(-\frac{1}{7}, \frac{1}{7}\right)\right\}.$$

We write $\mathcal{Z}_m(3/7, +, +)$ for the subset of $\mathcal{Z}_m(3/7, +)$ such that the intersection is with $\{e^{2\pi it} \mid t \in (\frac{2}{7}, \frac{1}{3})\}$. We also write

$$\mathcal{Z}(3/7, +) = \cup_{m>0} \mathcal{Z}_m(3/7, +), \quad \mathcal{Z}(3/7, -) = \cup_{m>0} \mathcal{Z}_m(3/7, -),$$

$$\mathcal{Z}(3/7, +, +) = \cup_{m>0} \mathcal{Z}_m(3/7, +, +).$$

Similarly we write $\mathcal{Z}_m(3/7, +, p)$ for subset of $\mathcal{Z}_m(3/7, +)$, such that the intersection with the unit circle is in

$$\left\{e^{2\pi it} : t \in \left(\frac{1}{3} - 2^{-p}\frac{1}{21}, \frac{1}{3} - 2^{-(p+1)}\frac{1}{21}\right) \cup \left(\frac{2}{3} - 2^{-(p+1)}\frac{1}{21}, \frac{2}{3} - 2^{-p}\frac{1}{21}\right)\right\}.$$

We define $\mathcal{Z}(3/7, +, p)$, and $\mathcal{Z}(3/7, +, +, p)$, analogously to the above.

The *parameter capture map* Φ (for V_3) is defined by

$$\Phi([\gamma]) = [\sigma_\gamma \circ s_q],$$

if γ is a capture path for s_q, for $q = \frac{1}{7}$, $\frac{6}{7}$ or $\frac{3}{7}$, and where $[\gamma]$ is the equivalence class of any capture path γ for s_q such that $\sigma_\gamma \circ s_q$ is equivalent to a rational map, and $[\sigma_\gamma \circ s_q]$ denotes the Thurston equivalence class of the critically finite branched covering $\sigma_\gamma \circ s_q$. We shall write Φ_m for the parameter capture map restricted to $([\gamma], s_q)$ for which the second endpoint of γ is of preperiod m under s_q. As we have seen, the parameter capture map restricted to pairs $([\gamma], s_{1/7})$ or $([\gamma], s_{6/7})$ is a bijection onto a naturally defined domain, and similarly restricted to pairs $([\gamma], s_{3/7})$ with $[\gamma] \in \mathcal{Z}(\frac{3}{7}, -)$. The case of $([\gamma], s_{3/7})$ for $[\gamma] \in \mathcal{Z}(3/7, +)$ is quite different. We will usually identify to domain of Φ in this case with $\mathcal{Z}(3/7, +)$. We shall also write Φ_m for the capture map Φ restricted to $\mathcal{Z}_m(3/7, +)$.

We can now state our main theorems.

Theorem 1.8. *For any capture path γ in $\mathcal{Z}(3/7, +, +, 0)$, the capture $\sigma_\gamma \circ s_{3/7}$ is equivalent to the centre of a type III hyperbolic component in $P_{3,m}(a_1, +) \subset V_3(a_1, +)$. Thus,*

$$\Phi(\mathcal{Z}(3/7, +, +, 0)) \subset P_{3,m}(a_1, +).$$

Theorem 1.9. *There is a constant $c > 0$ such that the number of type III hyperbolic components in $V_3(a_1, +)$ with second critical point in an immediate basin of preperiod m and represented by $\sigma_{\gamma_i} \circ s_{3/7}$ for exactly 2^r different capture paths γ_i in $\mathcal{Z}_m(3/7, +, +, 0)$, for $r = 1$ or any integer $r \geq 0$, is $\geq \lfloor 2^{-cr+m} \rfloor$. Hence, for $c_r = 2^{-cr}$,*

$$\#(\{[\gamma] \in \mathcal{Z}_m(3/7, +, +, 0) : \#(\Phi_m^{-1}([\sigma_\gamma \circ s_{3/7}]) = 2^r\}) \geq c_r.2^m.$$

1.10. So what?

Why does it matter? One of the most basic ideas in dynamics, especially in complex dynamics, is to model parameter space, at least local dynamics, by the parameter plane. The most famous example in complex dynamics is the combinatorial model for the Mandelbrot set for quadratic polynomials, which is based, perhaps surprisingly given the lack of immediate visual resemblance, on the dynamical plane of $z \mapsto z^2$. This results from the seminal work of Douady and Hubbard [**4**] which was reinterpreted by Thurston [**26**]. Thurston's framework is used in this article. The information obtained about the relative positions of hyperbolic components, and the dynamics in each hyperbolic component, just from the dynamics of one map, is extraordinarily detailed, and is complete. A conjectural topological model for the parameter space of quadratic polynomials is thus obtained. Important advances in proving that the model is the correct one were initially obtained by Yoccoz (described in [**6, 24**]). Nearly thirty years after the model was first obtained, it remains a conjectural model, although a great deal has been proved about it (for example, [**13**]) and important advances have been made recently [**11, 2, 7, 8, 10, 9**]. The topological equivalence of the combinatorial model to the actual parameter space is equivalent to the conjecture that the Mandelbrot set is locally connected (MLC).

Given that the information about combinatorial structure of quadratic polynomial parameter space is so detailed, it is natural to attempt a model for parameter space of quadratic rational maps in terms of the one for polynomials. This is the route that has been followed. It would also be natural to try to model parts of the parameter space on the dynamical planes of polynomials within the parameter space by simply using captures. Restricting to the subset V_3 of rational parameter space, this means using the dynamical planes of h_a for $a = a_0$, $\overline{a_0}$ and a_1, or, equivalently, the dynamical plane s_q for $q = \frac{1}{7}$, $\frac{6}{7}$ and $\frac{3}{7}$. It has been clear for some time that it is possible to do this in some parts of parameter space and essentially impossible in others. If we fix attention on V_3, then parts of the Julia sets of h_a for $a = a_0$, $\overline{a_0}$ and a_1, are visible in $V_3(a_0)$, $V_3(\overline{a_0})$ and $V_3(a_1, -)$. It can be proved — following the lines of analogous results of Aspenberger-Yampolsky [**1**] and of Timorin [**27**] for V_2 — that parts of these Julia sets translate into subsets of $V_3(a_0)$, $V_3(\overline{a_0})$. There is hope of proving something along these lines for $V_3(a_1, -)$. But the situation for capture paths in $\mathcal{Z}(a_1, +)$ is quite different. Theorem 1.8 shows that the image is contained in $V(a_1, +)$. The map cannot be surjective onto type III centres in $V_3(a_1, +)$, because there are approximately twice as many centres of preperiod m as there are capture paths in $\mathcal{Z}_m(3/7, +)$. It follows from the results of [**23**] that the map cannot be injective. But Theorem 1.9 quantifies this. The way in which the proof proceeds also suggests that the capture map from $\mathcal{Z}(3/7, +)$ to $V_3(3/7, +)$ does not extend continuously to a larger set. So a transfer of topological information from dynamical plane to parameter plane is, at least, problematic. But the proof of 1.9 does indicates ways in which discontinuities might arise.

1.11. The parameter capture map is not typical

It is interesting to compare the parameter capture maps Φ_m on $\mathcal{Z}_m(3/7, +)$ with typical maps between finite sets. From 1.7 we see that the size of $\mathcal{Z}_m(3/7, +)$ is $\frac{1}{21}2^{m+1} + O(1)$, while from 1.3, the size of the range $P_{3,m}(a_1, +)$ is $\frac{2}{21}2^{m+1} + O(1)$, that is, twice the size of the domain, to within $O(1)$. So the ratio of the domain size to range size for Φ_m tends to $\frac{1}{2}$ as m tends to ∞. Typical behaviour of gaps between finite sets is a classical problem in probability, often described in terms of "balls in boxes", and involving Stirling numbers of the second kind. Saddle point methods are usually used in the solution. It is shown [**3**] that, with probability tending to 1 as N tends to ∞ for maps from a set of M elements to a set of N elements, and for $M = bN$ with $b > 0$, the number of points with r preimages is within $O(\sqrt{N})$ of

$$\frac{b^r}{r!}N,$$

and the image size is within $O(\sqrt{N})$ of $(1 - e^{-b})N$. In particular, the maximum inverse image size for a typical map, for b bounded above and below, is $\log N / \log\log N(1+o(1))$. Surprisingly, the maximum inverse image size of Φ_m is a bit larger than typical, at least $cm = c_1 \log(2^m)$. This is especially intriguing as we shall see that

$$\Phi_m \mid \mathcal{Z}_m(s_{3/7}, +, +, 0)$$

can be written as a composition of a sequence of maps which converge exponentially fast to the identity, in a reasonable sense. Therefore one might expect Φ_m to behave more like a bijection than a typical map. In many respects, this could well be the case. The question of the relation of the image size to domain size is quite open. In principle, this could be achieved by estimation, but the practicality is another matter.

1.12. Contrast with the subhyperbolic case

All the maps considered in this work are hyperbolic critically finite rational maps. Any critically finite rational map is either hyperbolic or *subhyperbolic*. Subhyperbolic maps are expanding on their Julia sets, with respect to a suitable adaptation of the spherical metric: Lipschitz equivalent to the spherical metric outside the postcritical set. A critically finite map is hyperbolic if and only if the forward orbit of every critical point contains a periodic critical point. The concept of mating or capture can be extended to critically finite maps which are subhyperbolic and not hyperbolic. In the non-hyperbolic case it might not be useful to distinguish between matings and captures, and the term mating is often used. Some very interesting studies of these have been made in the non-hyperbolic case, by Daniel Meyer [**14, 16, 15**]. It seems that the non-hyperbolic case is very different from the hyperbolic case. In particular, the number of different representations of a map by matings is much larger — or almost certainly so. No definitive upper bound on the number of equivalent maps captures is given in the hyperbolic case, in the current work.

1.13. Organisation of the paper

The organisation of this paper is as follows.

Chapter 2 concerns the Resident's View. The "Resident's View" is the term used here, and in earlier papers on the same general subject, for a particular manifestation of a very basic philosophy in dynamical systems.

The idea is that information about variation of dynamics in a parameter space can be obtained by scrutiny of a fixed dynamical system within the parameter space. It is more appropriate, here, to say "resident's views" since the focus of the study is on determining when different models – or different "views" – represent the same dynamical system. The representations that we are considering are all Wittner captures. Chapter 2 reinterprets Theorem 1.8 into the context of the Resident's View. The resulting statement, Theorem 2.8, thus becomes one of the main goals of the paper. The proofs of both Theorems 1.8 and 1.9 then use Theorem 2.8 and its context.

Theorem 2.8 is statement about details of a certain group action on paths in the dynamical plane of a quadratic polynomial, a group action which has been the subject of intense study, notably in [**22**], where a fundamental domain was found for the group action in a particular case. This is the main tool in the current work. Theorem 2.8 gives important information about successive translates of the fundamental domain which are crossed by "parameter capture paths". Chapter 3 is devoted to providing the foundations for proving Theorem 2.8. The group action is a twisted group action, where the twisted action comes from certain homemorphisms which can be regarded as conjugating homeomorphisms between different model dynamical spaces. Chapter 3 is essentially devoted to the study of these homeomorphisms.

Chapter 4 contains the proof of Theorem 2.8. In order to prove this theorem, and in preparation for the proof of Theorem 1.9, more detailed information about the translates of the fundamental domain are given. Theorem 2.8 implies that translates of the fundamental domain are paired off, each one with an adjacent translate. The results at the end of Chapter 4, in particular 4.10.2, give good estimates on the distance between the translates, that is, on the distance between the group elements by which translation is made. These estimates are vitally important in controlling the effect of the twisted group action, and hence for proving the theorems 1.8 and 1.9.

Chapter 5 presents key examples. The positive density sets which are the subject of Theorem 1.9 are then perturbations of these key examples. The main result is Theorem 5.2. This result says, surprisingly, that for some constant $C > 0$, and each integer $k > 0$ there is a type III quadratic rational map with one critical point of period 3 and the other of preperiod $\leq C2^k$ which can be represented by exactly 2^k different Wittner captures with capture paths in $\mathcal{Z}(3/7, +, +, 0)$. The proof is direct and does not need information about the sides of translates of the fundamental domain that are crossed by these capture paths. Nevertheless, considerable information is given about these, in preparation for the positive density results in Chapter 6, with the proof of theorem 1.9.

Chapter 6 gives the proof of the positive density results, in particular, 1.9. It might be interesting to compare the implementation of the technique with the now-classical results of positive measure sets of interval maps with characteristics of non-uniform hyperbolicity. In all theorems of this type, all estimates depend on some sort of a priori control. It never seems possible to avoid a chicken-and-egg argument, and to prove results without any sort of control. This means that, although results are obtained, concerning the abundance of certain phenomena, no absolute upper bound is achieved, and there are number of open questions.

CHAPTER 2

The resident's view of the main theorems

2.1. A notation for Thurston equivalence

The following notation was used in [**19, 20, 21, 22, 23**]. If critically finite branched coverings f and g are Thurston equivalent, then there is an orientation-preserving homeomorphism $\psi : \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ such that $\psi(X(f_0)) = X(f_1)$, and ψ preserves the numbering of critical values and there is a homotopy g_t from $\psi \circ f \circ \psi^{-1} = g_0$ and g_1 such that g_t is a critically finite branched covering for all t with $X(g_t) = X(g)$ for all t. The definition of Thurston equivalence is sometimes stated in this way. If such a ψ exists, then we write $f \simeq_\psi g$. The drawback of this notation is that $\simeq_\psi$ is not an equivalence relation. However, if $f \simeq_\psi g$ then $g \simeq_{\psi^{-1}} f$, and if also $g \simeq_\xi h$ then $f \simeq_{\psi\circ\xi} h$.

An extension of this notation has also been used, and will be used in this paper. Suppose that $X_1(f)$ is a finite set which contains $X(f)$ and such that $f(X_1(f)) \subset X_1(f)$. Suppose that $X_1(g)$ has similar properties relative to g. Then $(f, X_1(f))$ is said to be *Thurston equivalent to* $(g, X_1(g))$, written $(f, X_1(f)) \simeq (g, X_1(g))$ if there is a homotopy through $(f_t, X_1(f_t))$ such that:

- $(f_0, X_1(f_0)) = (f, X_1(f))$ and $(f_1, X_1(g)) = g$;
- f_t is a critically finite branched covering with $X(f_t) \subset X_1(f_t)$ and $f_t(X_1(f_t)) \subset X_1(f_t)$;
- $X_1(f_t)$ moves isotopically with t and numbering of critical points is preserved.

Similarly to the above, if $(f, X_1(f)) \simeq (g, X_1(g))$, then ψ and g_t exist as before, and in addition, $\psi(X_1(f)) = X_1(g)$ and $g_t(X_1(g)) \subset X_1(g)$ for all t. We then write $(f, X_1(f)) \simeq_\psi (g, X_1(g))$.

A fact which is used throughout [**19, 20, 21, 22, 23**] and which will be used here also, is that if $(f, X_1(f)) \simeq_{\psi_0} (g, X_1(g))$, then for each $n \geq 1$ there is ψ_n such that $(f, f^{-n}X_1(f)) \simeq_{\psi_n} (g, g^{-n}(X_1(g)))$, and such that ψ_n and ψ_k are isotopic via an isotopy which is constant on $f^{-k}(X_1(f))$, for each $0 \leq k \leq n$.

2.2. The boundary of the universal cover

The theory of the Resident's View uses the following facts and notation concerning boundaries of universal covers of Riemann surfaces. Suppose that $V = \overline{\mathbb{C}} \setminus P$ where $P \subset \overline{\mathbb{C}}$ is finite and $\#(P) \geq 3$, so that V has negative Euler characteristic. (This is the set-up in all our examples, but $\overline{\mathbb{C}}$ could be replaced by any compact Riemann surface S, provided that $S \setminus P$ has negative Euler characeristic). The holomorphic universal cover of V is the open unit disc D. For fixed $a \in V$, we define $\pi_1(V, P, a)$ to be the set of homotopy classes of paths $\beta : [0, 1] \to \overline{\mathbb{C}}$ such that $\beta([0, 1)) \subset V$, $\beta(0) = a$ and $\beta(1) \in P$, where we restrict to homotopies through such paths. Then for any $[\beta] \in \pi_1(V, P, a)$, and any lift $\tilde{\beta}$ of $\beta \mid [0, 1)$ to D, $\lim_{t \to 1} \tilde{\beta}(t)$ exists as in D, and is uniquely determined by $[\beta]$ and $\tilde{\beta}(0)$. Therefore, having fixed a lift $\tilde{a}$ of a to D, we can regard $\pi_1(V, a, P)$ as a subset of ∂D, by identifying $[\beta]$ with $\tilde{\beta}(1)$ for any lift $\tilde{\beta}$ with $\tilde{\beta}(0) = \tilde{a}$. This subset is dense.

Since D is holomorphic, the action on D of the covering group of V is by Möbius transformations, and hence this action extends continuously to ∂D. The covering group is naturally isomorphic to the fundamental group $\pi_1(V, a)$, which acts naturally on $\pi_1(V, P, a)$ by concatenation, that is, for $[\gamma] \in \pi_1(V, a)$ and $[\beta] \in \pi_1(V, P, a)$, we define $[\gamma].[\beta] = [\gamma * \beta]$. This coincides with the action by Möbius transformations on ∂D, using the identification of $\pi_1(V, P, a)$ with a subset of ∂D.

2.3. Action of mapping class groups

If $Y \subset \overline{\mathbb{C}}$ is finite, then we denote by $\mathrm{MG}(\overline{\mathbb{C}}, Y)$ the *mapping class group* of $(\overline{\mathbb{C}}, Y)$, that is, the group of isotopy classes of orientation preserving homeomorphisms of $\overline{\mathbb{C}}$ which leave Y invariant, modulo isotopies which are constant on Y. If $Z \subset Y$ then there is a natural homomorphism from $\mathrm{MG}(\overline{\mathbb{C}}, Y)$ to $\mathrm{MG}(\overline{\mathbb{C}}, Z)$, given by mapping the isotopy class of ψ in $\mathrm{MG}(\overline{\mathbb{C}}, Y)$ to the isotopy class of ψ in $\mathrm{MG}(\overline{\mathbb{C}}, Z)$. If $\#(Z) \geq 3$, any lift to the universal cover D of any homeomorphism of $\overline{\mathbb{C}}$ which preserves Z extends homeomorphically to $D \cup \partial D$. If $y \in Y \setminus Z$ and $\widetilde{y}$ is a fixed lift of y to D, and $[\psi] \in \mathrm{MG}(\overline{\mathbb{C}}, Y)$ with $\psi(y) = y$ and $\widetilde{\psi}$ is a lift with $\widetilde{\psi}(\widetilde{y}) = \widetilde{y}$, then $\widetilde{\psi} \mid \partial D$ is uniquely detemined by $\widetilde{y}$ and $[\psi] \in \mathrm{MG}(\overline{\mathbb{C}}, Y)$.

2.4. The Resident's View.

The Resident's View gives information about one punctured sphere in terms of another. As in 1.3, we define $P_{3,m}$ to be the set of centres of hyperbolic components of types II, and of type III of preperiod $\leq m$. We also define

$$V_{3,m} = V_3 \setminus P_{3,m} = \overline{\mathbb{C}} \setminus (\{0, \infty\} \cup P_{3,m}).$$

As we saw in 1.2, there are just two type II hyperbolic components, with centres 1 and -1. So $V_{3,0} = \mathbb{C} \setminus \{0, \pm 1\}$, and, for every $m \geq 0$, $V_{3,m}$ is the complement in the Riemann sphere of four or more punctures. Hence it is certainly a surface of negative Euler characteristic, and the holomorphic universal cover identifies with the open unit disc D. Therefore, as above, $\pi_1(V_{3,m}, P_{3,m}, a)$ identifies with a subset of ∂D for a fixed choice $\tilde{a} \in D$ of lift of a, and $\pi_1(V_{3,m}, a)$ acts on $D \cup \partial D$ by Möbius transformations, where this action coincides with the usual action on $\pi_1(V_{3,m}, P_{3,m}, a) \subset \partial D$ by contenation of paths.

The other punctured sphere, with which we want to compare the first, is a punctured dynamical plane. For $a = a_0$ or $\overline{a_0}$ or a_1, we write

$$Z_m(h_a) = h_a^{-m}(\{0, \infty, 1\}) = h_a^{-m}(\{0, h_a(0), h_a^2(0)\}),$$

and

$$Y_m(h_a) = Z_m(h_a) \cup \{v_2(a)\},$$

where $v_2(a) = \frac{2a}{a+1}$ is the second critical point of h_a, which is fixed by h_a for these values of a and hence is also the second critical value. Thus, $Y_0(h_a)$ is the postcritical set $X(h_a)$ and $Y_m(h_a) = h_a^{-m}(Y_0(h_a))$ for all $m \geq 0$, for these values of a. Similarly (and as in 1.7), for $q = \frac{1}{7}$ or $\frac{6}{7}$ or $\frac{3}{7}$, we write

$$Z_m(s_q) = s^{-m}(\{s_q^n(0) : n \geq 0\},$$

and

$$Y_m(s_q) = Z_m(s_q) \cup \{\infty\},$$

remembering that ∞ is the fixed critical point of s_q. Since $\overline{\mathbb{C}} \setminus Z_m$, for $Z_m = Z_m(h_a)$ or $Z_m(s_q)$ for any of these values of a or q, the holomorphic universal cover is D once again, and we can again regard $\pi_1(\overline{\mathbb{C}} \setminus Z_m(h_a), Z_m(h_a), v_2(a))$ (or $\pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), Z_m(s_q), \infty)$) as a subset of ∂D. Under our definition of Thurston equivalence, the Thurston equivalence class of (f, Y) for any critically finite branched covering f with $\#(Y) \geq 3$ is simply connected. There is therefore a natural isomorphism between $\pi_1(\overline{\mathbb{C}} \setminus Z_m(h_a), v_2(a))$ and $\pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), \infty)$, and a natural identification between $\pi_1(\overline{\mathbb{C}} \setminus Z_m(h_a), Z_m(h_a), v_2(a))$ and $\pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), Z_m(s_q), \infty)$ for $(a, q) = (a_0, \frac{1}{7})$ or $(\overline{a_0}, \frac{6}{7})$ or $(a_1, \frac{3}{7})$.

From now on we fix lifts $\widetilde{a}$ and $\widetilde{\infty}$ to the universal cover D, of $a \in V_{3,m}$ and $\infty = v_2(s_q) \in \overline{\mathbb{C}} \setminus Z_m(s_q)$, in order to define the identifications of $\pi_1(V_{3,m}, P_{3,m}, a)$ and $\pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), Z_m(s_q), \infty)$ with subsets of ∂D.

Given these identifications, Resident's View gives two maps, both of which are called ρ, or $\rho(., s_q)$ if more precision is needed, in [**22**] (and called ρ_2 and ρ in [**21**]):

$$\rho : \pi_1(V_{3,m}, P_{3,m}, a) \to \pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), Z_m(s_q), \infty),$$

$$\rho : \pi_1(V_{3,m}, a) \to \pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), \infty).$$

Both these maps ρ are injective. The first map ρ is also monotone from the dense subset $\pi_1(V_{3,m}, P_{3,m}, a)$ of ∂D into $\pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), Z_m(s_q), \infty) \subset \partial D$. The image is not dense, except in the case when $m = 0$, and hence the map ρ cannot extend continuously to ∂D. However, it does extend continuously except at countably many points, where right and left limits exist. We define D' to be the convex hull of the closure of the image of ρ in ∂D. The convex hull can be taken with respect to either the hyperbolic or Euclidean metric on D, since the two convex hulls are homeomorphic. There is a slight extension of ρ to the domain $\pi_1(V_{3,m}, P_{3,m} \cup \{0, \infty\}, a)$ such that the images of $[\beta]$ is one of countably many points of $\partial D \cap \partial D'$ if β ends at 0, and is the closure of one of the countably many components of $\partial D' \cap D$ if β ends at ∞. In the latter case, we identify $\rho([\beta])$ with the set of paths from $\widetilde{\infty} = \widetilde{\rho(\beta)}(0)$ to $\overline{\partial D' \cap D}$, modulo homotopy equivalence fixing the first endpoint and keeping the second endpoint in $\overline{\partial D' \cap D}$.

The set $\rho(\pi_1(V_{3,m}, P_{3,m}, a))$ is the set of all paths in $\pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), Z_m(s_q), \infty)$ which lift to paths from $\widetilde{\infty}$ to $\partial D \cap \partial D'$. In particular, it includes all paths in $\mathcal{Z}_m(q)$.

The second map ρ, with domain $\pi_1(V_{3,m}, a)$, is *not* a group homomorphism. Nevertheless, the action of $\pi_1(V_{3,m}, a)$ on ∂D translates to an action on $\partial D'$, and, indeed on ∂D, in the following way. Suppose that $\gamma \in \pi_1(V_{3,m}, a)$ and that $\rho(\gamma) = \alpha$. Then there is $[\psi] \in \mathrm{MG}(\overline{\mathbb{C}}, Y_{m+1}(s_q))$ such that

$$s_q \simeq_\psi \sigma_\alpha \circ s_q$$

and for all $\beta \in \pi_1(V_{3,m}, P_{3,m}, a)$,

$$\rho([\gamma * \beta]) = [\alpha * \psi(\rho(\beta))]$$

where $\rho(\beta)$ is any path representing $\rho([\beta])$. Moreover, $[\psi]$ is uniquely determined and the map

$$\Xi : [\gamma] \mapsto [\psi]$$

is a group isomorphism of $\pi_1(V_{3,m}, a)$ onto a subgroup G of

$$\{[\psi] \in \mathrm{MG}(\overline{\mathbb{C}}, Y_{m+1}(s_q)) : \ \psi(\infty) = \infty, \ \exists \alpha \in \pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), \infty), \ s_q \simeq \sigma_\alpha \circ s_q\}.$$

(The notation Ξ has not been used in the earlier papers being quoted, where, for reasons which we will not go into, an anti-isomorphism, rather than an isomorphism, was used.) Here, $[\psi]$ determines $[\alpha]$ uniquely, and vice versa. For $[\psi] \in G$ and α as above, we define $\widetilde{\psi}$ to be the lift of ψ which maps $\widetilde{\infty}$ to $\widetilde{\alpha}(1)$, where the path α has domain $[0, 1]$ and the lift $\widetilde{\alpha}$ of α is such that $\widetilde{\alpha}(0) = \widetilde{\infty}$. Then $[\psi] \to \widetilde{\psi}|\partial D$ is an isomorphism onto its image, and therefore an action of G on ∂D is defined by

$$[\psi].x = \widetilde{\psi}(x) \quad (x \in \partial D).$$

If $x = [\zeta] \in \pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), Z_m(s_q), \infty)$, then $\widetilde{\psi}([\zeta]) = [\alpha * \psi(\zeta)]$. The action of $\pi_1(V_{3,m}, a)$ on ∂D therefore transfers under ρ to the restriction (to $\partial D' \cap \partial D$) of a natural action of G on ∂D.

2.5. Key features of the fundamental domain

The fundamental domain for $V_{3,m}$ that was chosen in [**22**] is an ideal polygon $F_{3,m}$ in the universal covering space of $V_{3,m}$, that is, a region of D that is bounded by finitely many geodesics (in the hyperbolic metric), with closure intersecting ∂D in just finitely many points, which are all in $\pi_1(V_{3,m}, P_{3,m} \cup \{0, \infty\}, a)$. Then $F_{3,m}$ is clearly determined by its vertices. Sides of $F_{3,m}$ are determined by pairs of adjacent vertices. Since $F_{3,m}$ is a fundamental domain, any adjacent pair of vertices has to be identified with some other pair by an element of $\pi_1(V_{3,m}, a)$. Any element of $\pi_1(V_{3,m}, a)$ identifies at most one pair of vertices of $F_{3,m}$, and the identifying elements freely generate $\pi_1(V_{3,m}, a)$. The set of paths chosen in [**22**] for the vertices of $F_{3,m}$ is Ω_m, where

$$\Omega_m = \gamma_0' * \Omega_m(a_0) \cup \gamma_0 * \Omega_m(\overline{a_0}) \cup \Omega(a_1, -) \cup \Omega(a_1, +)$$

where:

- γ_0' and γ_0' are paths from a_1 to a_0 and $\overline{a_0}$;
- the first endpoint of paths in $\Omega_m(a)$ is at a, and the second endpoints are in $V_3(a)$ for $a = a_0$ and $a = \overline{a_0}$;
- the first endpoint of paths in $\Omega_m(a_1, \sigma)$ is at a_1 and the second endpoints are in $V_3(a_1, \sigma)$, for $\sigma = \pm$.

Since $\rho : \pi_1(V_{3,m}, P_{3,m}, a) \to \pi_1(\overline{\mathbb{C}} \setminus Z_m(s_q), Z_m(s_q), \infty)$ is monotone and injective, it makes sense to find the subset of $\pi_1(V_{3,m}, P_{3,m}, a)$ which forms a vertex set for an ideal polygon which is a fundamental domain, by finding the image under ρ, because ρ preserves the order of vertices, and adjacency of vertices. This was the technique employed in [**22**]. A finite subset of $\pi_1(\overline{\mathbb{C}} \setminus Z_m(s_{3/7}), Z)_m(s_{3/7}), \infty)$ was found, together with a pairing of pairs of elements of this subset under the action of G. It was shown that under very mild extra conditions (which were satisfied) the preimage under ρ is indeed a fundamental domain for $V_{3,m}$, or, more precisely, for the action for the covering group $\pi_1(V_{3,m}, a_1)$ on D. Actually, $\rho = \rho(., s_{3/7})$ is not used directly to describe all of Ω_m in [**22**]. Using four particular paths (called ω_1, ω_{-1}, ω_1' and ω_{-1}') the space $V_{3,m}$ is split into four regions which contain one each of $V_3(a_0)$, $V_3(\overline{a_0})$ and $V_3(a_1, \pm)$. We use $\rho(., s_{3/7})$ to describe paths of $\Omega_m(a_1, \pm)$ — that is, to describe paths in $\Omega_m(a_1, +)$, which are between

ω_1 and ω_1' (including these) and paths in $\Omega_m(a_1,-)$ which are between ω_{-1} and ω_{-1}', including these. We use $\rho(.,s_{1/7})$ to describe paths in $\Omega_m(a_0)$, all of which are strictly between $\overline{\gamma_0'}*\omega_1'$ and $\overline{\gamma_0'}*\omega_{-1}'$, up to homotopy, and $\rho(.,s_{6/7})$ is used to describe paths in $\Omega_m(\overline{a_0})$, which are all strictly between $\overline{\gamma_0}*\omega_1$ and $\overline{\gamma_0}*\omega_{-1}$, up to homotopy.

The description then includes

$$\rho(\Omega_m(a_0),s_{1/7})=\mathcal{Z}_m(1/7),\quad \rho(\Omega_m(\overline{a_0}),s_{6/7})=\mathcal{Z}_m(6/7),$$

$$\rho(\Omega_m(a_1,-),s_{3/7})=\mathcal{Z}_m(a_1,-).$$

The set $\Omega_m(a_1,+)$ is defined by its image under $\rho(.,s_{3/7})$. It has some intersection with $\mathcal{Z}_m(a_1,+)$, but $\rho(\Omega_m(a_1,+))$ neither contains it, nor is contained in it. But part of the result of 1.8 is that each capture $\sigma_\gamma\circ s_{3/7}$, for $\gamma\in\mathcal{Z}_m(a_1,+)$, is equivalent to $\sigma_\beta\circ s_{3/7}$, for some $\beta\in\Omega_m(a_1,+)$. This is the same as saying that, under the identification of $\mathcal{Z}_m(a_1,+)$ with a subset of $\partial D'\cap\partial D$, the G-orbit of $\rho(\Omega_m(a_1,+))$ contains $\mathcal{Z}_m(a_1,+)$.

2.6. Key features of the fundamental domain paths

We now summarize some key features of the paths in $\rho(\Omega_m(a_1,+),s_{3/7})$ from [**22**], and the corresponding generators which identify adjacent pairs. From now on, we write s for $s_{3/7}$ except when this might cause confusion.

The paths in $\rho(\Omega_m(a_1,+),s_{3/7})$ are split into two sets, called R_m and R_m'. The set R_m is totally ordered using an ordering $<$. One way to describe the ordering is just in terms of the ordering of the endpoints in $\partial D\cap\partial D'$. Apart from one common element which is maximal in R_m, the sets R_m and R_m' are disjoint. Adjacent paths in R_m (or R_m') are adjacent in the ordering. Each adjacent pair in R_m is matched with exactly one adjacent pair in R_m' by exactly one adjacent pair in R_m' by exactly one element of G, and vice versa. More precisely, for each adjacent pair (β_1,β_2) of paths in R_m with $\beta_1<\beta_2'$, there is an adjacent pair (β_1',β_2') in R_m' and $[\psi]\in G$ such that $[\psi].[\beta_j']=[\beta_j]$ for $j=1$, 2, and (β_1',β_2'). Using the description of the action of G in 2.5 this means that there is $[\alpha]\in\pi_1(\overline{\mathbb{C}}\setminus Z_m(s),\infty)$ such that $s\simeq_\psi\sigma_\alpha\circ s$ (and $[\alpha]$ is uniquely determined by this property) and:

$$[\alpha*\psi(\beta_j')]=[\beta_j],\quad j=1,2.$$

The common maximal path in R_m and R_m' is a path called $\beta_{1/3}$ from ∞ to $e^{2\pi i(1/3)}$, and contained in $\{z:|z|>1\}$ apart from that endpoint. This is the only path in $R_m\cup R_m'$ whose endpoint is not in a lift of $Z_m(s)$, although it is in $\partial D'$. It is the image under ρ of the path along the negative real axis from a_1 to ∞ (enlarging the domain of ρ slightly, as described in 2.5). In general, if $e^{2\pi ix}$ is on the boundary of a gap of $L_{3/7}$, we write β_x for the capture path which crosses the unit circle at $e^{2\pi ix}$: this is unambiguous, because $e^{2\pi ix}$

is in the boundary of at most one gap. We write $q_p = \frac{1}{3} - 2^{-2p}\frac{1}{21}$. Then $\beta_{q_p} \in \mathcal{Z}_m(a_1, +)$ for each $0 \leq 2p \leq m$: the path $\beta_{2/7}$ is a type II capture path, but the others are all of type III. There is a further decomposition of R_m as

$$R_m = \cup_{0 \leq 2p \leq m} R_{m,p} \cup \{\beta_{1/3}\},$$

where $\beta_{q_{p+1}} \in R_{m,p} \cap R_{m,p+1}$, and $R_{m,p}$ is the set of all paths β in R_m with $\beta_{q_p} \leq \beta \leq \beta_{q_{p+1}}$. The minimal path $\beta_{2/7}$ in R_m is matched with $\beta_{5/7} \in R'_m$, and more generally the minimal path β_{q_p} in $R_{m,p}$ is matched with β_{1-q_p} in $R'_{m,p}$, where we write $R'_{m,p}$ for the set of paths in R'_m which are matched with paths in R'_m.

The paths in $R'_{m,p}$ were never described explicitly in [**22**], and we shall not do so here. They are determined inductively from the paths in $R_{m,0}$, using the ordering on paths of R_m and the matching with paths in R'_m, starting from the matching of $\beta_{2/7}$ with $\beta_{5/7}$. However, we shall use the following later. Let $\beta_1 < \beta_2 \in R_{m,p}$, and let β_i be matched with $\beta'_i \in R_{m,p}$. Let $(\beta_1, \beta_{1,2})$ and $(\beta_{2,1}, \beta_{2,2})$ be adjacent pairs including β_1 and β_2, with $\beta_1 < \beta_{1,2}$ and $\beta_{2,1} < \beta_2$, matched with $(\beta'_1, \beta'_{1,2})$ and $(\beta'_{2,1}, \beta'_2)$ in $R'_{m,p}$, and let (α_1, ψ_1) and (α_2, ψ_2) effect the matching, that is,

$$(s, Y_m(s)) \simeq_{\psi_j} (\sigma_{\alpha_j} \circ s, Y_m(s))$$

and

$$[\alpha_j * \psi_j(\beta'_j)] = [\beta_j], \quad [\alpha_1 * \psi_1(\beta'_{1,2})] = [\beta_{1,2}], \quad [\alpha_2 * \psi_2(\beta'_{2,1})] = [\beta_{2,1}].$$

Then $\alpha_1 * \overline{\alpha_2}$ is trivial in $\pi_1(\overline{\mathbb{C}} \setminus Z_{p+1}(s), \infty)$. This follows from the fact that, up to free homotopy in $\overline{\mathbb{C}} \setminus Z_{2p+2}(s)$, the loop $\alpha_1 * \overline{\alpha_2}$ lies in a set U^p — which is defined precisely in [**22**], and which is also described in the next chapter — and which is disjoint from $Z_{2p+2}(s)$.

2.7. Transferring the fundamental domain under ρ

We have seen that

$$R_m \cup R'_m \subset \rho(\Omega_m).$$

By the monotonicity of ρ, and by the form chosen for $F_{3,m}$, the geodesics homotopic to paths $\overline{\rho(\omega_1)} * \rho(\omega_2)$, for all adjacent pairs (ω_1, ω_2) in Ω, bound a polygon in D', which by abuse of notation we call $\rho(F)$. The translates of F by G cover D' by the Resident's View and the interiors of the translates of F are disjoint, also by the Resident's View. We cannot expect the translates to remain disjoint if we project down from D' to $\overline{\mathbb{C}} \setminus Z(s_q)$, because the covering group of $\overline{\mathbb{C}} \setminus Z(s_q)$ is not G. Nevertheless it is useful to view the translates in the projection. If $(\rho(\omega_1), \rho(\omega_2)) = (\beta_1, \beta_2)$ then an adjacent pair (β_1, β_2) in R_m, then

$$\overline{\rho(\omega_1)} * \rho(\omega_2) = \overline{\beta_1} * \beta_2,$$

and we can identify this with its left to $D' \subset D$, using the lifts of β_1 and β_2 which start from $\widetilde{\infty}$ Similarly if $[\gamma] \in \pi_1(V_{3,m}, a_1)$ with $\rho[\gamma] = \alpha$ and $\Xi([\gamma]) = [\psi]$,then

$$\overline{\rho(\gamma * \omega_1)} * \rho(\gamma * \omega_2) = \psi(\overline{\beta_1} * \beta_2),$$

where, this time, we use the lift of $\alpha * \psi(\beta_1)$ (or $\alpha * \psi(\beta_2)$) starting from $\widetilde{\infty}$ to determine which preimage of $\psi(\overline{\beta_1} * \beta_2)$ we have in D'.

Then we have the following theorem. The proof of this will only be completed in Chapter 4. Remember (as was recalled in 2.4) that $\mathcal{Z}_m(3/7) \subset \rho(\pi_1(V_{3,m}, P_{3,m}, a_1))$, and so for any $\beta \in \mathcal{Z}_m(3/7)$ the point on $\partial D'$ which is identified with β, must be in the orbit under $\pi_1(V_{3,m}, a_1)$ of Ω_m . This theorem immediately gives Theorem 1.8, because the branched coverings $\sigma_\zeta \circ s$, for $\zeta \in R_m$, are in $V_m(a_1, +)$, and it gives the structure for proving 1.9.

THEOREM 2.8. *Let* $\beta \in \mathcal{Z}_m(3/7, +, +, 0)$. *Let*

$$((\omega_{2i-1}, \omega_{2i}) : 1 \leq i \leq n(\beta))$$

and

$$(\gamma_i : 1 \leq i \leq n(\beta))$$

be the unique sequences of, respectively, adjacent pairs in Ω_m *and elements of* $\pi_1(V_{3,m}, a_1)$, *such that* β *successively crosses the arcs* $\overline{\rho(\gamma_i * \omega_{2i-1})} * \rho(\gamma_i * \omega_{2i-1})$, *with* $\beta = \rho(\gamma_{n(\beta)} * \omega_{2n(\beta)})$. *Then the following hold.*

1. *For* $1 \leq i \leq n(\beta)$, *there is an adjacent pair* $(\beta_{2i-1}, \beta_{2i})$ *in* $R_{m,0}$ *and matched with* $(\beta'_{2i-1}, \beta'_{2i})$ *in* $R'_{m,0}$ *such that*

$$(\rho(\omega_{2i-1}), \rho(\omega_{2i-1})) = \begin{cases} (\beta_{2i-1}, \beta_{2i}) \text{ if } i \text{ is odd,} \\ (\beta'_{2i-1}, \beta'_{2i}) \text{ if } i \text{ is even.} \end{cases}$$

Moreover, $\sigma_\beta \circ s$ *is Thurston equivalent to* $\sigma_{\beta_{2n(\beta)-1}} \circ s$, *and* $\Phi(\beta) = \beta_{2n(\beta)-1}$.

2. *Let* $[\psi_{2j-1}]$ *and* α_{2j-1} *effect the matching between the adjacent pairs* $(\beta'_{2j-1}, \beta'_{2j})$ *in* $R'_{m,0}$ *and* $(\beta_{2j-1}, \beta_{2j})$ *in* $R_{m,0}$, *that is:*

$$(s, Y_m) \simeq_{\psi_{2j-1}} (\sigma_{\alpha_{2j-1}} \circ s, Y_m)$$

and

$$\alpha_{2j-1} * \psi_{2j-1}(\beta'_{2j-1}) = \beta_{2j-1}, \quad \alpha_{2j-1} * \psi_{2j-1}(\beta'_{2j}) = \beta_{2j}.$$

Write

$$\psi_{4j-3,4j-1} = \psi_{4j-3} \circ \psi_{4j-1}^{-1}, \quad \alpha_{4j-3,4j-1} = \alpha_{4j-3} * \psi_{4j-3,4j-1}(\overline{\alpha_{4j-1}}),$$

and, for $j < k$,

$$\psi_{4j-3,4k-1} = \psi_{4j-3,4j-1} \circ \cdots \circ \psi_{4k-3,4k-1}$$

$$\psi_{4j-3,4k+1} = \psi_{4j-3,4k-1} \circ \psi_{4k+1},$$

$$\alpha_{4j-3,4k-1} = \alpha_{4j-3,4j-1} * \psi_{4j-3,4j-1}(\alpha_{4j+1,4k-1}) = \alpha_{4j-3,4k-5} * \psi_{4j-3,4k-5}(\alpha_{4k-3,4k-1}),$$

and

$$\alpha_{4j-3,4k+1} = \alpha_{4j-3,4j-1} * \psi_{4j-3,4j-1}(\alpha_{4j+1,4k+1}) = \alpha_{4j-3,4k-1} * \psi_{4j-3,4k-1}(\alpha_{4k+1}).$$

Then, for $r = 0$ or 1, taking $\psi_{1,-1}$ to be the identity, and $\alpha_{1,-1}$ to be trivial,

$$\rho(\gamma_i) = \alpha_{1,2i-3}, \quad \Xi(\gamma_i) = [\psi_{1,2i-3}],$$

$$\rho(\gamma_i * \omega_{2i-r}) = \begin{cases} \alpha_{1,2i-3} * \psi_{1,2i-3}(\beta_{2i-r}) \text{ if } i \text{ is odd,} \\ \alpha_{1,2i-3} * \psi_{1,2i-3}(\beta'_{2i-r}) = \alpha_{1,2i-1} * \psi_{1,2i-1}(\beta_{2i-r}) \text{ if } i \text{ is even.} \end{cases}$$

In other words

$$\rho(\gamma_i * \omega_{2i-r}) = \alpha_{1,4\lfloor i/2\rfloor-1} * \psi_{1,4\lfloor i/2\rfloor-1}(\beta_{2i-r}).$$

Consequently if i is odd then $\rho(\gamma_i * \omega_{2i-1})$ and $\rho(\gamma_i * \omega_{2i})$ are arbitrarily small perturbations of

$$\beta_1 * \psi_1(\overline{\beta'_1} * \beta'_3) * \psi_{1,3}(\overline{\beta_3} * \beta_5) \cdots * \psi_{1,2i-5}(\overline{\beta'_{2i-5}} * \beta'_{2i-3}) * \psi_{1,2i-3}(\overline{\beta_{2i-3}} * \beta_{2i-1}),$$

and

$$\beta_2 * \psi_1(\overline{\beta'_2} * \beta'_4) * \psi_{1,3}(\overline{\beta_4} * \beta_6) \cdots * \psi_{1,2i-5}(\overline{\beta'_{2i-4}} * \beta'_{2i-2}) * \psi_{1,2i-3}(\overline{\beta_{2i-2}} * \beta_{2i}),$$

and if i is even they are arbitrarily small perturbations of

$$\beta_1 * \psi_1(\overline{\beta'_1} * \beta'_3) * \psi_{1,3}(\overline{\beta_3} * \beta_5) \cdots * \psi_{1,2i-5}(\overline{\beta_{2i-5}} * \beta_{2i-3}) * \psi_{1,2i-3}(\overline{\beta'_{2i-3}} * \beta'_{2i-1}),$$

and

$$\beta_2 * \psi_1(\overline{\beta'_2} * \beta'_4) * \psi_{1,3}(\overline{\beta_4} * \beta_6) \cdots * \psi_{1,2i-5}(\overline{\beta_{2i-4}} * \beta_{2i-2}) * \psi_{1,2i-3}(\overline{\beta'_{2i-2}} * \beta'_{2i}).$$

In particular,

$$\Phi(\beta) = \psi^{-1}_{1,4\lfloor n(\beta)/2\rfloor-1}(\overline{\alpha_{1,4[n(\beta)/2]-1}} * \beta).$$

2.9. Remarks

It does not seem to be true that if $\beta \in \mathcal{Z}_m(3/7,+,-,p)$ then $\omega_{2n(\beta)-1}(\beta) \in \Omega_{m,p}$, even for $p = 0$. Some calculations in Chapter 6 suggest that it might be true that, for such a β,

$$\omega_{2n(\beta)-1}(\beta) \in \Omega_{m,p} \cup \Omega_{m,p+1}.$$

The main statement in Theorem 2.8 is item 1. Since an adjacent pair $(\beta_{2i-1}, \beta_{2i})$ in $R_{m,0}$ is only matched with an adjacent pair $(\beta'_{2i-1}, \beta'_{2i})$ in $R'_{m,0}$, and indeed any adjacent pair in Ω_m is only matched with one other, γ_i is uniquely determined by $(\omega_{2j-1}, \omega_{2j})$ for $j < i$. More precisely $\overline{\gamma_i} * \gamma_{i+1}$ is determined by $(\omega_{2i-1}, \omega_{2i})$, and hence the same is true for the images under $\rho$and Ξ. This is true whichever pairs $(\omega_{2i-1}, \omega_{2i})$ arise. But item 1 tells us(remembering that Ξ is a group homomorphism and ρ is not one, but is a twisted group homomorphism), that

$$\rho(\overline{\gamma_i} * \gamma_{i+1}) = \begin{cases} \alpha_{2i-1} \text{ if } i \text{ is odd,} \\ \psi_{2i-1}^{-1}(\overline{\alpha_{2i-1}}) \text{ if } i \text{ is even.} \end{cases}$$

and

$$\Xi(\overline{\gamma_i} * \gamma_{i+1}) = \begin{cases} [\psi_{2i-1}] \text{ if } i \text{ is odd,} \\ [\psi_{2i-1}^{-1}] \text{ if } i \text{ is even} \end{cases}$$

So item 2 follows from item 1.

Part of the proof needs some detailed analysis of the homeomorphisms $\psi_{2j-1,2k-1}$. The proof of this result is therefore given in two stages, partly in Chapter 3 and partly in Chapter 4.

2 of 2.8 also suggests an algorithm for determining when two different captures $\sigma_\beta \circ s$ and $\sigma_\zeta \circ s$ are equivalent, for β and $\zeta \in \mathcal{Z}_m(a_1,+,+,0)$.

Write

$$\beta_i(\beta), \quad \psi_{2i-1,\beta}, \quad \psi_{2j-1,2k-1,\beta}, \quad \alpha_{2j-1,\beta}, \quad \alpha_{2j-1,2k-1,\beta}$$

for β_i, ψ_{2i-1}, $\psi_{2j-1,2k-1}$, α_{2j-1} $\alpha_{2j-1,2k-1}$ in Theorem 2.8. Then $\sigma_\beta \circ s$ and $\sigma_\zeta \circ s$ are Thurston equivalent if and only if $(\beta_{2n(\beta)-1}(\beta), \beta_{2n(\beta)}(\beta)) = (\beta_{2n(\zeta)-1}(\zeta), \beta_{2n(\zeta)}(\zeta))$. If $\beta \neq \zeta$, then it must be the case that $\beta_i(\beta) \neq \beta_i(\zeta)$ for at least one i. Theorem 1.9 will be proved simply by exploring how this is possible.

CHAPTER 3

Symbolic dynamics, paths in $R_{m,0}$, and homeomorphisms

The aim of this chapter is to analyse the homeomorphism which will arise as the homeomorphisms $\psi_{2i-1,2j-1}$ which arise in Theorem 2.8. This will enable us to do much of the proof of Theorem 2.8 at the end of the chapter.

3.1. Symbolic Dynamics

Symbolic dynamics with respect to a Markov partition for h_{a_1}, or equivalently, for $s_{3/7}$ (using the semiconjugacy from $s_{3/7}$ to h_{a_1}), underpin the analysis in this paper. We shall work almost exclusively with $s_{3/7}$, in preference to h_{a_1}.

The Markov partition for $s_{3/7}$ is a partition of the unit disc. It is drawn below. This partition was also used in [**22**], and introduced in section 2.9 of that paper. The boundaries of the sets of the partition, within the unit disc, are the vertical lines joining $e^{\pm 2\pi ir/14}$ for $1 \leq r \leq 6$. The labels of the closures of the sets of the partition are, as shown in Figure 3, L_j and R_j for $1 \leq j \leq 3$, and the central region C. We shall write $D(X)$ for the closed set labelled X. We also denote by $D(BC)$ and $D(UC)$ the components of the complement in $D(C)$ of the central gap of $L_{3/7}$, where $D(BC)$ is the component intersecting the lower unit circle and $D(UC)$ is the component intersecting the upper unit circle, and the labels of these sets are BC and UC.

The dynamics under $s_{3/7}$ are as follows:

$$s_{3/7}(D(L_1)) = s(D(R_1)) = D(R_1) \cup D(R_2),$$

$$s_{3/7}(D(L_2)) = s_{3/7}(D(R_2)) = D(R_3) \cup D(C),$$

$$s_{3/7}(D(L_3)) = s_{3/7}(D(R_3)) = D(L_3) \cup D(L_2), \quad D(C) = D(L_1).$$

Also, $s_{3/7}(D(BC)) = s_{3/7}(D(UC))$ is the complement in $D(L_1)$ of the gap containing $s(0)$. Note that $s_{3/7}$ is a homeomorphism onto image, restricted to $D(L_j)$ or $D(R_j)$ or $D(BC)$ or $D(UC)$, but a degree two branched cover restricted to $D(C)$.



Figure 1. Markov partition for $s_{3/7}$

As usual in symbolic dynamics, we use words over the alphabet $\{L_j, R_j : 1 \leq j \leq 3\} \cup \{C, BC, UC\}$ to label subsets in the unit disc, so that for a word $w = X_0X_1 \cdots X_{n-1}$ labels the subset $D(w)$ where

$$D(w) = \cap_{j=0}^{n-1} s_{3/7}^{-j}(D(X_j)).$$

Then, as usual, $D(w) \neq \emptyset$ if and only if w is admissible, that is, $D(X_j) \cap s_{3/7}(D(X_{j-1})) \neq \emptyset$ for $0 < j \leq n-1$. We will use BC or UC rather than C, except for the last letters of words, but do not allow BC or UC for a last letter, using C instead. Under this convention, we have, for all admissible words $X_0 \cdots X_{n-1}$,

$$s_{3/7}(D(X_0 \cdots X_{n-1})) = D(X_1 \cdots X_{n-1}).$$

If w is an admissible word of length $m > 1$ (or $m > 2$) ending in L_2C (or R_1R_2C), then $D(w)$ contains a unique point of preperiod $m-1$ (or $m-2$) under s. We denote this point by $p(w)$, and in this circumstance we will sometimes talk of w having preperiod $m-1$ (or $m-2$).

3.2. Paths in $R_{m,0}$

All paths in $R_{m,0}$ have endpoints in $U^0 \cap Z_m$, where U^0 is the set defined by

$$U^0 = \cup_{i=0}^{\infty} D(L_3^{3i+1}L_2) \cup D(BC) \cup_{i=0}^{\infty} D(L_3^{2+2i}L_2BC) \setminus \cup_{i=0}^{\infty} D(L_3^{1+3i}L_2BC).$$

Not all points in $U^0 \cap Z_m$ are endpoints of paths in $R_{m,0}$, and some points are endpoints of two such paths in $R_{m,0}$, but no point in $U^0 \cap Z_m$ is the endpoint of more than one path in $R_{m,0}$, and each path in $R_{m,0}$ is uniquely determined up to isotopy preserving Z_m by its first S^1-crossing point and its final endpoint.

The description of the paths uses finite sequences of words

$$(w_i(w,0) : 1 \leq i \leq n(w)), \quad (w_i'(w,0) : 1 \leq i \leq n(w)),$$

for each w ending in C with $D(w) \subset U^0$and

$$(w_i'(w,x) : 1 \leq i \leq n(w)).$$

each w ending in C for $D(w) \subset U^x$, for a certain set U^x which will be defined in 3.7. Since we are only concerned in detail with $R_{m,0}$, and not with $R_{m,p}$ for $p > 0$, we write

$$w_i(w,0) = w_i(w), \quad w_i'(w,0) = w_i'(w).$$

For $\beta \in R_{m,0}$, we write $w(\beta)$ for the word such that the final endpoint of β is $p(w(\beta))$, and also write

$$w_i(\beta) = w_i(w(\beta)), \quad w_1'(\beta) = w_1'(w(\beta), \quad w_i'(\beta) = w_i'(w(\beta), w_1'(\beta)).$$

The i'th unit-disc crossing of β is determined by $w_i'(\beta)$ as follows. If the last letter of $w_i'(\beta)$ is L_3 or R_3, then the i'th crossing of β is along the set $D(w_i'(\beta)(L_2R_3L_3)^\infty)$, which is a leaf of $L_{3/7}$. To determine the direction of the crossing, we use the unique gap G of $L_{3/7}$ which has this leaf in its boundary. The direction of the crossing is in an anticlockwise direction along the boundary of G. If $i = n(x)$ then β crosses into G and ends at the point of $G \cap Z_m$, starting at what would be the start point of a crossing of the leaf which is part of an anticlockwise circuit round G.

Now suppose that the last letter of $w_i'(\beta)$ is BC. In this case, $i < n(x)$. Let u' be the word obtained from w_i' by replacing the last letter by UC. Then $i > 1$, and u' and w_i' both have w_{i-1}' as a prefix, while u' is a prefix of w_j' for $i < j \leq n(w(\beta))$. The words u' and w_i' then label disjoint subsets $\partial D(u')$ and $\partial D(w_i')$ of $\{z : |z| \leq 1\}$. The i'th crossing by $\beta(w,x,0)$ of $\{z : |z| \leq 1\}$ is then from the first point of $\partial D(w_i')$ to the first point of $\partial D(u')$, using anticlockwise order on these intervals.

3.3. Properties of the sequences w_i, w_i', $w_i'(.,x)$

The following properties hold for the sequences. Property 4 allows $w_i(w)$ and $w_i'(w)$ to be defined by induction on i, because there is always at least one occurrence of L_3 or BC or UC after the last letter of $w_{i-1}(w)$ and at or before the last letter of $w_i(w)$. Similarly, Properties 5 and 6 will allow for an inductive definition of $w_i'(w,x)$ if $D(w) \subset U^x$.

1. $w_i(w)$ is a prefix of w of length increasing with i, and is also a prefix of $w_{i+1}'(w)$ if $i < n(w)$.
2. $w_i'(w)$ always ends in L_3 or R_3 or BC, and is a function only of i and $w_i(w)$.
3. $w_1'(w,x) = x$ if $D(w) \subset U^x$

4. Let $w = uy$ and $D(y) \subset U^0$. If $|u| < w_1(w)$, then $w_1(uy) = uw_1(y)$ and $w_1'(uy) = uw_1'(y)$. If $w = uy$ and $|w_{i-1}(w)| < |u| \leq |w_i(w)|$, then $w_j(w) = uw_{j-i}(y)$ and $w_j'(uy) = uw_{j-i}'(y)$ for $i < j \leq n(w)$. If $w = uUCy_1$ and $w_{i-1}(w)| < |uUC| \leq |uUC|$, then $w_j(w) = uUCy_2$ where $w_{j-i}(BCy) = BCy_2$, for $i < j \leq n(x)$.
5. If $w = uy$, where $u = u_1u_2$ and $D(v_1u_2) \subset U^x$, with
$$|w_{i-1}(w)| < |u_1| \leq |w_i(w)|$$
and
$$|w_1(v_1u_2y)| \leq |v_1| < |w_2(v_1u_2y)| \leq |v_1u_2|,$$
then
$$w_j'(w, x) = uv_2$$
for $i < j \leq n(x)$, where
$$v_1v_2 = w_{j-i+1}'(v_1u_2y, x).$$
6. If $w_i(w)$ ends in BC or UC, then $w_i'(w, x) = w_i'(w)$.

Thus w_i and w_i' and $w_i'(.,x)$ can be defined inductively from w_1 and w_1' and $w_1'(.,x)$. The definitions are used explicitly in some sections of this article. We therefore include them here. For full details, see Chapter 7 of [**22**].

3.4. Definition of w_1

We define $w_1(w)$ for w with $D(w) \subset U^0$ and with w ending in C. The definition also works for infinite words. The first letter of w is either BC or L_3. If BC is the first letter, then $w_1(w) = BC$. If L_3 is the first letter of w, and w starts with L_3^4 or $L_3^2L_2BC$, then $w_1(w) = L_3^4$ or L_3^2 respectively. Otherwise, w starts L_3L_2, and we look for the first occurrence of one of the following. One of these must occur if w is preperiodic, that is, ends in C.

0. An occurrence of C.
1. An occurrence of xBC for any $n \geq 0$, where x is a maximal word in the letters L_3, L_2, R_3 with an even number of L letters.
2. An occurrence of xUC, where x is a maximal word in the letters L_3, L_2, R_3, with an odd number of L letters.
3. An occurrence of L_1R_2UC.
4. An occurrence of R_1R_2BC.
5. An occurrence of a string $v_1 \cdots v_nL_3$ which is the end of w, or is followed in w by L_2, where:
   - $v_i = (L_3L_2R_3)^{q_i}L_3(L_2R_3)^{m_i}L_3^{r_i-1}$ for $i \geq 2$, where $m_i + r_i = 3$ and $q_i \geq 0$;

- either $v_1 = (L_3L_2R_3)^{q_1}L_3(L_2R_3)^{m_1}L_3^{r_1-1}$ for $q_1 \geq 0$ and $m_1 \geq 1$ and an odd $m_1+r_1 \geq 3$, or $v_1 = XL_1R_1^{p_1}R_2R_3(L_2R_3)^{m_1}L_3^{r_1-1}$ for $3 \leq r_1 \leq 5$, and $m_1 \geq 0$, and $p_1 \geq 0$, and $X = BC$ or UC;
- v_nL_3 is either followed by L_2, or is at the end of w;
- the string $v_1 \cdots v_nL_3$ is maximal with these properties;
- n is odd.

We note that the term $(L_3L_2R_3)^{q_i}$ with $q_i \geq 0$. was missing in the definition of [**22**]. This is relevant to the calculations in this paper, because we shall have examples with $q_i = 1$ for some i.

We define $w_1(w)$ to be the prefix of w ending in the occurrence listed – except in case 1 when $n > 0$, and in case 4 when w ends in $L_1R_1R_2BC$, and in case 5. In case 4, we define $w_1(w)$ to be the word up to, and not including, this occurrence of $L_1R_1R_2BC$, and $w_2(w,0)$ to be the prefix of w ending in this occurrence of $L_1R_1R_2BC$. In case 5, if the maximal string of words of this type is $v_1 \cdots v_n$, then we define $w_1(w,0)$ to be the prefix ending with v_nL_3.

3.5. Definition of $w_2(w)$ if $D(w) \subset D(BC)$

Now suppose that the first letter of w is BC. Then w starts with $(BCL_1R_2)^k$ for a maximal $k \geq 0$. Then we define $w_2(w) = w$ if there is no occurrence of cases 1 to 5 above. We define $w_2(w)$ to be the prefix of w defined exactly as $w_1(w)$ is defined when w starts with L_3, if there is an occurrence of one of 2 to 5.

3.6. Definition of w_1'

Now we define $w_1'(w,0) = w_1'(w_1,0)$. If $w = C$ or if w_1 starts with BC or L_3^2, then we define $w_1'(w,0) = L_3$. Now suppose that w starts with L_3L_2, and we assume that one of cases 0 to 5 holds, which, as pointed out, is always true if w ends in C.

0. w_1' is the same length as w_1, ending in L_3, if preceded by a maximal subword in the letters L_3, L_2 or R_3 and with an even number of L letters, or ending in R_3 if preceded by a maximal subword in the letters L_3, L_2 or R_3 and with an even number of L letters, such that $D(w_1')$ and $D(w_1)$ are adjacent.
1. Let w_1'' be obtained from w_1 by deleting the last $(UCL_1R_2)^nBC$, and adding C. Then w_1' is the same length as w_1'', ending in L_3, such that $D(w_1')$ and $D(w_1'')$ are adjacent. If $n > 0$ then, w_2' is the same length as w_1'', ending in R_3, such that $D(w_2')$ and $D(w_1'')$ are adjacent.

2. Let w_1'' be obtained from w_1 by deleting the last UC and adding C. Then w_1' is defined similarly to 1, but ending in R_3.
3. w_1' is obtained from w_1 by replacing the last letter by BC.
4. If w_1 ends in $R_1^2R_2BC$, then w_1' is obtained from w_1 by replacing the last letter by $R_3L_2R_3L_3$. If w_1 is extended in w by $L_1R_1R_2BC$, then $w_1'(w,0)$ is the word of the same length as w_1, with $D(w_1')$ to the left of $D(v')$, where v' is obtained from w_1 by replacing the last letter by C. We then also define $w_2'(w,0) = v''$, where v'' is of the same length as v', with $D(v')$ between $D(w_1')$ and $D(v'') = D(w_2')$.
5. The word w_1' is obtained from w_1 by deleting all but the first letter of v_nL_3 if $n > 1$, or if $n = 1$, $m_1 + r_1 = 3$ and the first letter of v_1 is L_3. In the other cases with $n = 1$, the word w_1' is obtained from w_1 by replacing the last L_3^3 by $L_2R_3L_3$ if $r_1 \geq 3$, by deleting the last $L_2R_3L_3^2$ if $r_1 = 2$, and by deleting the last $(L_2R_3)^2L_3$ if $r_1 = 1$.

3.7. Definition of $w_1'(.,x)$

This definition generalises the definition given in [**22**], where the only possible value of $w_1'(.,x)$ was x, and this was defined only for w with $D(w) \subset U^x$, for a proper subset U^x of U^0.

For $i \geq 1$ we define $w_i'(w,x) = w_i'(w)$ except if $w = uy_1v$ or uy_2v, where $w_i(w) = uy_1L_3$ or uy_2L_3 and $D(v) \subset D(L_3L_2)$ is to the right of $D(x(L_2R_3L_3)^\infty)$ and

$$(y_1, y_2) = (L_3(L_2R_3)^{2j}L_3^4, L_3(L_2R_3)^{2j+2})$$

for some $j \geq 0$, or

$$(y_1, y_2) = (L_3(L_2R_3)^{2j}L_3^2, L_3(L_2R_3)^{2j-1}L_3^2L_2R_3)$$

for some $j \geq 1$. We then define

$$w_i'(uy_1v, x) = w_i'(uy_2v) = w_i'(uy_2L_3L_2), \quad w_i'(uy_2v, x) = w_i'(uy_1v) = w_i'(uy_1L_3L_2).$$

We then define U^x to be the union of all $D(w)$ with $w_1'(w,x) = x$. This agrees with the definition given in [**22**].

The definition of $w_i'(.,x)$ might seem as if it comes from nowhere. It is actually related to certain homeomorphisms which determine the group action on the universal cover of $V_{3,m}$, and hence suggest possible fundamental domains for this action. Homeomorphisms of a similar type also play an important role in the current work, since each capture path is equivalent under the group action to precisely two vertices of the fundamental domain. In both cases, the homeomorphisms act on both paths and on Z_m, and, in both cases, *exchangeable pairs* are important indefining the homeomorphisms.

3.8.

For the proofs of the main theorems of this paper, we need to analyse the homeomorphisms $\psi_{4j-3,4j-1,\beta} = \psi_{4j-3,\beta} \circ \psi^{-1}_{4j-1,\beta}$, which are defined in 2.8 and 2.9. These definitions assume the existence of $\beta_\ell(\beta)$ for $\ell \leq 4j$ with the properties claimed in 2.8. The proof of the existence of the $\beta_\ell(\beta)$ will be inductive, and the existence of $\beta_{4j+t}(\beta)$ for $1 \leq t \leq 4$ will involve an analysis of the homeomorphisms $\psi_{4k-3,4k-1,\beta}$ for $k \leq j$.

More generally, we consider homeomorphisms $\psi_{\zeta,\eta} = \psi_\zeta \circ \psi_\eta^{-1}$ whenever (ζ, ζ_2) and (η, η_2) are adjacent pairs in $R_{m,0}$, with $\zeta < \zeta_2$ and $\eta < \eta_2$, and ψ_ζ matches (ζ, ζ_2) with the adjacent pair (ζ', ζ_2') in $R'_{m,0}$, while ψ_η matches (η, η_2) with (η', η_2'). This means that there are $[\alpha_\zeta]$ and $[\alpha_\eta] \in \pi_1(\overline{\mathbb{C}} \setminus Z_m, \infty)$ such that

$$s \simeq_{\psi_\zeta} \sigma_{\alpha_\zeta} \circ s$$

$$[\alpha_\zeta * \psi_\zeta(\zeta')] = [\zeta], \quad [\alpha_\zeta * \psi_\zeta(\zeta_2')] = [\zeta_2]$$

and similarly for η.

We start with a very common lemma which has been used many times in the course of preceding work to compute the homeomorphism ψ such that $(s, Y_m) \simeq_\psi (\sigma\alpha \circ s, Y_m)$, and, in particular, to bound the support of ψ.

Lemma 3.8.1. *Suppose that α is a closed lop based at $v_2 = \infty$ which is trivial in $\pi_1(\overline{\mathbb{C}} \setminus Y_0, v_2)$. and suppose that ψ is a homeomorphism such that $(s, Y_m) \simeq_\psi (\sigma_\alpha \circ s, Y_m)$. Then ψ can be defined inductively by the equations: ψ_1 is the identity in* $\mathrm{Mod}(\overline{\mathbb{C}}, Y_0)$*, $\psi = \psi_m$, and*

$$s = \sigma_\alpha \circ s \circ \psi_1$$

and

$$\psi_i = \psi_{i+1} \text{ in } \mathrm{Mod}(\overline{\mathbb{C}}, Y_i)$$

and

$$\psi_i \circ s = \sigma_\alpha \circ s \circ \psi_{i+1}$$

Alternatively we can define ψ_{i+1} by

$$\psi_{i+1} = \xi_i \circ \psi_i$$

where

$$\xi_0 = \psi_1$$

and

$$\xi_i = \text{identity in } \mathrm{Mod}(\overline{\mathbb{C}}, Y_i)$$

and

$$\sigma_\alpha \circ s \circ \xi_{i+1} = \xi_i \circ \sigma_\alpha \circ s.$$

PROOF. It is clear that if ψ_i are defined by the first set of equations, and $\psi = \psi_m$, then ψ has the required properties. So it remains to show that the alternative definitions are equivalent to the original ones. If we have

$$\psi_i \circ s = \sigma_\alpha \circ s \circ \psi_{i+1} \tag{3.8.1.1}$$

and

$$\psi_{i+1} \circ s = \sigma_\alpha \circ s \circ \psi_{i+2}, \tag{3.8.1.2}$$

then writing $\psi_{i+1} = \xi_i \circ \psi_i$ and $\psi_{i+2} = \xi_{i+1} \circ \psi_{i+1}$ in the second equation we have

$$\xi_i \circ \psi_i \circ s = \sigma_\alpha \circ s \circ \xi_{i+1} \circ \psi_{i+1}$$

Combining these, we obtain

$$\xi_i \circ \sigma_\alpha \circ s \circ \psi_{i+1} = \sigma_\alpha \circ s \circ \xi_{i+1} \circ \psi_{i+1}$$

which gives

$$\xi_i \circ \sigma_\alpha \circ s = \sigma_\alpha \circ s \circ \xi_{i+1} \tag{3.8.1.3}$$

Conversely, we see that (3.8.1.3) implies (3.8.1.2) given (3.8.1.1), as required. □

This can be used to bound the support of $\psi_{1,4j-1}$, but, as it turns out, a variant of 3.8.1 is more useful. We recall from 2.8 that

$$s \simeq_\psi \sigma_\alpha \circ s$$

where $\alpha = \alpha_{1,4j-1}$ where

$$\alpha_{1,4j-1} = \alpha_{1,3} * \psi_{1,3}(\alpha_{5,7}) * \cdots * \psi_{1,4j-5})(\alpha_{4j-3,4j-1}).$$

Writing

$$\alpha_{4j-3,4j-1,1} = \psi_{1,4j-5}(\alpha_{4j-3,4j-1}),$$

we have

$$\alpha_{1,4j-1} = \alpha_{1,3} * \alpha_{5,7,1} * \cdots * \alpha_{4j-3,4j-1,1}.$$

We then write $E_{4j-3,4j-1,1}$ for the topological disc enclosed by the closed loop $\alpha_{1,4j-3,4j-1}$. We are only interested in $E_{4j-3,4j-1,1}$ up to homotopy preserving Z_∞. We have

$$\sigma_\alpha = \sigma_{\alpha_{4j-3,4j-1,1}} \circ \cdots \circ \sigma_{\alpha_{1,3}}.$$

Hence if we define $\psi_{1,4j-1,1}$ to be homotopic to the identity relative to Z_0, and to satisfy the equation

$$s = \sigma_\alpha \circ s \circ \psi_{1,4j-1,1}$$

then we have

$$\psi_{1,4j-1,1} = \psi_{1,3,1,1} \circ \cdots \circ \psi_{4j-3,4j-1,1,1}$$

where

$$s = \sigma_{\alpha_{4i-3,4i-1}} \circ s \circ \psi_{4i-3,4i-1,1,1}.$$

the loop $\alpha_{4i-3,4i-1}$ is a perturbation of $\alpha_{4i-3} * \psi_{4i-3,4i-1}(\overline{\alpha_{4i-1}})$. This is awkward to work with, simply because the effect of $\psi_{4i-3,4i-1}$ on α_{4i-1} is unclear. We will therefore employ a slightly different approach.

We write $E_{\zeta,\eta}$ for an arbitrarily small neighbourhood of the closed disc bounded by a loop homotopic to $\alpha_\zeta * \overline{\alpha_\eta}$, where the homotopy preserving Z_∞ and the disc does not contain $v_2 = c_2 = \infty$. Naturally, we wish to apply this in the case $(\zeta, \eta) = (\beta_{4i-3}(\beta), \beta_{4i-1}(\beta))$ for $\beta \in \mathcal{Z}_m(3/7, +, +, 0)$, once the existence of $\beta_\ell(\beta)$ as in Theorem 2.8 has been established for $\ell \leq 4i$. We shall therefore be particularly interested in the following in the case $(\zeta, \eta) = (\beta_{4i-3}(\beta), \beta_{4i-1}(\beta))$, for varying i.

LEMMA 3.8.2. *If k is the smallest integer for which $\alpha_{\zeta,\eta} = \alpha_\zeta * \overline{\alpha_\eta}$ is nontrivial in $\pi_1(\overline{\mathbb{C}} \setminus Z_k, v_2)$, then $\psi_\zeta \circ \psi_\eta^{-1} = \psi_{\zeta,\eta} = \psi_{\zeta,\eta,m+1}$ can be defined inductively as an element of* $\mathrm{Mod}(\overline{\mathbb{C}}, Y_{m+1})$ *by the equations*

$$\sigma_{\alpha_{\zeta,\eta}} \circ s \circ \psi_{\zeta,\eta,k+1} = s, \tag{3.8.2.1}$$

and for $k < \ell \leq m$

$$\psi_{\zeta,\eta,\ell+1} = \xi_{\zeta,\eta,\ell} \circ \psi_{\zeta,\eta,\ell} \tag{3.8.2.2}$$

$$\sigma_{\alpha_\zeta} \circ s \circ \xi_{\zeta,\eta,\ell} = \xi_{\zeta,\eta,\ell-1} \circ \sigma_{\alpha_\zeta} \circ s. \tag{3.8.2.3}$$

It follows that the support of $\psi_{\zeta,\eta,\ell}$, up to isotopy preserving Z_m, is

$$\cup_{1 \leq i \leq \ell - r} (\sigma_{\alpha_\zeta} \circ s)^{-i}(E_{\zeta,\eta}).$$

Proof. We have

$$\sigma_{\alpha_\zeta} \circ s \circ \psi_\zeta = \psi_\zeta \circ s \text{ rel } Y_{m+1}, \tag{3.8.2.4}$$

$$\sigma_{\alpha_\eta} \circ s \circ \psi_\eta = \psi_\eta \circ s \text{ rel } Y_{m+1}. \tag{3.8.2.5}$$

Here, $[\psi_\zeta]$ and $[\psi_\eta]$ are the only elements of $\mathrm{Mod}(\overline{\mathbb{C}}, Y_{m+1})$ satisfying (3.8.2.4) and (3.8.2.5) respectively. Now (3.8.2.5) gives

$$\psi_\eta^{-1} \circ \sigma_{\alpha_\eta} \circ s = s \circ \psi_\eta^{-1} \text{ rel } Y_{m+1}. \tag{3.8.2.6}$$

Then combining (3.8.2.4) and (3.8.2.6), we have

$$\sigma_{\alpha_\zeta} \circ s \circ \psi_{\zeta,\eta} = \psi_\zeta \circ s \circ \psi_\eta^{-1} = \psi_{\zeta,\eta} \circ \sigma_{\alpha_\eta} \circ s \text{ rel } Y_{m+1}. \tag{3.8.2.7}$$

If this equation is satisfied by $[\chi]$ replacing $[\psi_{\zeta,\eta}]$, then (3.8.2.4) is satisfied with $[\chi \circ \psi_\eta]$ replacing $[\psi_\zeta]$, which implies that $[\chi] = [\psi_{\zeta,\eta}]$. So (3.8.2.6) uniquely determines $[\psi_{\zeta,\eta}] \in \mathrm{MG}(\overline{\mathbb{C}}, Y_{m+1})$.

Now we define $\psi_{\zeta,\eta,k+1}$ by (3.8.2.1) and $[\psi_{k+1}] = [\text{identity}]$ in $\mathrm{MG}(\overline{\mathbb{C}}, Y_k))$, and define $\psi_{\eta,\zeta,k}$ to be the identity, we have

$$\xi_{\zeta,\eta,k} = \psi_{\zeta,\eta,k+1}.$$

Then we clearly want to define $\psi_{\zeta,\eta,\ell+1}$ in terms of $\psi_{\zeta,\eta,\ell}$ by $[\psi_{\zeta,\eta,\ell+1}] = [\psi_{\zeta,\eta,\ell}]$ in $\mathrm{MG}(\overline{\mathbb{C}}, Y_\ell)$ and:

$$\sigma^{-1}_{\psi_{\zeta,\eta,\ell}(\alpha_\eta)} \circ \sigma_{\alpha_\zeta} \circ s \circ \psi_{\zeta,\eta,\ell+1} = \psi_{\zeta,\eta,\ell} \circ s, \tag{3.8.2.8}$$

If we do this for $\ell \leq m+1$, then we will have $[\psi_{\zeta,\eta,m+2}] = [\psi_{\zeta,\eta,m+1}]$ in $\mathrm{MG}(\overline{\mathbb{C}}, Y_{m+1})$ and (3.8.2.6) is satisfied by $\psi_{\zeta,\eta,m+1}$ replacing $\psi_{\zeta,\eta}$, which, as we have seen, implies that $[\psi_{\zeta,\eta,m+1}] = [\psi_{\zeta,\eta}]$ Now (3.8.2.8) is equivalent to:

$$\sigma^{-1}_{\psi_{\zeta,\eta,\ell}(\alpha_\eta)} \circ \sigma_{\alpha_\zeta} \circ s \circ \xi_{\zeta,\eta,\ell} \circ \psi_{\zeta,\eta,\ell} = \xi_{\zeta,\eta,\ell-1} \circ \psi_{\zeta,\eta,\ell-1} \circ s. \tag{3.8.2.9}$$

But then this gives, by (3.8.2.8) with $\ell - 1$ replacing ℓ,

$$\sigma^{-1}_{\psi_{\zeta,\eta,\ell}(\alpha_\eta)} \circ \sigma_{\alpha_\zeta} \circ s \circ \xi_{\zeta,\eta,\ell} \circ \psi_{\zeta,\eta,\ell} = \xi_{\zeta,\eta,\ell-1} \circ \sigma^{-1}_{\psi_{\zeta,\eta,\ell-1}(\alpha_\eta)} \circ \sigma_{\alpha_\zeta} \circ s \circ \psi_{\zeta,\eta,\ell}$$

Then this gives

$$\sigma^{-1}_{\psi_{\zeta,\eta,\ell}(\alpha_\eta)} \circ \sigma_{\alpha_\zeta} \circ s \circ \xi_{\zeta,\eta,\ell} \circ \psi_{\zeta,\eta,\ell} = \sigma^{-1}_{\psi_{\zeta,\eta,\ell}(\alpha_\eta)} \circ \xi_{\zeta,\eta,\ell-1} \circ \sigma_{\alpha_\zeta} \circ s \circ \psi_{\zeta,\eta,\ell},$$

which gives

$$\sigma_{\alpha_\zeta} \circ s \circ \xi_{\zeta,\eta,\ell} = \xi_{\zeta,\eta,\ell-1} \circ \sigma_{\alpha_\zeta} \circ s$$

which gives (3.8.2.3), from which we see that the support of $\xi_{\zeta,\eta,\ell}$ is

$$(\sigma_{\alpha_\zeta} \circ s)^{k-\ell}(E_{\zeta,\eta}),$$

and hence we also have the required expression for the support of $\psi_{\zeta,\eta,\ell}$.
□

3.9. Basic exchangeable pairs

From 3.8.2, we see that $\psi_{\zeta,\eta}$ is a composition of exchanges between sets. Restricted to Z_m, each of these exchanges is just an involution. If $\alpha_\zeta * \overline{\alpha_\eta}$ is a simple closed loop up to homotopy, then $\psi_{\zeta,\eta}$ is a composition of disc exchanges, first between the two halves of $E_{\zeta,\eta}$ and then between two discs in each component of $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E_{\zeta,\eta})$ for increasing i.

An *exchangeable pair (for ζ)* is a notation for keeping track of inverse images of sets $E = E_{\zeta,\eta}$ under $(\sigma_{\alpha_\zeta} \circ s)^n$, and hence for describing the support of $\psi_{\zeta,\eta}$. We shall see later that a component of $(\sigma_{\alpha_\zeta} \circ s)^{-n}(E)$ coincides with a component of

$$s^{-n}(E) \cup (\sigma_\zeta \circ s)^{1-n}(s^{-1}\zeta).$$

In all cases that we shall consider, we shall have $E \subset D(L_3L_2)$ or $E \subset D(BC)$, and the intersection of ζ with the unit disc will be with $D(L_3L_2)$, or the union of $D(BC)$ and the single leaf of $L_{3/7}$ joining $e^{\pm 2\pi i(2/7)}$ respectively.

It follows any component C is coded by an *exchangeable pair for* ζ which is denoted by $v_1 \leftrightarrow v_2$, possibly with the words "top" or "bottom" attached to v_1 and v_2, where v_1E and v_2E are the components of $s^{-n}E$ within C, and "top" or "bottom" indicates that the component of $(\sigma_\zeta \circ s)^{1-n}(s^{-1}(\zeta))$ attached to (v_1E or v_2E) at a point on the upper or lower unit circle. We shall write $C = C(v_1 \leftrightarrow v_2, E)$ or $C(v_1 \leftrightarrow v_2)$ if it is clear what E is. Strictly speaking, we should include the path ζ in the notation, but in the cases which we consider, ζ is usually a path in $R_{m,0}$ for which $w_1'(\zeta)$ is known, and the words $w_j'(\zeta)$, which encode the unit-disc crossings of ζ, coincide with $w_j'(v_1) = w_j'(v_2)$, and hence ζ can be computed from v_1 (or v_2). At any rate, v_1 and v_2 are admissible of words of the same length n in the letters $\{L_j, R_j, BC, UC : 1 \leq j \leq 3\}$. Of course $C(v_1 \leftrightarrow v_2) = C(v_2 \leftrightarrow v_1)$, and so $v_1 \leftrightarrow v_2$ and $v_2 \leftrightarrow v_1$ are naturally equivalent. If $C = C(v_1 \leftrightarrow v_2)$ does not intersect ζ, then $(\sigma_\zeta \circ s)^{-1}(C) = s^{-1}(C)$. If in addition the left-most letters of v_1 and v_2 are the same, then the components of $(\sigma_\zeta \circ s)^{-1}(C) = s^{-1}(C)$ are $C(u_1 \leftrightarrow u_1v_2)$ and $C(u_2v_1 \leftrightarrow u_2v_2)$, where u_1 and u_2 are the letters such that u_1v_j and u_2v_j are admissible for one, and hence both, j.

The *length* of an exchangeable pair $v_1 \leftrightarrow v_2$ is the length $|v_1|$ of v_1 — which is equal to the length $|v_2|$ of v_2.

Basic exchangeable pairs are defined inductively. The only basic exchangeable pairs of length 1 are $L_3 \leftrightarrow R_3$ and $L_2 \leftrightarrow R_2$. We define $C = C(L_i \leftrightarrow R_i)$ to be the set obtained by joining $D(L_i)$ and $D(R_i)$ by an arc along the unit circle which joins a point on $\partial D(L_i)$ in the lower unit circle to a point in $\partial D(R_i)$ in the upper unit circle. Now let $n \geq 2$. Then $v_1 \leftrightarrow v_2$ is a basic exchange of length n if and only if a component of $s^{-n}(C)$ is the union of sets $D(v_1)$ and $D(v_2)$ and an arc joining them. Thus if $v_1 \leftrightarrow v_2$ is a basic exchange of length n, and v_1' and v_2' are the suffixes of v_1 and v_2 of length m for some $m < n$, then $v_1' \leftrightarrow v_2'$ is a basic exchange of length m. In particular the last letters of v_1 and v_2 are L_i and R_i for $i = 2$ or 3. If $i = 3$, we say that $v_1 \leftrightarrow v_2$ is *a basic exchange for* L_3 — because $s^{-1}(D(L_3)) = D(L_3^2) \cup D(R_3L_3)$ — and if $i = 2$ we say that $v_1' \leftrightarrow v_2'$ is *a basic exchange for* BC — because $s^{-1}(D(BC)) = D(L_2BC) \cup D(R_2BC)$.

It follows that a basic exchange $v_1 \leftrightarrow v_2$ is an exchange for ζ if, for any m with $1 < m < n$, the arc in $C(v_1' \leftrightarrow v_2')$ does not intersect ζ. It turns out that a basic exchange $v_1 \leftrightarrow v_2$ is automatically an exchange for $\zeta \in R_{m,0}$ if all the unit disc crossings of $\zeta \in R_{m,0}$ are in $D(L_3)$, and either v_1 and v_2 have no common letters in the the same positions, or there are common letters, and, among the common letters, if there is an occurrence of L_3, it is the first letter of each word. This becomes apparent from computing all the basic exchanges $v_1 \leftrightarrow v_2$. We now do this. This calculation was also carried out in [**22**]. If $v_1 \leftrightarrow v_2$ is a basic exchange, and the first letters of v_1 and v_2 are the same, or can admissibly be preceded by the same letter, then $uv_1 \leftrightarrow uv_2$ is a basic exchange for any word u such that uv_1 is admissible. Therefore we only need to compute those basic exchanges $v_1 \leftrightarrow v_2$ such that all the letters in the same positions of v_1 and v_2 are different. Also, of course, we only have to compute up to equivalence, that it, up to interchanging v_1 and v_2.

We start with basic exchangeable pairs for L_3 . The unique exchangeable pair of length one is

$$L_3(\mathrm{bot}) \leftrightarrow (\mathrm{top})R_3 \tag{3.9.0.10}$$

Since $L_3E \subset D(L_3^2L_2)$ and $R_3E \subset D(R_3L_3L_2)$, for any $\zeta \in R_{m,0}$, the only possible intersection between $C(L_3 \leftrightarrow R_3)$ and ζ is with the initial segment of ζ, before the first intersection with S^1. Up to homotopy preserving S^1 and Z_∞, we can assume that $C(L_3 \leftrightarrow R_3) \cap \{z : |z| > 1\}$ is close to the clockwise arc of the unit circle between R_3E and L_3E, mostly close to the lower unit circle, and not intersecting ζ. The only exchanges of length 2 are

$$L_3^2(\mathrm{top}) \leftrightarrow (\mathrm{bot})L_2R_3 \tag{3.9.0.11}$$

and

$$R_3L_3(\mathrm{bot}) \leftrightarrow (\mathrm{top})R_2R_3 \tag{3.9.0.12}$$

These are both basic. The only exchanges of length 3 are

$$R_3L_3^2(\mathrm{top}) \leftrightarrow (\mathrm{top})L_3L_2R_3, \tag{3.9.0.13}$$

$$L_3^3(\mathrm{bot}) \leftrightarrow (\mathrm{bot})R_3L_2R_3, \tag{3.9.0.14}$$

$$L_2R_3L_3(\mathrm{top}) \leftrightarrow (\mathrm{bot})L_1R_2R_3, \tag{3.9.0.15}$$

$$R_2R_3L_3(\mathrm{bot}) \leftrightarrow (\mathrm{top})R_1R_2R_3. \tag{3.9.0.16}$$

Of these, only the component represented by (3.8.2.4) might intersect ζ. If we assume, as will always be true, that $L_3L_2R_3E$ is disjoint from ζ – in fact it is always disjoint from $E \cup \zeta$– then this too is basic.

The following was computed in [**22**], and is easily checked. The exchanges are grouped together so that those in (3.9.1.1) are derived from (3.9.0.13), those in (3.9.1.2) are derived from (3.9.0.14) and those in (3.9.1.3) from (3.9.0.15), with (3.9.0.16) being the last one in the group (3.9.1.3).

LEMMA 3.9.1. *The complete list of basic exchangeable pairs $a \leftrightarrow b$ for ζ, for which the first letters of a and b are not the same, but for which the letters preceding a and b in any inverse image has to be the same, is as follows, where n is an integer ≥ 0 . In all cases, if one of a and b has all letters in the set $\{L_3, L_2, R_3\}$, then b has been chosen to have this property,* *and if both a and b have all letters in $\{L_3, L_2, R_3\}$ then $D(b)$ is chosen to be to the left of $D(a)$, if the exchange is along the bottom of the unit circle, and to the right if the exchange is along the top of the unit circle. Any other basic exchanges are represented by suffixes of these. In (3.9.1.1) to (3.9.1.3), the connection in C between $D(a)$ and $D(b)$ is between the nearest component of $\partial D(a) \cap S^1$ and $\partial D(b) \cap S^1$, while in (3.9.1.4) the connection is between the furthest component of $\partial D(a) \cap S^1$ and $\partial D(b) \cap S^1$.*

(3.9.1.1)

$$\begin{gathered}
L_2(UCL_1R_2)^nR_3L_3^2(\text{bot}) \leftrightarrow (\text{bot})L_3(L_3L_2R_3)^{n+1} \\
BCL_1R_2(UCL_1R_2)^nR_3L_3L_3(\text{bot}) \leftrightarrow (\text{bot})R_3L_2R_3(L_3L_2R_3)^{n+1} \\
R_1R_2(UCL_1R_2)^nR_3L_3L_3(\text{top}) \leftrightarrow (\text{top})R_2R_3(L_3L_2R_3)^{n+1}
\end{gathered}$$

(3.9.1.2)

$$\begin{gathered}
L_2(BCL_1R_2)^nR_3L_2R_3(\text{top}) \leftrightarrow (\text{top})L_3(L_3L_2R_3)^nL_3^3, \\
UCL_1R_2(BCL_1R_2)^nR_3L_2R_3(\text{top}) \leftrightarrow (\text{top})R_3L_2R_3(L_3L_2R_3)^nL_3^3, \\
R_1R_2(BCL_1R_2)^nR_3L_2R_3(\text{top}) \leftrightarrow (\text{top})R_2R_3(L_3L_2R_3)^nL_3^3, \\
,
\end{gathered}$$

$$\begin{gathered}
L_2(BCL_1R_2)^{n+1}R_3(\text{top}) \leftrightarrow (\text{top})L_3(L_3L_2R_3)^{n+1}L_3 \\
UCL_1R_2(BCL_1R_2)^nR_3(\text{top}) \leftrightarrow (\text{top})R_3L_2R_3(L_3L_2R_3)^nL_3 \\
R_1R_2(BCL_1R_2)^{n+1}R_3(\text{bot}) \leftrightarrow (\text{bot})R_2R_3(L_3L_2R_3)^{n+1}L_3, \\
R_2R_3L_3(\text{bot}) \leftrightarrow (\text{top})R_1R_2R_3
\end{gathered} \tag{3.9.1.3}$$

For a word u ending in L_3 or R_3, we define $i = i(u)$ to be the largest integer such that $w_i(uL_3L_2)$ is a proper prefix of uL_3L_2 — defining $i(u) = 0$ if $w_1(uL_3L_2) = uL_3L_2$. The exchanges $a \leftrightarrow b$ for which v can be chosen so that $i(vbL_3L_2) = i(vaL_3L_2) = 1$ and $w_1'(vbL_3L_2) \neq w_1'(vaL_3L_2)$, and both $D(va)$ and $D(vb)$ intersect the upper unit circle in $D(L_3L_2)$, are precisely those for which $v = v_0v_1$, where all letters of v_1 and a and b are in $\{L_3, L_2, R_3\}$ and v_0v_1 starts L_3L_2 and v_1 starts L_3 and all subsequent letters are in $\{L_2, R_3\}$, and $i(v_0L_3L_2) = v_0L_3L_2$. These are derived from the first lines of (3.9.1.1) and (3.9.1.2) and (3.9.1.3). The exchanges $v_1a \leftrightarrow v_1b$ are all of the form

$$\begin{gathered}
L_3(L_2R_3)^{2t+1}L_3^2L_2R_3(\text{bot}) \leftrightarrow (\text{bot})L_3(L_2R_3)^{2t+2}L_3^2, \\
L_3(L_2R_3)^{2t+2}(\text{bot}) \leftrightarrow (\text{bot})L_3(L_2R_3)^{2t}L_3^4.
\end{gathered} \tag{3.9.1.4}$$

For these exchanges we have

$$i(v_1b) = i(v_1a) = 1 = i(va) = i(vb),$$

$$w_1'(vaL_3L_2) = v_0w_1'(v_1aL_3L_2), \quad w_1'(vbL_3L_2) = v_0w_1'(v_1bL_3L_2),$$

and

$$w_1'(v_1a) \neq w_1'(v_1b).$$

In fact

$$\begin{gathered} w_1'(L_3(L_2R_3)^{2t+2}L_3L_2) = L_3(L_2R_3)^{2t}, \\ w_1'(L_3(L_2R_3)^{2t}L_3^5L_2) = L_3(L_2R_3)^{2t}L_3^2, \\ w_1'(L_3(L_2R_3)^{2t+1}L_3^2L_2R_3L_3^2L_2) = L_3(L_2R_3)^{2t}, \\ w_1'(L_3(L_2R_3)^{2t+2}L_3^3L_2) = L_3(L_2R_3)^{2t+3}L_3. \end{gathered}$$

The exchanges $a \leftrightarrow b$ for which v can be chosen so that vb starts with L_3L_2 with

$$i(vbL_3L_2) = i(vaL_3L_2) = 1, \;\; w_1'(vbL_3L_2) = w_1'(vaL_3L_2), \;\; w_1(vbL_3L_2) \neq w_1(vaL_3L_2),$$

and such that $D(vb)$ intersects the upper unit circle, and $D(va)$ does not, are obtained from the second lines of (3.9.1.1) and (3.9.1.2) and the first two lines of (3.9.1.3). They are as follows, where n is any integer ≥ 0

(3.9.1.5)

$$\begin{gathered} L_3(L_2R_3)^{2t}L_2BCL_1R_2(UCL_1R_2)^nR_3L_3^2(\mathrm{bot}) \leftrightarrow (\mathrm{bot})L_3(L_2R_3)^{2t+2}(L_3L_2R_3)^{n+1} \\ (BCL_1R_2)^{n+1}R_3L_2R_3(\mathrm{bot}) \leftrightarrow (\mathrm{bot})(L_3L_2R_3)^{n+1}L_3^3, \\ (BCL_1R_2)^{n+1}R_3(\mathrm{top}) \leftrightarrow (\mathrm{bot})(L_3L_2R_3)^{n+1}L_3 \end{gathered}$$

The first of these can be prefixed by any word v such that $w_1(vL_3L_2) = vL_3L_2$. The last two can be prefixed by (v_1L_2, v_1L_3) or by $(v_2UCL_1R_2, v_2R_3L_2R_3)$ for any v_1 and v_2 such that $w_1(v_1L_3^2L_2) = v_1L_3^2L_2$ and $w_1(v_2R_3L_2R_3L_3L_2) = v_2R_3L_2R_3L_3L_2$.

The pairs (a, b) as in (3.9.1.4) are precisely those for which $w_1'(vbL_3L_2) = y_2$ and $w_1'(vaL_3L_2) = y_1$ with $y_1 \neq y_2$, and such that the set U^{y_2} contains $D(vbv_2)$ if and only if $w_1'(v_2) > w_1'(y_2)$, and contains $D(vav_2)$ if and only if $w_1'(v_1) \leq w_1'(y_2)$, while the set U^{y_1} contains $D(vav_2)$ if and only if $w_1'(v_2) > y_1$, and contains $D(vbv_2)$ if and only if $w_1'(v_1) \leq y_1$. It follows that $D(vav_1) \subset U^{y_1} \cap U^{y_2}$ if $y_1 < w_1'(v_1) \leq y_2$ and that $D(vbv_1)$ is contained in neither U^{y_1} nor U^{y_2} if $y_1 < w_1'(v_1) \leq y_2$. In fact, $D(vbv_2)$ is not contained in any U^y if $y_1 < w_1'(v_2) \leq y_2$.

3.10. Exchangeable pairs for BC

For convenience, we list the basic exchangeable pairs for BC. These are obtained by changes to the later letters in the pairs above and are as follows:

$$\begin{gathered} BCL_1R_2(\mathrm{bot}) \leftrightarrow (\mathrm{bot})L_3^2L_2, \\ L_3^3L_2(\mathrm{top}) \leftrightarrow (\mathrm{top})L_2BCL_1R_2. \end{gathered} \tag{3.10.0.6}$$

$$L_3(L_3L_2R_3)^nL_3^2L_2(\mathrm{top}) \leftrightarrow (\mathrm{top})L_2(BCL_1R_2)^{n+1} \tag{3.10.0.7}$$

for some $n \geq 1$.

$$\begin{aligned}&R_2R_3L_3^2L_2 \leftrightarrow R_1R_2BCL_1R_2,\\ &R_3L_2R_3L_3^2L_2(\text{top}) \leftrightarrow (\text{top})UCL_1R_2BCL_1R_2.\end{aligned} \tag{3.10.0.8}$$

$$\begin{aligned}&R_2R_3L_2 \leftrightarrow R_1^2R_2),\\ &R_3L_2R_3L_3 \leftrightarrow UCL_1R_2R_3),\\ &L_3(L_3L_2R_3)^nL_2(\text{top}) \leftrightarrow (\text{top}))L_2(BCL_1R_2)^{n-1}BCL_1R_1R_2,\\ &R_2R_3(L_3L_2R_3)^nL_2 \leftrightarrow R_1R_2(BCL_1R_2)^{n-1}BCL_1R_1R_2),\\ &R_3L_2R_3(L_3L_2R_3)^nL_2(\text{top}) \leftrightarrow (\text{top})UCL_1R_2(BCL_1R_2)^n.\end{aligned} \tag{3.10.0.9}$$

3.11. Exchangeable pairs for (ζ, η)

Now we return to the question of when a component C of $s^{2-i}(\sigma_\zeta \circ s)^{-2}(E)$ coincides with a component of $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$, for $E = E_{\zeta,\eta}$, and can therefore be coded by basic exchangeable pairs. Coincidence fails when C is in $(\sigma_{\alpha_\zeta} \circ s)^{j-i}(C')$ for a component C' of $s^{2-j}(\sigma_\zeta \circ s)^{-2}(E))$ for some $j < i$ and $C' \cap \alpha_\zeta \neq \emptyset$. Now $C' = C_1' \cup C_2' \cup \ell_r$ where C_1' and C_2' are components of $s^{-j}(E)$ and $r = 1$ or 2, and ℓ_1 and ℓ_2 are the component of $s^{2-j}((\sigma_\zeta \circ s)^{-1}(s^{-1}(\zeta))$. In our cases it will always be true that

$$(C_1' \cup C_2') \cap \alpha_\zeta = \emptyset$$

and

$$\ell \cap \alpha_\zeta = \ell \cap \zeta.$$

This will be proved in 4.12.1. This means that components of $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ will be components of $(\sigma_{\zeta_0} \circ s)^{2-i}(\sigma_\zeta \circ s)^{-2}(E))$. We shall call these *exchangeable pairs for* (ζ, η).

This coding, as before, is inductive. Exchangeable pairs for (ζ, η) (and L_3 or BC) are the same as basic exchangeable pairs for length ≤ 3, and indeed all the exchangeable pairs $a \leftrightarrow b$ for $(\zeta, \eta$ for which corresponding letters in a and b are all different are the same as the basic ones. In general the exchangeable pairs $a \leftrightarrow b$ of length n are of the form $xa_1 \leftrightarrow xb_1$ for exchangeable pairs $a_1 \leftrightarrow b_1$ and letters x such that xa_1 is an admissible word. (In that case xb_1 is admissible also as we are assuming that a_1 and b_1 can be preceded by the same letter.) The exception is when the component of $s^{2-n}(\ell_r))$ corresponding to $a_1(\text{top}) \leftrightarrow (\text{top})b_1$ intersects ζ. The inverse image exchangeable pairs (for L_3)are then

$$L_3a_1(\text{bot}) \leftrightarrow R_3b_1, \quad L_3b_1(\text{bot}) \leftrightarrow R_3a_1$$

Then the coding proceeds inductively.

Relatively few letters can be different in an exchangeable pair $a \leftrightarrow b$. For a basic exchangeable pair the only pairs for which more than the last 5 letters differ are

- $UCL_1R_2R_3L_2R_3 \leftrightarrow R_3L_2R_3L_3^3$,
- pairs $u_1a_1 \leftrightarrow u_2b_1$ for which a_1 (and b_1) has length ≤ 4 and the first two letters of (u_1, u_2) are (L_2BC, L_3^2) or (UCL_1, R_3, L_2) or (R_1R_2, R_2R_3), the last letters are (R_2, R_3) and the rest is a suffix of $((BCL_1R_2)^n, (L_3L_2R_3)^n)$.

The exchangeable pairs for (ζ, η) are the same except that some more letters in a and b can be different. If $w_k'(\beta_1) = u^k$ then any exchangeable pair for (ζ, η) is of the form

$$u_ra_ru_{r-1}a_{r-1}\cdots u_1a_1 \leftrightarrow u_rb_ru_{r-1}b_{r-1}\cdots u_1b_1$$

where

- $a_i \leftrightarrow b_i$ is a basic exchangeable pair
- $D(u^{k_i})$ is between $D(u_ia_i)$ and $D(u_ib_i)$ for some k_i and $2 \leq i \leq r$.

We have the following extension to 3.9.1, which follows from the definition of the $w_j(u)$ and $w_j'(u)$.

LEMMA 3.11.1. *Let $v \leftrightarrow v'$, be an exchangeable pair for (ζ, η). For any prefixes u and u' of vL_3L_2 and $v'L_3L_2$ with $|u| = |u'|$, we have the following. The largest integer $i = i(u)$ such that $w_i(u)$ is a proper prefix of u is the same as the largest integer k such that $w_i(u')$ is a proper prefix of u'. Write*

$$v = u_1a_1\cdots u_ta_t, \quad v' = u_1b_1\cdots u_tb_t.$$

If $k(u_1L_3L_2) \geq 1$ then

$$w_1'(u_1a_1) = w_1'(u_1b_1) = w_1'(v) = w_1'(v'),$$

and this is also true if $i(u_1a_1) \geq 1$, except for certain choices of (a_1, b_1) as specified in (3.9.1.5).

3.12. The homeomorphisms $\psi_{m,q}$.

As in [**23**], we let $\beta_q : [0,1] \to \overline{\mathbb{C}}$ be the capture path which crosses the unit circle at $e^{2\pi iq}$ and ends at the point $\beta_q(1) \in Z(s)$ in the unique gap of $L_{3/7}$ with $e^{2\pi iq}$ in its boundary. Thus, β_q is defined for all q such that $e^{2\pi iq}$ is in the boundary of a gap of $L_{3/7}$. If $\beta_q(1)$ is strictly preperiodic, then β_q is a type III capture map and $\sigma_{\beta_q} \circ s$ is type III branched covering. If $\beta_q(1)$, is periodic then we define $\zeta_q : [0,1] \to \overline{\mathbb{C}}$ to be the unique path such that $s \circ \zeta_q = \beta_q$ and $\zeta_q(1)$ is periodic under s. In this case $\sigma_{\zeta_q}^{-1} \circ \sigma_{\beta_q} \circ s$ is a type II branched covering and β_q is a type II capture path (by definition). An example is given by $q = \frac{2}{7}$. This is the most important example for us.

In 3.3 of [**23**] we gave a proof (not new in any essential way, of course) that

$$\sigma_{\zeta_{5/7}}^{-1}\circ\sigma_{\beta_{5/7}}\circ s\simeq_{\psi_{0,2/7}}\sigma_{\zeta_{2/7}}^{-1}\circ\sigma_{\beta_{2/7}}\circ s$$

We then used $\psi_{0,2/7}$ to define a sequence of homeomorphisms $\psi_{m,2/7}$ and a sequence of closed loops $\alpha_{m,2/7}$, for all $m\geq 0$, and satisfying the following properties.

1. $\alpha_{m,2/7}$ is an arbitrarily small perturbation of $\beta_{2/7}*\psi_{m,2/7}(\overline{\beta_{5/7}})$ which bounds a disc disjoint from the common endpoint 0 of $\beta_{2/7}$ and $\beta_{5/7}$, for all $m\geq 0$.
2. $\psi_{m+1,2/7}=\xi_{m,2/7}\circ\psi_{m,2/7}$ for all $m\geq 0$.
3. $\psi_{m+1,2/7}$ and $\psi_{m,2/7}$ are isotopic relative to Y_m for all $m\geq 0$ and ξ_m is isotopic to the identity relative to Y_m for all $m\geq 0$.
4. $\sigma_{\alpha_{m,2/7}}\circ s\circ\psi_{m+1,2/7}=\psi_{m,2/7}\circ s$ and and $\sigma_{\beta_{2/7}}\circ s\circ\xi_{m+1,2/7}=\xi_{m,2/7}\circ\sigma_{\beta_{2/7}}$ for all $m\geq 1$.

In the definition of $\psi_{m,2/7}$ and $\alpha_{m,2/7}$ in [**22**], we started with the definition of $\psi_{0,2/7}$. But we might as well start from $\psi_{2,2/7}$ and $\xi_{1,2/7}$, because then 1 to 4 can be used to defined $\psi_{m,2/7}$ and $\xi_{m,2/7}$ and $\alpha_{m,2/7}$ inductively for all $m\geq 2$. (Of course, this will also give the definition of $\psi_{1,2/7}$, as $\psi_{2,2/7}=\xi_{1,2/7}\circ\psi_{1,2/7}$.) The homeomorphism $\psi_{2,2/7}$ is a composition of anticlockwise Dehn twists round the boundaries of four topological discs, two larger disjoint discs and two smaller discs, one contained in each of the others. The two larger discs are called D_+ and D_-, both contained in the unit disc, symmetrically placed about the origin, with D_+ bounded by the leaf of $L_{3/7}$ — just a vertical line up to isotopy preserving Z — joining $e^{2\pi i(5/14)}$ and $e^{2\pi i(9/14)}$ – and D_- is its symmetric copy bounded by the leaf joining $e^{2\pi i(1/7)}$ and $e^{2\pi i(6/7)}$. The two smaller discs are the components of $s^{-1}(D_-)$. The homeomorphism $\xi_{1,2/7}$ is the composition of anticlockwise Dehn twist round the boundaries of the components of $s^{-1}(D_{-,2/7})$, and of a clockwise twist in an annulus $A_{1,2/7}$. This annulus bounds a disc disjoint from ∞ and containing D_+. Its intersection with the unit disc is the set bounded by one vertical leaf joining $e^{2\pi i(2/7)}$ and $e^{2\pi(5/7)}$, and one joining $e^{2\pi i(5/14)}$ and $e^{2\pi i(9/14)}$. A clockwise twist in the annulus means, up to isotopy, a clockwise twist round the outer boundary composed with an anticlockwise twist round the inner boundary.

From the definitions, it will be seen that $\psi_{2,2/7}$ is the identity on $\beta_{5/7}$ and hence $\alpha_{2/2/7}$ is an arbitrarily small perturbation of $\beta_{2/7}*\overline{\beta_{5/7}}$. For any path γ based at ∞ we also have

$$\alpha_{m,2/7}*\psi_{m,2/7}(\gamma)=\beta_{2/7}*\psi_{m,2/7}(\overline{\beta_{5/7}}*\gamma).$$

For $\gamma \in \mathcal{Z}(3/7,+,0)$, $\psi_{2,2/7}$ is the identity on $\overline{\beta_{5/7}} * \gamma$ and hence for all $m \geq 3$,

$$\alpha_{m,2/7} * \psi_{m,2/7}(\gamma) = \beta_{2/7} * \xi_{m-1,2/7} \circ \cdots \circ \xi_{2,2/7}(\overline{\beta_{5/7}} * \gamma).$$

In order to proceed further we need to give some more information about the homeomorphisms $\xi_{i,2/7}$ for $i \geq 2$. The support of $\xi_{2,7}$ is a union of annuli $A_{i,2/7} \cup C_{i,2/7}$ where

$$A_{i,2/7} = (\sigma_{\beta_{2/7}} \circ s)^{-1}(A_{i-1,2/7})$$

for all $i \geq 2$ and

$$C_{i,2/7} = (\sigma_{\beta_{2/7}} \circ s)^{-1}(C_{i-1,2/7})$$

for all $i \geq 1$.

Previously in this chapter, we have used homeomorphisms which are compositions of disc exchanges. The homeomorphisms $\xi_{i,2/7}$ are a bit different, because each component of the support is a topological annulus rather than a topological disc. Within each annulus of $A_{i,2/7}$, components of intersection with the unit disc are cyclically permuted by $\xi_{2,2/7}$, up to isotopy preserving Z_m, for any fixed $m \geq i$, with each component of intersection sent to the next one in the clockwise direction. Within each annulus of $C_{i,2/7}$ the components of

$$s^{1-i}(\{z : |z| \leq 1\} \setminus \partial C_1 \cap \{z : |z| < 1\})$$

are cyclically permuted, with each one sent to the next one round in an anticlockwise direction. It is therefore possible to code the action of these homeomorphisms using words. This was done in section 7.2 of [**23**].

We have already defined $A_{1,2/7}$. The set $C_{0,2/7}$ is simply the complement in $D_{-,2/7}$ of a sufficiently small disc neighbourhood of $s(v_1)$ that the intersection of the neighbourhood with Z is simply $s(v_1)$. The sets $A_{i,2/7}$ and $C_{i,2/7}$ therefore have empty intersection with $Z_i \cup \{\infty\} = Y_i$, but do intersect Z_{i+1}. The branched covering $\sigma_{\beta_{2/7}} \circ s$ is of degree one or two on each component of $A_{i,2/7}$ and $C_{i,2/7}$, and both are possible.

We now give some pictures of these sets.

The permutations of sets $D(w)$ in these annuli include the following. The clockwise twist in $A_{1,2/7}$ preserves the single component of intersection $D(L_3)$ with the unit disc, up to isotopy preserving Z_m, for any fixed m. In $A_{2,/7}$ the permutation of components of intersection is indicated by:

$$L_3^2 \to (\text{top})R_3L_3 \to (\text{bot})L_3^2.$$

Here, "top" immediately preceding the word indicates a component of $A_{2,2/7} \cap \{z : |z| > 1$ ending on the top of the unit circle and "bot" immediately preceding the word indicates component of $A_{2,2/7} \cap \{z : |z| > 1$ ending on the bottom of the unit circle. In fact, in this example, one component of



FIGURE 2. $A_{1,2/7}$ and $C_{0,2/7}$



FIGURE 3. $A_{2,2/7}$



FIGURE 4. $A_{3,2/7}$



FIGURE 5. $A_{4,2/7}$



FIGURE 6. $A_{5,1,2/7}$

$A_{2,2/7} \cap \{z : |z| > 1$ is above the unit circle and one is below. Similarly, the action of $\xi_{3,2/7}$ in $A_{3,2/7}$ is indicated by

$$L_2R_3L_3(\text{top}) \rightarrow (\text{top})L_3^3\text{bot} \rightarrow (\text{top})R_2R_3L_3(\text{bot})$$
$$\rightarrow (\text{bot})R_3L_3^2(\text{top}) \rightarrow \text{bot}L_2R_3L_3.$$

For the two components of $A_{4,2/7}$, which we call $A_{4,1,2/7}$ and $A_{4,2,1/7}$, we have

$$L_1R_2R_3L_3(\text{top}) \rightarrow (\text{top})L_2R_3L_3^2(\text{bot}) \rightarrow \text{top})L_3L_2R_3L_3(\text{bot})$$
$$\rightarrow (\text{bot})L_3^4(\text{top}) \rightarrow (\text{bot})L_1R_2R_3L_3,$$



FIGURE 7. $A_{5,2,2/7}$



FIGURE 8. $A_{6,2,2/7}$

$$R_1R_2R_3L_3(\mathrm{bot}) \to (\mathrm{bot})R_2R_3L_3^2(\mathrm{top}) \to (\mathrm{bot})R_3L_2R_3L_3(\mathrm{top})$$
$$\to (\mathrm{top})R_3L_3^3(\mathrm{bot}) \to (\mathrm{top})R_1R_2R_3L_3.$$

For the single component $A_{5,1,2/7}$ of $A_{4,1,2/7}$, the action of $\xi_{5,2/7}$ is indicated by

$$(\mathrm{bot})BCL_1R_2R_3L_3 \to (\mathrm{bot})L_3L_2R_3L_3^2(\mathrm{top}) \to (\mathrm{top})R_3L_3L_2R_3L_3(\mathrm{bot}) \to$$
$$(\mathrm{bot})R_3L_3^4 \to (\mathrm{top})UCL_1R_2R_3L_3 \to (\mathrm{top})R_3L_2R_3L_3^2 \to (\mathrm{bot})$$
$$L_3^2L_2R_3L_3 \to (\mathrm{top})L_3^5 \to (\mathrm{bot})BCL_1R_2R_3L_3.$$

Since $\beta_{2/7}$ does not intersect any of the components of $A_{4,2,2/7} \cap \{z : |z| \geq 1\}$, the components of $(\sigma_{\beta_{2/7}} \circ s)^{-1}(A_{4,2,2/7})$ are actually components of $s^{-1}(A_{4,2,2/7})$. In fact since none of the points $e^{2\pi ir/7}$ for $r \in \{1, 2, 4\}$ are in any interval of the unit circle bounded by a component of $A_{4,2,2/7} \cap$



FIGURE 9. $C_{2,2/7}$

$\{z : |z| \geq 1\}$, all components of $A_{i,2/7}$ in $(\sigma_{\beta_{2/7}} \circ s)^{4-i}(A_{4,2,2/7})$, for all $i \geq 5$, are actually components of $s^{4-i}(A_{4,2,2/7})$. In fact all components of $s^{-p}(A_{i,2/7} \cap \{z : |z| \geq 1\}$ are disjoint from $\beta_{2/7}$ for all $p \geq 0$ except for those components represented by:

$$(L_3L_2R_3)^nL_3^2(\text{top}) \to (\text{top})(R_3L_3L_2)^nR_3L_3,$$

if $i = 3n + 2$ for any $n \geq 0$;

$$(R_3L_3L_2)^nR_3L_3^2(\text{top}) \to (\text{bot})(L_2R_3L_3)^{n+1}$$

if $i = 3n + 3$;

$$(L_2R_3L_3)^nL_3(\text{bot}) \to (\text{top})L_3(L_2R_3L_3)^n$$

if $i = 3n + 4$. For any other component represented by $Y \to Y'$, any component of any $A_{j,2/7} \cap \{z : |z| \geq 1\}$ in the backward orbit is represented by $ZXY \to ZX'Y$, where (X, X') is one of the following, or obtained from one of these by adjoining BC, L_3 or (BCL_1, L_3L_2), or a suffix of one of these if Z is empty:

$$(R_1R_2, R_2R_3),\ (UCL_1R_2(BCL_1R_2)^p, R_3L_2R_3(BCL_1R_2)^p)$$

for any $p \geq 0$. So, altogether, these give the same pairs as the basic exchanges in 3.9.1.

As for components of $C_{i,2/7}$, we have a similar pattern. Here are some pictures of some of the sets $C_{i,2/7}$.

The homemorphism $\xi_{2,2/7}$ restricted to $C_{2,2/7}$ is represented by:

$$UC \to BC \to UC.$$

The preimage $C_{3,2/7}$ of $C_{2,2/7}$ is connected and $\xi_{3,2/7}$ restricted to $C_{3,2/7}$ is represented by:

$$R_2BC \to R_2UC \to L_2BC \to L_2UC \to R_2BC.$$

This, and the following representations, should be read in conjunction with the pictures. In $C_{3,2/7}$, for example, the piece of $C_{3,2/7}$ between $D(R_2BC)$ and $D(R_2UC)$ is inside the unit circle, along the right boundary of $D(R_2C)$,



FIGURE 10. $C_{3,2/7}$



FIGURE 11. $C_{5,2/7}$

while the piece of $C_{3,2/7}$ between $D(L_2UC)$ and $D(R_2UC)$ traces an approximate diagonal path from the lower unit circle (adjacent to $D(L_2UC)$, then enters the unit disc again on the upper boundary of $D(R_2C)$ and travels down the left vertical boundary of $D(R_2C)$ to $D(R_2BC)$. The set $C_{4,2/7}$ has two components $C_{4,1,2/7}$ and $C_{4,2,2/7}$. The homeomorphism $\xi_{4,2/7}$ restricted to $C_{4,1,2/7}$ and $C_{4,2,2/7}$ is represented by, respectively:

$$L_1R_2BC \to L_1R_2UC \to L_3L_2BC \to L_3L_2UC \to L_1R_2BC,$$

$$R_1R_2BC \to R_1R_2UC \to R_3L_2BC \to R_3L_2UC \to R_1R_2BC.$$

The preimage of $C_{4,1,2/7}$ in $C_{5,2/7}$ has a single component $C_{5,1,2/7}$ and the homeomorphism $\xi_{5,2/7}$ restricted to $C_{5,1,2/7}$ is represented by:

$$BCL_1R_2BC \to BCL_1R_2UC \to L_3^2L_2BC \to L_3^2L_2UC \to$$

$$UCL_1R_2BC \to UCL_1R_2UC \to R_3L_3L_2BC \to R_3L_3L_2UC \to BCL_1R_2BC.$$

None of the components of $C_{4,2,2/7} \cap \{z : |z| \geq 1\}$ intersect $\beta_{2/7}$ and neither do any of the components of $s^{-n}(C_{4,2,2/7} \cap \{z : |z| \geq 1\})$ for any $n \geq 0$. Therefore the components of $(\sigma_{\beta_{2/7}} \circ s)^{-n}(C_{4,2,2/7} \cap \{z : |z| \geq 1\})$ are components of $s^{-n}(C_{4,2,2/7} \cap \{z : |z| \geq 1\})$. The same is true for all but two of the components of $C_{4,1,2/7} \cap \{z : |z| \geq 1\}$. The exceptional pair are the components represented by

$$L_1R_2UC \to L_3L_2BC$$

and

$$L_3L_2UC \to L_1R_2BC,$$

where both the components intersect the unit circle in $D(L_1R_2UC)$ and $D(L_3L_2UC)$, and there are adjacent components of $C_{4,1,2/7}$ in the boundaries of $D(L_1R_2C)$ and $D(L_3L_2C)$ connecting $D(L_1R_2UC)$ to $D(L_1R_2BC)$, and $D(L_3L_2UC)$ to $D(L_3L_2BC)$. Of the preimages of these in $C_{5,1,2/7}$, only those represented by

$$UCL_1R_2UC \to R_3L_3L_2BC$$

and

$$R_3L_3L_2UC \to UCL_1R_2BC$$

have backward orbits which intersect $\beta_{2/7}$. Similarly in $C_{i,,2/7} \cap \{z : |z| \geq 1\}$, the only components whose backward orbits intersect $\beta_{2/7}$ represented by

$$(R_2UCL_1)^nR_2UC \to (L_2R_3L_3)^nL_2BC$$

and

$$(L_2R_3L_3)^nL_2BC \to (R_2UCL_1)^nR_2UC$$

if $i = 3n + 3$, or

$$(L_1R_2UC)^{n+1} \to (L_3L_2R_3)^nL_3L_2BC$$

and

$$(L_3L_2R_3)^nL_3L_2BC \to (L_1R_2UC)^{n+1}$$

if $i = 3n + 4$, or

$$(UCL_1R_2)^nUC \to (R_3L_3L_2)^nBC$$

and

$$(R_3L_3L_2)^nBC \to (UCL_1R_2)^nUC$$

if $i = 3n + 5$.

The pre-images in $C_{i+n,2/7}$ of any component of $C_{i,2/7} \cap \{z : |z| \geq 1\}$ represented by $Y \to Y'$ whose backward orbit does not intersect $\beta_{2/7}$, that is, any but the cycle given above, are obtained by adjoining Z to $Y \to Y'$ if the first letters of Y and Y' are the same, or are in the set $\{R_3, BC, UC\}$, and adjoining ZX and ZX' to $Y \to Y'$, giving $ZXY \to ZX'Y'$, where

(X, X') is of one of the following forms, or a suffix of one of these if Z is empty:

$$(R_1, R_2),\ (UC(L_1R_2BC)^pL_1, R_3(L_2R_3L_3)^pL_2),$$

$$(L_2(BCL_1R_2)^pBCL_1R_1, L_3(L_3L_2R_3)^{p+1})$$

for any $p \geq 0$. These are the same forms as for the basic exchangeable pairs for BC in 3.10.

3.13. $\psi_{m,q}$

We saw in 3.3 of [**22**] that the equivalence between the type III captures $\sigma_{\beta_{1-q}}$ and $\sigma_{\beta_q} \circ s$ can be realised similarly for any $q \in (\frac{2}{7}, \frac{1}{3})$ such that β_q is a type III capture path, that is, $e^{2\pi iq}$ is on the boundary of a gap of $L_{3/7}$. The homeomorphisms $\psi_{m,q}$ and $\xi_{m,q}$ and loops $\alpha_{m,q}$ are defined in the same inductive fashion as the homeomorphisms $\psi_{m,2/7}$ and $\xi_{m,2/7}$ and loops $\alpha_{m,2/7}$ in 3.12. The support of $\xi_{m,q}$ is again a union of annuli

$$A_{m,q} \cup C_{m,q} = (\sigma_{\beta_q} \circ s)^{2-m}(A_{2,q} \cup C_{2,q})$$

and $\psi_{1,q}$ is again a composition of anticlockwise Dehn twists round the boundaries of discs $D_{\pm,q}$ and $s^{-1}(D_{-,q}) \subset D_{+,q} \cup D_{-,q}$.

3.14. First stage in the proof of Theorem 2.8

We are now ready to prove parts of Theorem 2.8. We prove the base of the induction and part of the inductive step. We cannot prove the whole of the inductive step at this stage, however. That will have to wait until Chapter 4. Nevertheless, the part of the inductive step that can be proved holds for $\mathcal{Z}_m(3/7, +, p)$ for any $p \geq 0$. We have the following.

Lemma 3.14.1. *Let* $\beta \in \mathcal{Z}_m(3/7, +, p)$. *Let*

$$((\omega_{2i-1}, \omega_{2i}) : 1 \leq i \leq n(\beta))$$

and

$$(\gamma_i : 1 \leq i \leq n(\beta))$$

be the unique sequences of, respectively, adjacent pairs in Ω_m *and elements of* $\pi_1(V_{3,m}, a_1)$, *such that* β *successively crosses the arcs* $\overline{\rho(\gamma_i * \omega_{2i-1})} * \rho(\gamma_i * \omega_{2i-1})$, *with* $\beta = \rho(\gamma_{n(\beta)} * \omega_{2n(\beta)})$. *Then the following hold.*

1. $(\rho(\omega_1), \rho(\omega_2)) = (\beta_1, \beta_2) \in R_{m,p}$ *if* $\beta \in \mathcal{Z}_m(+, +, p)$ *and* $(\rho(\omega_1), \rho(\omega_2) = (\beta_1', \beta_2') \in R'_{m,p}$ *if* $\beta \in \mathcal{Z}_m(+, -, p)$.

2. *Let $\beta \in \mathcal{Z}_m(3/7, +, +, p)$. Suppose that it is true that, for $j \leq k$, there are adjacent pairs $(\beta_{2j-1}, \beta_{2j})$ in $R_{m,p}$ matched with $(\beta'_{2j-1}, \beta'_{2j})$ in $R'_{m,p}$, and*

$$(\rho(\omega_{2j-1}), \rho(\omega_{2j-1})) = \begin{cases} (\beta_{2j-1}, \beta_{2j}) \text{ if } j \text{ is even,} \\ (\beta'_{2j-1}, \beta'_{2j}) \text{ if } j \text{ is odd.} \end{cases}$$

   Then $(\omega_{2k+1}, \omega_{2k+2})$ is an adjacent pair in R_m if k is even and in R'_m if k is odd.
3. *An analogous statement holds for $\beta \in \mathcal{Z}_m(3/7, +, -, p)$, with odd and even interchanged.*

3.14.2. Remarks. The inductive step in 2 (and 3) is not quite strong enough to deduce the full statement of Theorem 2.8. The proof would be complete if R_m and R'_m could be replaced by $R_{m,p}$ and $R'_{m,p}$. It has only proved to be possible to do that for $p = 0$, because of lack of detailed knowledge about $R_{m,p}$ for $p > 0$. In any case, the completion of the proof of Theorem 2.8 will have to wait until Chapter 4.

3.14.3.

Proof. The paths in $R_{m,p}$ have never been described in detail for $p > 0$, although the set of endpoints lie in a set U^p which is described in Theorem 2.10 of [**22**]. We now present the only facts which we shall need. All statements are up to homotopy preserving Y_∞.

- All unit disc crossings of paths in R_m and the last component of intersection with the unit disc, are either along leaves of $L_{3/7}$ or in gaps of $L_{3,7}$. The same is true for paths in R'_m.
- The first unit-disc-crossings of paths in $R_{m,p}$ (apart from the path $\beta_{q_{p+1}}$) start at a point $e^{2\pi it}$ for $t \in [q_p, q_{p+1})$.
- The sets U^p are not all disjoint, not all points in $U^p \cap Z_m(s)$ are second endpoints of paths in $R_{m,p}$, and it is possible for at most two different paths in $R_{m,p}$ to have the same endpoint. However, each path in R_m is uniquely determined by its first unit-disc crossing and its final endpoint.

Suppose first that $\beta \in \mathcal{Z}_m(3/7, +, +, p)$, that is, the S^1 crossing point is at $e^{2\pi ix}$ for $x \in (q_p, q_{p+1})$ for some $p \geq 0$. We do not need to consider $x = q_p$ for any p because $\beta_{q_p} \in R_{m,p}$.

First, we prove 1. By definition, a capture path has only one intersection with the unit circle before entering the gap of $L_{3/7}$ containing the second endpoint of β. So the first essential intersections with S^1 of the paths $\rho(\gamma_i * \omega_{2i-1})$ and $\rho(\gamma_i * \omega_{2i})$ must be distinct, and opposite sides of the intersection with S^1 of β, although one of them could coincide with this. If

$\omega_i \in \Omega_m(a_0)$ (or $\omega_i \in \Omega_m(\overline{a_0})$) for at least one of $i = 1$ or 2, then both $\rho(\omega_1)$ and $\rho(\omega_2)$ have first essential S^1-crossing at $e^{2\pi i(6/7)}$ (or $e^{2\pi i(1/7)}$). This is is incompatible with β crossing $\overline{\rho(\omega_1)} * \rho(\omega_2)$. The paths in $\Omega_m(a_1, -)$ are ruled out more simply as candidates for $\rho(\omega_1)$ or $\rho(\omega_2)$, because these are simply the paths $\mathcal{Z}_m(3/7, -)$, which have first and only crossings of S^1 at points $e^{2\pi ix}$ for $x \in (-\frac{1}{7}, \frac{1}{7})$. Similarly we can rule out $\Omega_m(a_1, -)$. So we must have $\omega_i \in \Omega_m(a_1, +)$, and $\rho(\beta_i) \in R_{m,p}$ for $i = 1$ and 2, in order to have β crossing $\overline{\beta_1} * \beta_2$ before the first S^1-crossing points of those paths.

Now we consider the inductive step. So we assume the inductive hypothesis of the lemma, which is assuming that suppose that 1 of 2.8 is true for β_i for $i \leq 2k$, for some $k \geq 1$. Then we consider what $(\omega_{2k+1}, \omega_{2k+2})$ can be. First, let k be even. Then the inductive hypothesis gives $\rho(\gamma_{k+1}) = \alpha_{1,2k-1}$ in the notation of 2.8, that is, $\alpha_{1,2k-1}$ is an arbitrarily small perturbation of

$$\alpha_{1,2k-1} = \alpha_{1,3} * \psi_{1,3}(\alpha_{5,7} * \cdots * \psi_{1,2k-5}(\alpha_{2k-3,2k-1})$$

where $\alpha_{4j-3,4j-1}$ is an arbitrarily small perturbation of

$$\beta_{4j-3} * \psi_{4j-3}(\overline{\beta'_{4j-3}} * \beta'_{4j-1}) * \psi_{4j-3,4j-1}(\overline{\beta_{4j-1}}).$$

This means that the number of vertical intersections of $\rho(\gamma_{k+1}) = \alpha_{1,2k-1}$ with the unit disc is even. So $\psi_{1,2k-1}(\rho(\omega_{2k+1}))$ and $\psi_{1,2k-1}(\rho(\omega_{2k+2}))$ must first cross the unit disc in the same direction as β, that is, across the upper unit circle first. Also, we now know that $\psi_{1,2k-1}$ is a composition of disc exchanges, and we have a complete list of disc exchanges, in particular of those which intersect the unit circle within the set $\{e^{2\pi it} : t \in [\frac{2}{7}, \frac{1}{3})\}$. None of these supports can cancel first S^1-crossing of $\rho(\omega_{2k+1})$ or $\rho(\omega_{2k+2})$ at $e^{2\pi ix}$ for $x \notin [\frac{2}{7}, \frac{1}{3})$. So we must have $(\rho(\gamma_{2k+1}, \gamma_{2k+2}) = (\beta_{2k+1}, \beta_{2k+2})$ where $(\beta_{2k+2}, \beta_{2k+2})$ is an adjacent pair in R_m.

If k is odd then the proof is similar. But this time,

$$\alpha_{1,2k-1} = \alpha_{1,3} * \psi_{1,3}(\alpha_{5,7} * \cdots * \psi_{1,2k-1}(\alpha_{2k-1})$$

and the number of intersections with the unit circle is odd.

Now we consider the case of $\beta \in \mathcal{Z}_m(3/7, +, -, p)$, that is, with S^1-intersection at $e^{2\pi ix}$ for $x \in (-q_{p+1}, -q_p)$. For the base of the induction, we consider $\alpha_{q_p} * \psi_{q_p}(\beta)$. Then the first arc of the form $\rho(\overline{\eta_1}) * \rho(\eta_2)$ crossed by this, for an adjacent pair $(\eta_1, \eta_2) \in \Omega_m$, is for an adjacent pair $(\rho(\eta_1), \rho(\eta_2)) = (\beta_1, \beta_2)$ in $R_{m,p}$. Then let $\psi_1, \alpha_1)$ effect the matching between (β_1, β_2) and $(\beta'_1 \beta'_2)$ where (β'_1, β'_2) is the matched adjacent pair in $R_{m,p}$. Then $\overline{\beta_1} * \beta_2 = \psi_1(\overline{\beta'_1} * \beta'_2)$ is also the first arc crossed by $\alpha_1 * \psi_1(\beta)$, which means that $\overline{\beta'_1} * \beta'_2$ is the first arc crossed by β. Then we proceed inductively as before. □

CHAPTER 4

Quadruples of paths

4.1. Notation

Our goal in this work is to prove our main theorem, Theorem 1.9, concerning capture paths $\beta \in \mathcal{Z}_m(3/7, +, +, 0)$. This means analysing the sequence of translates of the fundamental domain crossed by β, as explained in Chapter 1. A particular structure for this sequence is claimed in theorem 2.8. In particular the sequence of pairs $(\omega_{2i-1}, \omega_{2i})$ is related to a sequence $(\beta_{2i-1}.\beta_{2i})$ of adjacent pairs $R_{m,0}$, and a matching sequence of adjacent pairs $(\beta'_{2i-1}, \beta'_{2i})$ in $R'_{m,0}$. Theorem 2.8 has not yet been completely proved. But the base case of the induction has been completed, and much of the inductive step has been established. This is enough to work with. We will now assume the existence of $(\beta_{2j-1}, \beta_{2j})$ for $j \leq i$, with the properties claimed in 2.8, and will work to extend these. In the process, the proof of Theorem 2.8 will be completed.

From now on we consider capture paths $\beta \in \mathcal{Z}_m(3/7, +, +, 0)$, and the associated sequences $\beta_j(\beta)$ for $j \leq 2i$ in $R_{m,0}$, and the homeomorphisms $\psi_{j,\beta}$ and $\psi_{j-2k,\beta}$ for any $k \geq 1$, as defined in 2.9. We write Y_m and Z_m for $Y_m(s)$ and $Z_m(s)$. If β ends in the point $p(x)$ for a word x (see 2.1) then we will also write $\beta_j(x)$ and $\psi_{j,x}$ and $\psi_{j-2k,j,x}$ for these quantities.

A natural way to proceed is to consider what happens if $\psi_{j-2k,j,\beta}$ is always the identity. If that were the case then the sequence of arcs $\overline{\beta_{2j-1}} * \beta_{2j} = \psi_{2i-1}(\overline{\beta'_{2j-1}} * \beta'_{2j})$ which is crossed by β can be computed just by studying $R_{m,0}$. Much of this chapter is devoted to studying this sequence, which is denoted by ζ_j. The precise definition is given below. We call such sequences *ζ-sequences.* One aim is to relate $\beta_j(\beta)$ to various ζ-sequences. In the process, we also obtain information about quadruples $(\beta_{4j+t}(\beta) : -3 \leq t \leq 0)$. In the process we are led to study *ζ-quadruples* $(\zeta_{4j+t} : -3 \leq t \leq 0)$.

4.2. Principal arcs and levels

Let (ζ_1, ζ_2) be any pair of paths starting at ∞ and with endpoints in Z_m, such that unit-disc-crossings are either along leaves of $L_{3/7}$ or in gaps of $L_{3/7}$. Thus, ζ_1 and ζ_2 could be paths in $R_{m,0}$ or $R'_{m,0}$, and be possibly adjacent in one of these, but not necessarily so. We define $k = k(\zeta_1, \zeta_2)$ to be the least integer such that the k'th unit-disc-crossings of ζ_1 and ζ_2 are distinct, up to isotopy preserving $Z_\infty \cup \{\infty\}$. We call $k(\zeta_1, \zeta_2)$ the *level* of (ζ_1, ζ_2).

If ζ_1 and ζ_2 are in $R_{m,0}$, this means that $w_i'(\zeta_1) = w_i'(\zeta_2)$ for $i < k$, but $w_k'(\zeta_1) \neq w_k'(\zeta_2)$. Note that, if $\zeta_j \in R_{m,0}$, the i'th unit disc crossing of ζ_j is the same as the $i-1$'th crossing of $\psi_{\zeta_j}(\zeta_j')$ for $i \geq 2$, unless it is the last half-crossing of both ζ_j and $\psi_{\zeta_j}(\zeta_j')$, in which case they meet in the middle.

Now we specialise to (ζ_1, ζ_2) being an adjacent pair in $R_{m,0}$. Then the first unit-disc-crossing of ζ_1 starts at a point $e^{2\pi it}$ for $t \in [\frac{2}{7}, \frac{9}{28}]$ and similarly for ζ_2. If k is odd, then the k'th crossing of ζ_1 starts to the right of that of ζ_2 and if k is even it starts to the left. If $k > 1$ and all letters of $w_{k-1}'(\zeta_1) = w_{k-1}'(\zeta_2)$ are in the set $\{L_3, L_2, R_3\}$, then the k'th crossings of ζ_1 and ζ_2 start on the upper unit circle if and only if k is odd. If the word $w_{k-1}'(\zeta_1)$ contains some other letter, then it must contain BC or UC, and the prefix ending in this first occurrence of BC or UC is a prefix of $w_j'(\zeta_1) = w_j'(\zeta_2)$ for all $i \leq j \leq k-1$ for some i. If the first occurrence is of BC is preceded by an odd number of letters in $\{L_3, L_2\}$, or the first occurrence is UC, preceded by an even number of letters in $\{L_3, L_2\}$, then i is odd, and all crossings from the i'th onwards have both endpoints on the upper unit circle. In the opposite cases, i is even and all crossings from the i'th onwards have both endpoints on the upper unit circle.

We define $D(\zeta_1, \zeta_2)$ to be the region between the k'th unit-disc intersections of ζ_1 and ζ_2, where $k = k(\zeta_1, \zeta_2)$, replacing the k'th unit-disc intersection by the leaf of $L_{3/7}$ containing the point on S^1 at the start of the intersection, if is is not a complete crossing.

The *principal arc of* (ζ_1, ζ_2) is the arc of $\overline{\zeta_1} * \zeta_2$ between the k'th unit-disc-crossings of ζ_1 and ζ_2.We denote it by $I(\zeta_1, \zeta_2)$. It is the unique arc bounded by endpoints of essential unit-disc crossings of ζ_1 and ζ_2 which are also essential unit-disc crossings on $\overline{\zeta_1} * \zeta_2$, up to homotopy. The principal arc can have both endpoints on either the upper or lower unit circle. The arc of the unit circle between the endpoints of $I(\zeta_1, \zeta_2)$ is in $D(\zeta_1, \zeta_2)$.

There are two types of principal arcs of adjacent pairs in $R_{m,0}$:

Type 1. $w_k'(\zeta_1) = u_1$ and $w_k'(\zeta_2) = u_2$ are adjacent values of w_k', subject to the fixed value of $w_{k-1}'(\zeta_1) = w_{k-1}'(\zeta_2)$ if $k > 1$, and $w_{k-1}'(\zeta_1)$ is not between $w_k'(\zeta_1)$ and $w_k'(\zeta_2)$.

Type 2. $k > 1$, and for the fixed value of $w_{k-1}'(\zeta_1) = w_{k-1}'(\zeta_2) = u$, with the k'th crossing of ζ_1 starting furthest left and that of ζ_2 starting furthest right if k is odd, and that of ζ_1 starting furthest right and that of ζ_2 starting furthest left if k is even.



FIGURE 1. $D(\zeta_1,\zeta_2)$ and the principal arc $I(\zeta_1,\zeta_2)$



FIGURE 2. A type 2 principal arc

For both types, the path ζ_1 is maximal among among paths in $R_{m,0}$ with $w'_k(\zeta_1)=u_1$, while ζ_2 is minimal among paths $\zeta\in R_{m,0}$ with $w'_k(\zeta_2)=u_2$. We shall refer to (ζ_1,ζ_2) being *of type 1* (or 2) if $I(\zeta_1,\zeta_2)$ is of type 1 (or 2). The figures show examples of a principal arcs $I(\zeta_1,\zeta_2)$. On the first figure, the set $D(\zeta_1,\zeta_2)$ below $I(\zeta_1,\zeta_2)$ is also marked. The principal arc in the first figure could be of type 1 or type 2, depending on whether $\zeta_1<\zeta_2$ or $\zeta_2<\zeta_1$. If $\zeta_2<\zeta_1$ then there could be at least one path ζ_3 with $k(\zeta_1,\zeta_2)=k(\zeta_1,\zeta_3)=k(\zeta_2,\zeta_3)$ and $w'_{k-1}(\zeta_3)=w'_{k-1}(\zeta_1)=w'_{k-1}(\zeta_2)$, and such that $\zeta_1<\zeta_3$, and the k'th unit-disc crossing of ζ_3 is strictly between the k'th unit-disc crossings of ζ_1 and ζ_2. The second figure must be a type two arc, because, for such a configuration, we must have $\zeta_1<\zeta_2$, and a path ζ_3 with $\zeta_2<\zeta_3$ and $w'_{k-1}(\zeta_1)=w'_{k-1}(\zeta_2)=w'_{k-1}(\zeta_3)$, the the k'th unit-disc crossing of ζ_3 between those of ζ_1 and ζ_2, must exist.

4.3. Principal arcs for $[\psi]$ and $R_{m,0}$

For the results to be proved, we need principal arcs of $\alpha*\psi(R_{m,0})$, for $[\psi]\in G$ and $[\alpha]\in\pi_1(\overline{\mathbb{C}}\setminus Z_m(s),\infty)$ with $s\simeq_\psi\sigma_\alpha\circ s$. In fact we define *principal arcs for* $[\psi]$ *and* $R_{m,0}$ *over* β, for $\beta\in\mathcal{Z}_m(3/7,+,+)$ as follows, for any adjacent pair (ζ_1,ζ_2) in $R_{m,0}$ such that $\psi(\overline{\zeta_1}*\zeta_2)$ crosses β at least once. We define exactly one principal arc for $[\psi]$ over β for each such pair. This is simply the first arc of $\psi(\overline{\zeta_1}*\zeta_2)$ which is crossed by β, up to isotopy preserving $S^1 \cup Z_m(s)$, where $\psi(\overline{\zeta_1} * \zeta_2)$ is chosen up to isotopy preserving $Z_m(s)$ to have only essential intersections with S^1. If $\psi = \psi_{1,4\lfloor i/2\rfloor -1,\beta}$, and the existence of the sequence $\beta_j(\beta)$ as in Theorem 2.8 has been established for $j \leq 2i$, then the principal arc for $(\beta_{2i-1}(\beta), \beta_{2i}(\beta), [\psi])$ is indeed the arc of $\psi(\overline{\beta_{2i-1}(\beta)} * \beta_{2i}(\beta))$ such that the lift determined by $\alpha_{1,4\lfloor i/2\rfloor -1} * \psi_{1,4\lfloor i/2\rfloor -1,\beta}(\beta_{2i-1}(\beta))$ is crossed by the lift of β. This is the first arc of $\psi(\overline{\zeta_1} * \zeta_2)$, for any adjacent pair (ζ_1, ζ_2) in $R_{m,0}$, which is crossed by β after crossing the principal arc for $(\beta_{2i-3}(\beta), \beta_{2i-2}(\beta), [\psi_{1,4\lfloor (i-1)/2\rfloor -1}])$. This is useful to know if $i \geq 3$, because, as we shall see in 4.10.4, for i even and ≥ 4, $\psi_{2i-3,2i-1,\beta}$ is the identity on the principal arc for $[\psi_{1,2i-1,\beta}]$ over β which is immediately beneath the principal arc for $(\beta_{2i-3}(\beta), \beta_{2i-2}(\beta), [\psi_{1,2i-5,\beta}])$ over β.

4.4. The ζ-sequence $\zeta_i(x)$

Fix any infinite admissible word x ending in $(L_2R_3L_3)^\infty$ and such that $D(x)$ intersects $\{e^{2\pi it} : t \in (\frac{2}{7}, \frac{9}{28})$. We are thinking of $p(x)$ as being the last S^1 crossing-point of a path in $\mathcal{Z}(3/7, +, +, 0)$. We define the *$\zeta$-sequence for* x, $\zeta_j(x)$, for $\leq 2n(x)$ by the following properties:

1. $(\zeta_{2i-1}(x), \zeta_{2i}(x))$ is an adjacent pair in $R_{m,0}$ with $\zeta_{2i-1}(x) < \zeta_{2i}(x)$;
2. $D(x) \subset D(\zeta_{2i-1}(x), \zeta_{2i}(x))$ and $I(\zeta_{2i-1}(x), I(\zeta_{2i}(x))$ is over $D(x)$;
3. $I(\zeta_1(x), \zeta_2(x))$ is the outermost arc as in 2, and for $i \geq 1$, the principal arc $I(\zeta_{2i+1}, \zeta_{2i+1}(x))$, if it exists, is not separated from $I(\zeta_{2i-1}(x), \zeta_{2i}(x))$ by any other principal arc $I(\omega_1, \omega_2)$.

There is one more property, which we now explain.The pair $(\zeta_1(x), \zeta_2(x))$ is always defined, but it can happen that $(\zeta_{2i+1}(x), \zeta_{2i+2}(x))$ is not defined for any $i \geq 1$, satisfying conditions 1 to 3. This happens whenever x is not the endpoint of any path in $R_{m,0}$. In the language of [**22**], this happens precisely when $D(x)$ is not contained in U^y for any value y of $w_1'(.)$. The x for which $\zeta_i(x)$ is defined only for $i = 1$, 2, and for which $x \neq w(\zeta_1(x))$, is precisely the set of x such that $D(x) \subset D(vbv_2)$, for v_2 such that $y_1 < w_1'(v_2) \leq y_2$, and y_1 and y_2 as in 4.7.2. Also, more simply, $(\zeta_{2i+1}(x), \zeta_{2i+2}(x))$ is not defined if $x = w(\zeta_{2i-1}(x))$. In that case, we define $i = n(x)$. If $(\zeta_3(x), \zeta_4(x))$ is defined, then $x = w(\zeta_{2n(x)-1}(x))$. If there is more than one adjacent pair (η_1, η_2) in $R_{m,0}$ such that $I(\eta_1, \eta_2)$ is bounded by $I(\zeta_{2i-1}(x), \zeta_{2i}(x))$ and is not separated from $I(\zeta_{2i-1}(x), \zeta_{2i}(x))$ by any other principal arc, and $D(x)$ is bounded by $I(\eta_1, \eta_2)$, then there are exactly two such. If $i > 1$, then it is always true that $w_1'(\zeta_{2i-1}(x)) = w_1'(\zeta_{2i}(x)$ and there is exactly one such pair (η_1, η_2) with $w_1'(\eta_1) = w_1'(\eta_2) = w_1'(\zeta_{2i-1}(x))$. and in this case we choose (η_1, η_2) to be this $(\zeta_{2i+1}, \zeta_{2i+2}(x))$. If $i = 1$ then the two adjacent pairs (η_1,η_2) and (η_3,η_4) in $R_{m,0}$ are such that $I(\eta_1,\eta_2)=I(\eta_3,\eta_4)$ and $\eta_1<\zeta_1(x)<\zeta_2(x)<\eta_3(x)$. Condition 4 is then as follows:

4. Write $w_2'(x)=w_1(x)u_2$. Then $(\zeta_3(x),\zeta_4(x))$ is chosen so that $\zeta_3(x)<\zeta_1(x)$ if $L_3u_2>w_1'(\zeta_1(x))$ and $\zeta_3(x)>\zeta_1(x)$ if $L_3u_2\leq w_1'(x)$. If $i>1$ then $w_1'(\zeta_{2i+1}(x))=w_1'(\zeta_{2i+2}(x))$.

The reason for the rather complicated definition of (ζ_3,ζ_4) is that $w_2'(.)$ is not injective on all words, as shown by the definitions in section 3. But there are never more than two adjacent pairs (η_1,η_2) and (η_3,η_4) such that $I(\eta_1,\eta_2)$ and $I(\eta_3,\eta_4)$ both bound $D(x)$ and are bounded by $I(\zeta_{2i-1}(x),\zeta_{2i}(x))$, and are not separated from $I(\zeta_{2i-1}(x),\zeta_{2i}(x))$ by any other principal arc. In this case there are $y_1\neq y_2$ such that

$$w_1'(\eta_1)=w_1'(\eta_2)=y_1,\quad w_1'(\eta_3)=w_1'(\eta_4)=y_2,$$

and

$$I(\eta_1,\eta_2)=I(\eta_3,\eta_4)\subset(U^{y_1}\cap U^{y_2}).$$

We can also define a *lower ζ sequence* for x such that $D(x)$ intersects $\{e^{2\pi it}:t\in(\frac{19}{28},\frac{5}{7})\}$. The definition is similar but the first pair (ζ_1,ζ_2) in a lower ζ sequence satisfies $k(\zeta_1,\zeta_2)=2$.

We can also define the sequence $\zeta_i(x)$ for $x\in D(BC)$. and $x=p(\gamma)$ with $\gamma\in\mathcal{Z}(3/7,+,-)$. For the moment , we just do it for $x=p(\beta)$ with $\beta\in\mathcal{Z}(3/7,+,-)$ and for $x\in D(BC)\setminus D(BCL_1R_2UC)$. The outermost principal arc $I(\zeta_3(x),\zeta_4(x))$ is the same for all such x, and is the unique principal arc $I(\zeta_3,\zeta_4)$ such that $w_1'(\zeta_3)=w_1'(\zeta_4)$ and the third intersections of ζ_3 and ζ_4 with the unit circle are the furthest right and left respectively. We then define the sequence $\zeta_{2i-1}(x),\zeta_{2i}(x)$ for $i\geq 3$ as before satisfying conditions 1 to 3 above. The complications of condition 4 do not arise, as there is always exactly one choice for $(\zeta_{2i+1}(x),\zeta_{2i+2}(x))$ so that 1 to 3 are satisfied, once $(\zeta_{2i-1}(x),\zeta_{2i}(x))$ is given.

Finally for $x\in D(BCL_1R_2)^kUC\backslash D((BCL_1R_2)^{k+1}UC)$, any pair (η_1,η_2) such that $I(\eta_1,\eta_2)$ bounds $D(x)$ is such that η_1 and η_2 have at least $k+1$ common unit-disc crossings. We therefore only define $(\zeta_{2i-1}(x),\zeta_{2i}(x))$ for $i\geq k$.

LEMMA 4.4.1. 1. $k(\zeta_1,\zeta_2)=1$

2. *The principal arc $I(\zeta_{2i-1},\zeta_{2i})$ is of Type 1 as in 4.2 if i is odd, and of type 2 if i is even.*
3. *$k(\zeta_{2i-1},\zeta_{2i})$ is always odd, $1<k(\zeta_{2i-1},\zeta_{2i})$ if $i>1$, and for all $i<n(x)$,*

$$k(\zeta_{2i-1},\zeta_{2i})\leq k(\zeta_{2i+1},\zeta_{2i+2})\leq k(\zeta_{2i-1},\zeta_{2i})+2.$$

4. *If i is even, and $i<n(x)$, then $k(\zeta_{2i+1},\zeta_{2i+2})=k(\zeta_{2i-1},\zeta_{2i})$.*

Proof. Clearly $I(\zeta_1,\zeta_2)$ has to be of type 1, and $k(\zeta_1,\zeta_2)=1$. Also, it is clear that if $i>1$ then $k(\zeta_{2i-1},\zeta_{2i})>1$.

Now let $i>1$ and $k=k(\zeta_{2i-1},\zeta_{2i})$ and assume inductively that k is odd. If the longest prefix of $w'_k(\zeta_{2i-1})$ which ends in BC or UC does not coincide with the longest such prefix of $w'_k(\zeta_{2i})$, then there is no principal arc strictly inside $I(\zeta_{2i-1},\zeta_{2i}(x))$ and i is maximal. Now we assume that i is not maximal, because otherwise there is nothing to prove.

Now suppose that $\zeta\in R_{m,0}$ and j is minimal with $w'_j(\zeta)$ strictly between $w'_k(\zeta_{2i-1})$ and $w'_k(\zeta_{2i})$, and such that the j'th unit-disc-crossing of ζ intersects the unit-circle arc bounded by $I(\zeta_{2i-1},\zeta_{2i})$. Then the j'th unit-disc-crossing of ζ must intersect both the circle arcs bounded by the k'th unit-disc crossings of ζ_{2i-1} and ζ_{2i}. Write $u=w'_j(\zeta)L_2$. We claim that $j\geq k-1$. Since ζ intersects the circle arcs bounded by the k'th unit-disc crossings of ζ_{2i-1} and ζ_{2i}, and letter BC or UC in $w'_j(\zeta)$ must be in the longest common prefix of $w'_k(\zeta_{2i-1})$ and $w'_k(\zeta_{2i})$. So $w'_j(\zeta)$ must end in L_3 or R_3. Then $u=w'_j(\zeta)L_2$ is an admissible word. Now either $w'_{k-1}(\zeta_{2i-1})=w'_{k-1}(u,w'_1(x))$, or $w_{k-1}(\zeta_{2i-1})$ is a proper prefix of $w_{k-1}(u,w'_1(x))$ and $w'_{k-1}(\zeta_{2i-1})$ is a proper prefix of $w'_{k-1}(u,w'_1(x))$. The latter only happens if, for $n\geq 3$ and odd,

$$w_{k-1}(u)=u_0u_1u_2\cdots u_n,\quad w_{k-1}(\zeta_{2i-1})=w_{k-1}(\zeta_{2i})=u_0u_1,$$

$$w'_{k-1}(u)=u_0\cdots u_{n-1},\quad w'_{k-1}(\zeta_{2i-1})=w'_{k-1}(\zeta_{2i})=u_0,$$

where $u_\ell=(L_3L_2R_3)^{r_\ell}u'_\ell$ for some $r_\ell\geq 0$ and

$$u'_\ell\in\{L_3L_2R_3L_3,\ L_3(L_2R_3)^2,L_3^3\}$$

for $1\leq\ell\leq n-1$. Now $u_0u_1u_2$ cannot be a prefix of $w'_k(\zeta_{2i-1})$ or $w'_k(\zeta_{2i})$ — which both extend $w_{k-1}(\zeta_{2i-1})=u_0u_1$ — because otherwise $w'_{k-1}(\zeta_{2i-1})$ would be defined differently. It follows that in this case $D(u_0u_1u_2)$ is strictly between $w'_k(\zeta_{2i-1})$ and $w'_k(\zeta_{2i})$. Also $w_{k-2}(u)$ is a prefix of u_0. Hence $j\geq k-1$.

Also, j must have the opposite parity to k, because the j'th crossing of ζ must start on the circle arc bounded by $w'_k(\zeta_{2i-1})$ and $w'_k(\zeta_{2i})$ which is not bounded by $I(\zeta_{2i-1},\zeta_{2i})$. But then $w'_{k-1}(\zeta)$ is equal to, or extends, $w'_{k-1}(\zeta_{2i-1})$. We have seen that if $w'_{k-1}(\zeta_{2i-1})$ is a proper prefix of $w'_{k-1}(\zeta)$, then $D(w'_{k-1}(\zeta_{2i-1})u_1L_3L_2)$ is strictly between $w'_k(\zeta_{2i-1})$ and $w'_k(\zeta_{2i})$ and hence $w'_{k-1}(\zeta)$ is strictly between $w'_k(\zeta_{2i-1})$ and $w'_k(\zeta_{2i})$ and hence $j=k-1$. If $w'_{k-1}(\zeta_{2i-1})=w'_{k-1}(\zeta)$ then we claim that $j=k+1$. For suppose not so. Then $D(w_{k+1}(\zeta))$ intersects the region between $w'_k(\zeta_{2i-1})$ and $w'_k(\zeta_{2i})$ and $D(w'_{k+1}(\zeta))$ intersects the complement. Since $w_k(\zeta)$ is a prefix of both $w_{k+1}(\zeta))$ and $D(w'_{k+1}(\zeta)$, it must be the case that $w_k(\zeta)$ is a prefix of one of

$w_k'(\zeta_{2i-1})$ and $w_k'(\zeta_{2i})$. But then $w_k(\zeta)=w_k(\zeta_{2i-1})$ is a prefix of $w_k'(\zeta_{2i-1})$, or similarly for ζ_{2i}. Both of these are impossible.

It follows that the $k-1$'th or the $k+1$'th unit disc crossing of ζ_{2i+1} and ζ_{2i+2} is strictly between the k'th unit-disc crossings of ζ_{2i-1} and ζ_{2i}. If the $k-1$'th crossing is between and $I(\zeta_{2i-1},\zeta_{2i})$ is of type 1, then $w_{k-1}'(\zeta_{2i+1})\neq w_{k-1}'(\zeta_{2i-1})$ and $I(\zeta_{2i-1},\zeta_{2i+1})$ must be of type 2. If the $k+1$'th unit disc crossing of ζ_{2i+1} and ζ_{2i+2} is the first to be strictly between, then again $I(\zeta_{2i+1},\zeta_{2i+2})$ must be of type 2. Now suppose that $I(\zeta_{2i-1},\zeta_{2i})$ is of type 2. Then if $i<n(x)$, there is at least one other path ζ between ζ_{2i-1} and ζ_{2i} with $w_{k-1}'(\zeta)=w_{k-1}'(\zeta_{2i-1})$. It follows that $w_{k-1}'(\zeta_{2i+1})=w_{k-1}'(\zeta_{2i+2})$ and $k(\zeta_{2i+1},\zeta_{2i+1})=k(\zeta_{2i-1},\zeta_{2i})$ and $I(\zeta_{2i+1},\zeta_{2i+2})$ is of type 1. This completes the proof of 2, 3 and 4. □

4.5. Quadruples

We have the following immediate corollary to Lemma 4.4.1

Corollary 4.5.1. *For any quadruple* $(\zeta_{4i+s}(x):-3\leq s\leq 0)$*, for any* $i\geq 1$*,* $(\zeta_{4i-3}(x),\zeta_{4i-2}(x))$ *is of type 1 and* $(\zeta_{4i-1}(x),\zeta_{4i}(x))$ *is of type 2. In addition, there are two possibilities for the quadruple* $(\zeta_{4i+s}:-3\leq s\leq 0)$ *for* $i\geq 1$*.*

A. $k(\zeta_{4i-1}(x),\zeta_{4i}(x))=k(\zeta_{4i-3}(x),\zeta_{4i-2}(x))+2$;
B. $k(\zeta_{4i-1}(x),\zeta_{4i}(x))=k(\zeta_{4i-3}(x),\zeta_{4i-2}(x))$.

We shall refer to these as *quadruples* of *type A* and *type B*. Any quadruple $(\zeta_{4i+t}(x):-3\leq t\leq 0)$ in a sequence $\zeta_j(x)$ will be referred to as a ζ*-quadruple*. If $\zeta_j(x)$ is an upper ζ sequence then we will refer to $(\zeta_{4i+s}(x):s\leq 0$ as an *upper* ζ *quadruple*. Lower ζ quadruples are similarly defined. We recall the notation $E_{\zeta,\eta}$ of 3.8 to denote an arbitrarily small disc neighbourhood of the closed loop $\alpha_\zeta*\overline{\alpha_\eta}$, for paths ζ and $\eta\in R_{m,0}$. We continue to use this notation for quadruples of various types. So if $(\zeta_i:1\leq i\leq 4)$ is a quadruple, we use the notation E_{ζ_1,ζ_3} to denote an arbitrarily small neighbourhood of $\alpha_{\zeta_1}*\overline{\alpha_{\zeta_3}}$. The figures show examples of type A and type B quadruples, together with the shaded region E_{ζ_1,ζ_3} in each case. In the examples drawn we have $\zeta_3<\zeta_4<\zeta_1<\zeta_2$ for the type A quadruple, and $\zeta_1<\zeta_2<\zeta_3<\zeta_4$ for the type B quadruple. Examples like this will arise later, but these are not the only possibilities.

4.6. Quadruples of types C and AC

We shall see later that the sets of the loops $(\beta_{4i+j}(\beta):1\leq j\leq 4)$ are always closely related to quadruples or lower quadruples, of types A or B or *type C* or *type AC*, which we now describe. We say that $(\eta_i:1\leq i\leq 4)$



FIGURE 3. Example of a type A quadruple



FIGURE 4. Example of a type B quadruple

is of *type C* if (η_1, η_2) and (η_3, η_4) are two adjacent pairs with $\eta_1 < \eta_2$ and $\eta_3 < \eta_4$ and one of the following holds.

C-1 There is a basic exchangeable pair (a, b) which is one of the pairs of 3.9.1.4, such that the following hold.

$$k(\eta_1, \eta_2) = 1, \quad k(\eta_3, \eta_4) = 3,$$

$$w_1(\eta_3) = v_1 a L_3, \quad w_1'(\eta_3) = x = w_1'(v_1 a L_3 L_2), \quad w_2'(\eta_3, x) = v_1 a v',$$

$$w_1'(v_1 a L_3 L_2)(L_2 R_3 L_3)^\infty < v'(L_2 R_3 L_3)^\infty < w_1'(v_1 b L_3 L_2)(L_2 R_3 L_3)^\infty,$$

$$w_1'(\eta_1)(L_2 R_3 L_3)^\infty < v_1 b L_3 L_2$$

and η_1 is maximal with respect to this property, and (η_1, η_2) of type 1 and (η_3, η_4) of type 2. Moreover,

$$I(\eta_3, \eta_4) \subset u E_{\eta_1, \eta_3},$$

where $u \leftrightarrow u'$ is an exchange for η_1 which is prefixed by $v_1a \leftrightarrow v_1b$ such that $w_2(uL_3L_2) = uL_3L_2$ and $w_2(u'L_3L_2) = u'L_3L_2$. Such a quadruple is said to be *type C of level 1*

C-2. There is a quadruple $(\zeta_i : 1 \leq i \leq 4)$ which is of type C of level 1 such that

$$w_1'(\eta_i) = w_1'(\zeta_3) = w_1'(\zeta_4),$$

(η_1, η_2) is of type 1 and (η_3, η_4) is of type 2 and for some odd $k_1 \geq 3$,

$$k(\eta_1, \eta_2) = k_1, \quad k(\eta_3, \eta_4) = k_1 + 2,$$

and, writing $v = v_1aL_3L_2v_2$ for a word v_2,

$$I(\eta_j, \eta_{j+1}) = vI(\zeta_j, \zeta_{j+1}), \quad E_{\eta_1,\eta_3} = vE_{\zeta_1,\zeta_3}.$$

The definition of type AC is exactly similar to type C except that the basic exchange $va \leftrightarrow vb$ is one of the types listed in 3.9.1.4. This makes for some important differences, the chief of which are that $I(\eta_3, \eta_4)$ spans an arc on the lower unit circle, and that now $w_1'(vaL_3L_2) = w_1'(vbL_3L_2)$.

4.7. Intersections between sets E_{η_1,η_3} and $s^{-n}E_{\omega_1,\omega_3}$

If E any set with $E \subset D(L_3)$, then a component of $s^{-n}(E)$ is of the form uE for some word u such that uL_3 is admissible if $E \subset D(L_3)$. This means simply that $uE = \{p(ux) : p(x) \in E\}$. The definition and properties of $w_j(.)$ and $w_j'(.)$ and $w_j'(.,y)$, for any value y of w_1', ensure that if $(\eta_i : 1 \leq i \leq 4)$ is an upper or lower quadruple and $w_1'(uL_3L_2) = w_1'(\eta_3) = y$, then there is an upper or lower ζ quadruple $(\eta_i : 5 \leq i \leq 8)$ such that $w_1'(\eta_i) = y$ for all $5 \leq i \leq 8$ and

$$uE_{\eta_1,\eta_3} = E_{\eta_5,\eta_7} \subset U^y$$

The condition 4 in the definition of the zeta sequence ensures that $w_{k_1}'(.,y)$ is constant on

$$\{\eta \in R_{m,0} : (\eta, \eta^2) \text{ is an adjacent pair in } R_{m,0}, \quad I(\eta, \eta^2) \subset E_{\eta_5,\eta_7}\},$$

where $k_1 = k(u) + k(\eta_1, \eta_2)$ where $k(u)$ is the largest integer k such that $w_k(uL_3L_2)$ is a proper prefix of uL_3L_2. (This condition is only needed if $k(\eta_1, \eta_2) = 1$.) A necessary and sufficient condition for $(\eta_i : 5 \leq i \leq 8)$ to be an upper ζ quadruple with $k(\eta_5, \eta_6) \geq 3$ is that $k(u)$ be even and $k(u) + k(\eta_1, \eta_2) \geq 3$.

Lemma 4.7.1. *Let $(\eta_i : 1 \leq i \leq 4)$ and $(\omega_j : 1 \leq j \leq 4)$ be two upper or lower ζ quadruples, such that*

$$w_1'(\eta_3) = w_1'(\omega_3),$$

and $(\eta_1,\eta_2)=(\omega_1,\omega_2)$ *if* $k(\eta_1,\eta_2)=k(\omega_1,\omega_2)=1$. *Write* $E=E_{\eta_1,\eta_3}$. *Let* E' *be a component of* $s^{-n}(E_{\omega_1,\omega_3})$ *for some* $n>0$. *Then* $E\cap E'\neq\emptyset$ *if and only if one of* E' *and* E *is contained in the other.*

Remark

1. For this lemma to be useful, we still have to show that for $\alpha=\alpha_{\eta_1,\eta_3}$, a component of $(\sigma_\alpha\circ s)^{-n}(E_{\eta_1,\eta_3})$ is, up to homotopy preserving Z_∞, the union of two components of $s^{-n}(E_{\eta_1,\eta_3})$ and a path joining the two. This will be shown in later lemmas.
2. The conditions of a ζ quadruple ensure that the second unit-disc-crossing of η_3 is not contained in $\cup_{n\geq 0}s^{-n}(E_{\eta_1,\eta_3})$.

PROOF. Let u be the word such that $E'=uE_{\omega_1,\omega_3}$. Let $(\omega_i:5\leq i\leq 8)$ be the quadruple such that $uE_{\omega_1,\omega_3}=E_{\omega_5,\omega_7}$. If $k(u)\geq 1$, then $w_1'(\omega_i)=w_1'(u)$ for all $5\leq i\leq 8$ and $k(\omega_7,\omega_8)\geq k(\omega_5,\omega_6)>1$.

If $k(u)=0$, then $w_1'(\omega_i)=uw_1'(\omega_{i-4})$ for $5\leq i\leq 8$. If $k(\omega_1,\omega_2)>1$ then $w_1'(\omega_i)=w_1'(\omega_1)$ for all $1\leq i\leq 4$ and

$$w_1'(\omega_i)=w_1'(\omega_5)=uw_1'(\omega_3)=uw_1'(\eta_3)$$

for $5\leq i\leq 7$. It follows that E' is contained in or disjoint from E in this case, except possibly when $k(\eta_1,\eta_2)=1$. But then the only case to consider is when $w_1'(\omega_i)=w_1'(\eta_1)$ for all $5\leq i\leq 8$ or $w_1'(\omega_i)=w_1'(\eta_2)$ for all $5\leq i\leq 8$. But then η_1 (or η_2) is either the maximal (or minimal) path in $R_{m,0}$ with this value of w_1', and depending on whether $\eta_4\leq\eta_1$ or $\eta_2\leq\eta_3$, as well as whether $w_1'(\omega_i)=w_1'(\eta_1)$ or $w_1'(eta_2)$, E' is contained in or disjoint from E.

So now suppose that $k(u)>0$. Write

$$k_1=k(\eta_1,\eta_2),\quad k_2=k(\eta_3,\eta_4),\quad k_3=k(\omega_5,\omega_6),\quad k_4=k(\omega_7,\omega_8).$$

If $I(\omega_5,\omega_6)=I(\eta_1,\eta_2)$, then we have disjointness of E and E' if ω_7 and η_3 are on opposite sides of η_1, and one of E and E' is contained in the other if they are on the same side. So now suppose that $I(\omega_5,\omega_6)\neq I(\eta_1,\eta_2)$. Now we consider the boundary of E, up to Z_∞-preserving homotopy. Either the k_1-1'th or the k_1-2'th unit-disc-crossings of the η_i are in common (up to Z_∞-preserving homotopy) depending on whether the quadruple is of type A or B. Lock these common crossings together. Lock η_1 and η_2 together up to the last common unit-disc crossing, and similarly for η_3 and η_4. Finally, lock together $I(\eta_1,\eta_2)$ and $I(\eta_3,\eta_4)$. In the figures shown earlier of the sets E_{ζ_1,ζ_3}, for both type A and type B quadruples, the horizontal arcs of the boundary of E_{ζ_1,ζ_3} denote the locking of the principal arcs $I(\zeta_1,\zeta_2)$ and $I(\zeta_3,\zeta_4)$. Returning to the quadruple $(\eta_i:1\leq i\leq 4)$, we obtain a set bounded by the locked-together principal arcs and by common segments of the paths η_i. This set is E up to homotopy, and so we denote it by E. It has directed boundary with direction given by the paths η_i. No path in $\zeta \in R_{m,0}$ with $w_1'(\zeta) = w_1'(\eta_3)$ travels through any of the "locks" except for the lock between the common crossings of all the η_i, as (η_1, η_2) is of type 1 and (η_3, η_4) is of type 2. So the only way for a path $\zeta \in R_{m,0}$ to enter E is between η_4 and η_1, or between η_2 and η_3 (depending on whether $\eta_3 < \eta_1$ or $\eta_1 < \eta_3$). We choose locks similarly for E'. Then either one of E and E' is contained in the other or they are disjoint. For suppose none of these hold. Then the boundaries of E and E' must intersect transversally, and since both E and E' are topological discs, there must be two transversal intersections. But the only place for a transversal intersection is between the common lock of all the η_i and the common lock of all the ω_i. So this is impossible, and one of E and E' must be contained in the other and they are disjoint.

We have to use a slight variant on this if $k_1 = 1$, because in that case there is no common crossing of all the η_i. But there is still only one way in: through the top.

$\square$

COROLLARY 4.7.2. *Let $(\eta_i : 1 \leq i \leq 4)$ be an upper or lower ζ quadruple, let $E = E_{\eta_1,\eta_3}$ and let E' be a component of $s^{-n}E$ for some $n > 0$. Then $E' \cap I(\eta_j, \eta_{j+1}) = \emptyset$ for $j = 1$, 3 and either $E' \subset E$ or $E' \cap E = \emptyset$.*

PROOF. We have $E' = E_{\eta_5,\eta_7}$ for a quadruple $(\eta_{4+j} : 1 \leq j \leq 4)$ and with $k(\eta_5, \eta_6) \geq k(\eta_1, \eta_2)$. If E' contains $I(\eta_1, \eta_2)$, then $k(\eta_5, \eta_6) = k(\eta_1, \eta_2)$. We clearly cannot have $E \subset E'$, so we must have $E' \subset E$ and $E' \neq E$. Since $I(\eta_1, \eta_2)$ is the furthest left (or right) of the principal arcs in E of level k_1, then if $I(\eta_1, \eta_2)$ is in E' it must be equal to $I(\eta_5, \eta_6)$ which implies that $I(\eta_1, \eta_2)$ is fixed by s^i. This is impossible. So $I(\eta_1, \eta_2)$ is not contained in E'. Now suppose that $I(\eta_3, \eta_4) \subset E'$. Since $I(\eta_3, \eta_4)$ intersects ∂E, and $E' \subset E$, this is only possible if $I(\eta_3, \eta_4) \subset \partial E'$. But this, in turn is only possible if $I(\eta_3, \eta_4) \subset s^{-i}(\partial E)$. But ∂E is a union of principal arcs of level k_1, with $I(\eta_1, \eta_2)$ at one end, and the k_1'th unit-disc crossing of η_3 at the other end, and $I(\eta_3, \eta_4)$. We cannot have $I(\eta_3, \eta_4) \subset s^{-i}(I(\eta_3, \eta_4))$, and anything else in ∂E mapping to $I(\eta_3, \eta_4)$ under s^{-i} leads to E' not contained in E. So $I(\eta_3, \eta_4)$ is not contained in E' either. $\square$

We need a variant on 4.7.1 for quadruples of type C or AC.

LEMMA 4.7.3. *Let $(\eta_i : 1 \leq i \leq 4)$ be of type C or AC. Write $k_1 = k(\eta_1, \eta_2)$ and $E = E_{\eta_1,\eta_3}$.*

1. *If $k_1 > 1$ then $I(\eta_1, \eta_2)$ and $I(\eta_3, \eta_4)$ are not contained in $s^{-n}(E$ for any $n > 0$.*

2. . *Let* u_1 *and* u_2 *be admissible words. Then the only way that* u_1E *and* u_2E *can intersect without one being contained with the other is if* $u_2 = u_1va$ *or* $u_1 = u_2va$.
3. *If* $k_1 = 1$, *then* $I(\eta_1, \eta_2)$ *is disjoint from* $s^{-n}(E)$ *and* $I(\eta_3, \eta_4)$ *is contained in a component of* $s^{-n}(E)$ *for a single value of* $n > 0$.

Proof. 1. Suppose that $I(\eta_3, \eta_4) \subset uE$ for some nontrivial u. Since $k(\eta_1, \eta_2) > 1$ by the definition of type C or AC we have $w_1'(\eta_1) = w_1'(\eta_3) = x$, say.There is at most one $y$with $U^x \cap U^y \neq \emptyset$ and they have a common prefix apart form the last at most four letters. If $I(\eta_1, \eta_2)$ or $I(\eta_3 . \eta_4)$ is in uE then we need $ux = x$ or $uy = x$, both of which are impossible.
2. We can assume without loss of generality that u_2 is trivial. The question is then how it is possible to have $u_1E \cap E \neq \emptyset$. We have seen in 4.7.2 that it is impossible to have the first segment of η_1 in the unit disc in $u_1E_{\eta_1,\eta_3}$. It is similarly impossible to have the first segment of η_3 in the unit disc in u_1E because then $w_1'(\eta_3)$ would have to be between $u_1w_1'(\eta_3)$ and $u_1'(w_1'(\eta_1)$. The second unit-disc-crossing of η_3 is in vaE and . So va must be a prefix of u_1, and since $I(\eta_3, \eta_4) \subset \partial vaE$, we must have $va = u_1$.
3. We now have $k(\eta_1, \eta_2) = 1$. From the definition of type C, or AC, there is a basic exchange $a \leftrightarrow b$ and an admissible word v_1 such that $I(\eta_3, \eta_4) \subset v_1aE$. WE have $v_1aE \setminus E \neq \emptyset$, and by 2, v_1a is the unique nontrivial word with these properties.

□

4.8. A basic result about the quadruple homeomorphisms

One of our aims is to obtain a good bound on the supports of the homeomorphisms $\psi_{4i-3,4i-1,\beta}$, for any capture path $\beta \in \mathcal{Z}_\infty(3/7, +, +, 0)$. Any bound obviously has to be obtained inductively, because the support of $\psi_{4i-3,4i-1,\beta}$ is obtained from $E_{\beta_{4i-3},\beta_{4i-1},\beta}$, which is, in turn, obtained from $(\beta_{4i+t} : -3 \leq t \leq 0)$, which in turn depends on $\psi_{4j-3,4j-1,\beta}$ for $j < i$. It is clearly important to know how the supports of $\psi_{4j-3,4j-1,\beta}$, for varying j, intersect each other, and also how they intersect the boundaries of sets $E'_{\beta_{4i-3},\beta_{4i-1},x}$ for varying i.

We start with a preliminary lemma.

Lemma 4.8.1. *Let* (ζ_1, ζ_2) *be an adjacent pair in* $R_{m,0}$ *and let* $\zeta_{1,0}$ *be the initial segment of* ζ_1. *No component of* $R_{m,0} \cap \{z : |z| \geq 1\}$ *or of* $(\sigma_{\zeta_{1,0}} \circ s)^{-n}(s^{-1}(\zeta_{1,0}))$ *comes between* $I(\zeta_1, \zeta_2)$ *and* $I(\zeta_3, \zeta_4)$ *for any quadruple* $(\zeta_i : 1 \leq i \leq 4)$ *of type A or B, unless it is a component of* $\sigma_{\zeta_{1,0}} \circ s)^{-n}(s^{-1}(\zeta_{1,0}))$ *which is isotopic to* $I(\zeta_3, \zeta_4)$ *up to isotopy preserving* $Z_\infty \cup S^1$.

Proof. Any component of $(\sigma_{\zeta_{1,0}} \circ s)^{-n}(s^{-1}(\zeta_{1,0}))$ is isotopic, up to isotopy preserving $Z_\infty \cup S^1$, to $I(\zeta_5, \zeta_6)$, for some adjacent pair (ζ_5, ζ_6) in $R_{m,0}$.

By definition, there are no principal arcs separating $I(\zeta_1, \zeta_2)$ and $I(\zeta_3, \zeta_4)$. The result follows. □

We have the following.

THEOREM 4.8.2. *Let* $(\zeta^i : 1 \leq i \leq 4)$ *be an upper or lower quadruple or type C or AC. Write*

$$\zeta^1 = \zeta, \quad \psi = \psi_{\zeta,\zeta^3}, \quad E = E_{\zeta,\zeta^3}, \quad k_1 = k(\zeta, \zeta^2).$$

Write ζ_n *for the portion of* ζ *up to the n'th intersection with* S^1*.*

Then each component C *of* $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ *is a component of* $(\sigma_{\zeta_{2k_1-1}} \circ s)^{-i}(E)$*, and takes the form*

$$E_1 \cup E_2 \cup \omega_{1,m(C),n(C,1)} \cup \omega_{2,m(C),n(C,2)} \cup \ell,$$

where:

- E_1 *and* E_2 *are components of* $s^{-i}(E)$*;*
- C *is either contained in* U^0*, or disjoint from it, or* $\ell(C)$ *is an arc with endpoints on the unit circle on either side of* $e^{2\pi i(2/7)}$*, and for one of* $r = 1$ *or* 2*,* $E_r \cup \omega_{r,m(C),n(C,r)}$ *is contained in* U^0 *apart from the first segment on* $\omega_{r,m(C),n(C,r)}$*;*
- *If* $E_r \cup \omega_{r,n(C),m(C)}$ *intersects* U^0*, then* $\omega_{r,n(C),m(C)}$ *is the segment of* $\omega_r \in R_{m,0}$ *from the* $m(C)$*'th intersection with the unit circle to the* $n(C,r)$*'th for* $r = 1$ *and* 2*, and* $m(C) \leq 2k_1$*;*
- ℓ *is a component of* $s^{-j}(\zeta_0)$ *for some* $j \leq i$*;*
- *the only intersections between* C *and* α_ζ *are between* ℓ *and* ζ_{2k_1-1}*.*

In particular, the support of ψ *is contained in* $\cup_{i \geq 1}(\sigma_{\zeta_{2k_1-1}} \circ s)^{-i}(E)$*.*

PROOF. We start by showing that any intersections between $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ and α_ζ must be intersections between $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ and ζ, and in fact must be intersections with ζ_{2k_1}. Recall that α_ζ is an arbitrarily small perturbation of $\zeta * \psi_\zeta(\zeta')$. We need to cut down the intersections between $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ and α_ζ. We first concentrate on cutting down the intersection between $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ and $\psi_\zeta(\zeta')$. Now $E \subset U^0$, where U^0 is defined in 3.2, and apart from the component of $\alpha_\zeta \cap \{z : |z| > 1\}$ which contains ∞, the loop α_ζ is also contained in U^0. It follows by induction on i that each component of $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ is contained in a component of

$$(\sigma_{\zeta_1} \circ s)^{-i}(U^0),$$

up to isotopy preserving Z_m and S^1, where ζ_1 is the initial segment of ζ, up to the first intersection with S^1. These are the same as the components of

$$(\sigma_{\beta_{2/7}})^{-i}(U^0).$$

Next we show that there are no intersections between this set and the piece of $\psi_\zeta(\zeta')$ which is common with $\psi_{m,2/7}(\beta_{5/7})$ up to S^1 and Z_m-preserving isotopy, where ζ' is the path in $R'_{m,0}$ which is matched with ζ. To do this, we simply show that there are no intersections at all with $\psi_{m,2/7}(\beta_{5/7})$. To do this, since $U^0 \cap Z_2 = \emptyset$, it suffices to show that that $\psi_{m,2/7}(\beta_{5/7})$ is disjoint from T^0 where

$$T^0 = \cup_{i \leq m-3}(\sigma_{\beta_{2/7}} \circ s)^{-i}(U^0).$$

It suffices to prove this by induction on m. Let D_1 be the disc bounded by $\overline{\alpha_{2/7}} * \alpha_{9/28}$, that is, bounded by an arbitrarily small perturbation of $\overline{\beta_{2/7}} * \beta_{5/7} * \overline{\beta_{19/28}} * \beta_{9/28}$. We denote by $\partial' D_1$ the part of the boundary of D_1 which is an arbitrarily small perturbation of $\beta_{5/7} * \overline{\beta_{19/28}}$. The support of $\psi_{2,2/7}$, illustrated in the figures in 3.12 is disjoint from D_1 and so $\partial' D_1 = \psi_{2,2/7}\partial' D_1$. We note that $\psi_{2,2/7}(\beta_{5/7})$ is disjoint from D_1, and $\psi_{3,2/7}(\beta_{5/7})$ is disjoint from $D_1 \cup s^{-1}(D_1)$. Now U^0 is obtained by adjoining to D_1 the region bounded by $\partial' D_1$ and $\psi_{m,2/7}(\partial' D_1)$. It follows that $\psi_{m,2/7}(\beta_{5/7})$ is disjoint from $\psi_{m,2/7}(\partial' D_1 \cup s^{-1}(\partial' D_1)$. But the intersection of $\psi_{m,2/7}(\beta_{5/7})$ with the unit disc can be homotoped to $\partial' U^0 = \psi_{m,2/7}(\partial' D_1)$ via a homotopy preserving Z. Each component of $s^{-i}U^0$ is either contained in, or disjoint from, U^0. It follows that $\psi_{m,2/7}(\beta_{5/7})$ is disjoint from T_0. Hence, any intersection between $\psi_\zeta(\zeta')$ and $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ must be contained in U^0 and coincides, up to isotopy preserving $Z \cup S^1$, with an intersection between $\psi_{m,2/7}(\overline{\beta_{5/7}}) * \psi_\zeta(\zeta')$ and $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$. Following [**22**], we denote by $\partial' U^0$ the subset of ∂U^0 which is isotopic, up to isotopy preserving $Z \cup S^1$, to $\psi_{m,2/7}(\overline{\beta_{5,2/7}}) * \psi_{m,9/28}(\beta_{19/28})$. Then $\psi_{m,2/7}(\overline{\beta_{5/7}}) * \psi_\zeta(\zeta')$ can be isotoped, up to isotopy preserving $Z \cup S^1$, into $\partial' U^0 \cup \zeta$. Hence the same is true for α_ζ. So any intersection between α_ζ and $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ must be contained in $\partial' U^0 \cup \zeta$. But $\partial' U$ has no transversal intersections with $(\sigma_{\beta_{2/7}} \circ s)^{-i}(U^0)$, and hence none with $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ either. (In fact there is at most one component of $(\sigma_{\beta_{2/7}} \circ s)^{-i}(U^0)$ which intersects U^0 without being contained in it, for which the corresponding component of $\sigma_{\beta_{2/7}} \circ s)^{-i}(\beta_{2/7})$ intersects the unit circle at $e^{2\pi i(2/7)}$. This is possible if and only if i is divisible by 3.) It follows that any intersection between α_ζ and $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ is an intersection between ζ and $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$, up to isotopy preserving $Z \cup S^1$. Furthermore, since $\zeta \setminus \zeta_{2k_1}$ coincides with $\psi_\zeta(\zeta')$, all intersections between $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ and α_ζ must be with ζ_{2k_1}.

Suppose that the theorem is true for i, and we consider pre-images of a component C of $(\sigma_{\alpha_\zeta} \circ s)^{-i}(E)$ under $\sigma_{\alpha_\zeta} \circ s$. So by the inductive hypothesis C is a component of $(\sigma_{\zeta_{2k_1-1}} \circ s)^{1-i}(s^{-1}(E) \cup s^{-1}(\zeta_{2k_1-1}))$ and

$$C = E_1 \cup E_2 \cup \omega_{1,n(C),m(C,1)} \cup \omega_{2,n(C),m(C,2)} \cup \ell_C$$

where E_1 and E_2 are components of $s^{-i}E$ and $\omega_r \in R_{m,0}$ if $\omega_{r,n(C),m(C)} \cap U^0 \neq \emptyset$. and $\omega_{r,n(C),m(C)}$ denotes the segment of ω_r from the $m(C)$'th intersection with S^1 to the $n(C,r)$'th. It is possible that $m(C) = n(C,r)$, in which case by convention these segments are absent. If C intersects U^0 then it is contained in U^0, unless $\ell(C)$ passes over $e^{2\pi i(2/7)}$. In that case, exactly one of the sets $E_r \cup \omega_{r,n(C),m(C,r)}$ intersect U^0. We assume without loss of generality that this is true for $r = 1$. If this happens then $w_1'(\omega_1) \neq w_1'(\zeta)$, and $E_1 \cap \zeta = \emptyset$. In fact we have $s^{-i}E \cap \zeta_1 = \emptyset$ for all$i > 0$. Let the first unit disc crossings of ζ^1 and ζ^3 be represented by

$$D(v_1) = D((L_3L_2R_3)^{k_1}u_1(L_2R_3L_3)^{\infty})$$

and

$$D(v_2) = D((L_3L_2R_3)^{k_2}u_2(L_2R_3L_3)^{\infty})$$

respectively, where k_1 and k_2 are maximal. Then $k_1 = k_2$ or $k_2 = k_1 \pm 1$ and $D(v_2)$ is strictly between $D(v_1)$ and $D((L_3L_2R_3)^{\pm 1}v_1)$. This is all that is needed.

Since any intersection between ζ_{2k_1} and $\ell(C)$ must occur outside the unit disc, it cannot occur on $\zeta_{2k_1} \setminus \zeta_{2k_1-1}$, and must occur on ζ_{2k_1-1}. There are no transversal intersections between ζ_{2k_1} and $\omega_{r,n(C),m(C,r)}$, because if $\omega_{r,n(C),m(C,r)}$ intersects U^0 then ω_r is a path in $R_{m,0}$ and cannot have transversal intersections with $\zeta \in R_{m,0}$, because paths in $R_{m,0}$ have no transversal intersections [**22**]. It remains to consider how ζ_{2k_1} can intersect $E_1 \cup E_2$. By 4.5, there are no intersections between ∂E and $\partial E_1 \cup \partial E_2$ if $(\zeta^j : 1 \leq j \leq 4)$ is an upper or lower quadruple. and $\zeta_{2k_1} \setminus \zeta_{2k_1-1}$ is in ∂E Since E_1 and E_2 are contained in the unit disc, and $\zeta_{2i+1} \setminus \zeta_{2i}$ is in the exterior for all $i \geq 1$, the only possible intersections between ζ_{2k_1} and $E_1 \cup E_2$ are with ζ_{2k_1-2}. We now show that no intersection is possible in these cases. The case of $k_1 = 1$ is dealt with since then $2k_1 - 2 = 0$ and ζ_0 is empty. So we can assume that $k_1 > 1$. The subsets E_1 and E_2 of C are bounded by paths ω_r and ω_{r+2} in $R_{m,0}$ (for $r = 1$ and 2), where ω_1 and ω_2 are as above. By the invariance properties, $k(\omega_r, \omega_{r+2}) \geq k(\zeta^1, \zeta^3) \geq k_1 - 1$ or $\geq k_1$, depending on whether the quadruple $(\zeta^i : 1 \leq i \leq 4)$ is of type A or of type B. We can extend the notion of type A or B to lower quadruples, and if $k_1 = 2$ then $(\zeta^i : 1 \leq i \leq 4$ is a type A quadruple. Suppose, for contradiction, that ζ_{2k_1} intersects the component of E_r between ω_r and ω_{r+2}. This is certainly impossible if $w_1'(\zeta) = w_1'(\omega_r)$ because then ζ_{2k_1-2} or ζ_{2k_1-4} has to coincide with $\omega_{r,2k_1-2}$ or $\omega_{r,2k_1-4}$ (or $\omega_{r+2,2k_1-2}$ or $\omega_{r+2,2k_1-4}$). WE necessarily have ζ_{2k_1-2} coinciding with $\omega_{r,2k_1-2}$ or $\omega_{r+2,2k_1-2}$ if $k_1 = 2$, because then $(\zeta^j : 1 \leq j \leq 4)$ is of type A. It is also impossible if $w_1'(\zeta) \neq w_1'(\omega_r)$ because although paths ζ and ω in $R_{m,0}$ can occasionally intersect if $w_1'(\zeta) \neq w_1'(\omega)$, intersections can only be between r'th components of intersection with the exterior of the unit disc for the same r. So ζ_{2k_1}cannot intersect $E_1 \cup E_2$ This completes the proof for upper and lower quadruples.

Now we consider the case of $(\zeta^i : 1 \leq i \leq 4)$ being of type C or AC. There is no intersection if $k(\omega_r, \omega_{r+2}) > k_1$. Since $k(\zeta, \zeta^3) = k_1$ the only possibility is that $k(\omega_r, \omega_{r+2}) = k_1$. If $k_1 > 1$ then there can only be an intersection if $w_1'(\omega_j) = y$ and $w_1'(\zeta) = x$ and $U^x \cap U^y \neq \emptyset$. But y also has to be in $s^{-i}(x)$, and this is never true in the cases when $U^x \cap U^y \neq \emptyset$, except in the case $i = 0$. So the only possibility is that $k_1 = 1$. The proof for type AC is then exactly as for type A, so now we consider type C only, and we write $w_1'(\zeta^3) = uL_3$ and $w_2'(\zeta^3) = uau'$ for a basic exchangeable pair (a, b). Then $D(w_1'(\zeta)(L_2R_3L_3)^\infty)$ is between $D(uL_3(L_2R_3L_3)^\infty)$ and $D(ubL_3L_2)$ and is maximal to the right of $D(ubL_3L_2)$. If $D(w_1'(\zeta)(L_2R_3L_3)^\infty)$ is also between $D(vuL_3(L_2R_3L_3)^\infty)$ and $D(vubL_3L_2)$ then u must be a prefix of vu and so $u = v^n v'$ for some $n \geq 1$ or $v = uu''$. But $D(ubL_3L_2)$ also has to be in $D(vu)$. So $v^{n+1}v'$ has to be a prefix of $v^n v' bL_3L_2$. So vv' has to be a prefix of $v'bL_3L_2$. This is impossible because $w_1(vv') = vv'$ and $w_1(v'bL_3L_2)$ is a proper prefix of vbL_3L_2.

□

4.9. The first four elements of the β_i sequence

In this section we prove Theorem 2.8 for the first four elements of the sequence $\omega_i = \omega_i(\beta)$, given a capture path $\beta \in \mathcal{Z}_m(3/7, +, +, 0)$. As in 2.8, we let γ_i be the sequence of elements of $\pi_1(V_{3,m}, a_1)$ and $(\omega_{2i-1}, \omega_{2i})$ the sequence of adjacent pairs in Ω_m such that $\rho(\beta)$ successively crosses $\rho(\gamma_i * \omega_{2i-1}) * \rho(\gamma_i * \omega_{2i})$.

Lemma 4.9.1. *Then there is a quadruple* $(\beta_i : 1 \leq i \leq 4)$ *of type A, AC or C such that*

$$(\rho(\omega_1), \rho(\omega_2)) = (\beta_1, \beta_2),$$
$$(\rho(\omega_3), \rho(\omega_4)) = (\beta_3', \beta_4'),$$

where (β_3', β_4') *is the adjacent pair in* $R'_{m,0}$ *which is matched with* (β_3, β_4). *Moreover the following holds. Either* $\psi_{1,3,\beta} = \psi_{\beta_1,\beta_3}$ *preserves* $D(\beta_1, \beta_2)$ *or there is a single basic exchange* $a \leftrightarrow b$ *and prefix* v *such that* $C(va \leftrightarrow vb, D(X)$ *intersects both* $D(\beta_1, \beta_2)$ *and its complement, where* $X = L_3$ *or* BC, *depending on whether* $a \leftrightarrow b$ *is a basic exchange for* L_3 *or* BC. *If* $(\beta_i : 1 \leq i \leq 4)$ *is of type C or AC then* $a \leftrightarrow b$ *is the basic exchange occurring in the definition of types C and AC.*

Proof. We already know that $(\beta_1, \beta_2) = (\zeta_1(x), \zeta_2(x))$, where $(\zeta_i(x))$ is the ζ-sequence for x. Write $\zeta_i(x) = \zeta_i$ and let $(\zeta_{2i-1}', \zeta_{2i}')$ be the adjacent pair in $R)_{m,0}$ which is matched with $(\zeta_{2i-1}, \zeta_{2i})$. Write $E = E_{\zeta_1,\zeta_3}$. Clearly, if

$\zeta_i(x)$ is defined for $i = 3$, then $(\rho(\omega_3), \rho(\omega_4)) = (\zeta_3', \zeta_4')$ and $(\zeta_3, \zeta_4) = (\beta_3, \beta_4)$ has the required properties, unless $I(\zeta_3, \zeta_4)$ intersects the support of ψ_{ζ_1,ζ_3}. The only way that this can happen is if $C(u \leftrightarrow u', E)$ intersects $I(\zeta_3, \zeta_4)$ for some exchange $u \leftrightarrow u'$ for ζ_1 which is prefixed by some basic exchange $va \leftrightarrow vb$. The only way in which $C(va \leftrightarrow vb, D(L_3L_2))$ (or $C(va \leftrightarrow vb, D(BC))$) cannot be contained in $D(\zeta_1, \zeta_2) = D(\beta_1, \beta_2)$ is if $a \leftrightarrow b$ is one of the exchanges of (3.9.1.4) (or (3.9.1.5)). If one of these happens then we have

$$D(u'X) \cap D(\beta_1, \beta_2) \neq \emptyset$$

where $X = L_3L_2$ or $X = BC$, as appropriate. Let ξ_1 be the exchange supported by $C(u \leftrightarrow u', E_{\beta_1,\zeta_3})$ if $\zeta_i(x)$ is defined for $i \geq 3$ and supported by $C(u \leftrightarrow u', D(L_3L_2))$ if $\zeta_3(x)$ is not defined fo $i \geq 3$. In both cases there is an adjacent pair (β_3, β_4) in $R_{m,0}$ such that $I(\beta_3, \beta_4) \subset D(uJL_3L_2)$ and $\xi_1(I(\beta_3, \beta_4))$ is under $I(\beta_1, \beta_2)$ and not separated from it by anything else in the image of ξ_1. If $\zeta_i(x)$ is defined for $i \geq 3$ then ξ_1 interchanges $I(\zeta_3(x), \zeta_4(x)) = I(\zeta_3, \zeta_3)$ and $I(\beta_3, \beta_4)$, and also $E_{\beta_1,\zeta_3} \subset E_{\beta_1,\beta_3}$. In any case if ξ_1' is the exchange supported by $C(u \leftrightarrow u', E_{\beta_1,\beta_3})$, then

$$\xi_1'(I(\beta_3, \beta_4)) = \xi_1(I(\beta_1, \beta_3).$$

Also, since $I(\beta_3, \beta_4)$ is in the boundary of E_{β_1,β_3} it is clear that $I(\beta_3, \beta_4)$ is not in any component of $s^{-n}(E_{\beta_1,\beta_3})$ for any $n > 0$, apart from uE_{β_1,β_3}. It follows that

$$\rho(\omega_i) = \beta_i' \text{ for } i = 3,\ 4,$$

where (β_3', β_4') is the adjacent pair of $R_{m,0}'$ which is matched with (β_3, β_4). So in all cases we have a quadruple $(\beta_i : 1 \leq i \leq 4)$ of type A, C or AC with the required properties. It is also true that

$$\begin{aligned} &D(\beta_3, \beta_4) \subset uD(X)), \\ &u'D(\beta_3, \beta_4) \subset D(\beta_1, \beta_2) \end{aligned} \tag{4.9.1.1}$$

where $X = L_3L_2$ or BC, as appropriate.

We claim that there is at most one basic exchange $va \leftrightarrow vb$ such that $C = C(va \leftrightarrow vb, D(X))$ intersects both $D(\zeta_1, \zeta_2) = D(\beta_1, \beta_2)$ and its complement. We see this as follows. The set C never intersects $I(\zeta_1, \zeta_2)$ because there are no values of w_1' between $w_1'(\zeta_1)$ and $w_1'(\zeta_2)$, and $w_1'(\zeta)$ is never between two values of $w_i'(., y)$ for any value y of w_1' and $i \geq 1$. We only need to consider exchanges $va \leftrightarrow vb$ as in (3.9.1.4) for which $w_1'(vaL_3L_2) \neq w_1'(vbL_3L_2)$, or as in (3.9.1.5) for which $w_1(vaL_3L_2) \neq w_1(vbL_3L_2)$. For all of these exchanges, since there are no intersections with $I(\zeta_1, \zeta_2)$, we only need to consider those with $w_1(vaL_3L_2) = vaL_3$ and $w_1(vbL_3L_2) = vbL_3L_2$. This condition ensures that for any two such, say $v_1a_1 \leftrightarrow v_1b_1$ and $v_2a_2 \leftrightarrow v_2b_2$, neither of $v_1a_1L_3L_2$ or $v_1b_1L_3L_2$ can be a proper prefix of $v_2a_2L_3L_2$ or $v_2b_2L_3L_2$. It follows that if these are distinct basic pairs then

$$(D(v_1a_1L_3L_2) \cup D(v_1b_1L_3L_2)) \cap (D(v_2a_2L_3L_2) \cup D(v_2b_2L_3L_2)) = \emptyset.$$

Now we show that $D(v_1b_1L_3L_2)$ and $D(v_2b_2L_3L_2)$ are not in $D(\zeta_1, \zeta_2)$ for the same pair $I(\zeta_1, \zeta_2)$. But if $D(v_1b_1L_3L_2) \subset D(\zeta_1, \zeta_2)$ then $w_1'(\zeta_1)$ must be the maximal value of w_1' to the right of $D(v_1b_1L_3L_2)$ which means that $w_1'(\zeta_1)$ has a common prefix with v_1b_1 of length $|v_1b_1| - 1$. It is not possible for this to be true also for v_2b_2 if $v_2b_2 \neq v_1b_1$. So the claim is proved.

If $va \leftrightarrow vb$ exists as in (3.9.1.4) then, using the same argument, the only basic exchange $v_2a_2 \leftrightarrow v_2b_2)$ for which the homeomorphism ξ supported by $C(v_2a_2 \leftrightarrow v_2b_2)$ does not preserve $vaD(\zeta_1, \zeta_2)$ is $vava \leftrightarrow vavb$. By induction it follows that ψ_{β_1,β_3} preserves $\cup_{n\geq 0}(va)^nD(v)$

□

4.10. A general property of β quadruples

In order to prove Theorem 2.8 in its entirety, we need some inductive information about the form that a quadruple $(\beta_{4i+t}(\beta) : -3 \leq t \leq 0)$ can take, assuming the existence of $\beta_j(\beta)$ as in 2.8 for $j \leq 4i$.

Lemma 4.10.1. *Corollary 4.7.2 and Lemma 4.8.2 remain true if we generalise to include quadruples* $(\zeta^t : 1 \leq t \leq 4)$ *of the following form. There are two quadruples* $(\zeta^i : -1 \leq i \leq 2)$ *and* $(\zeta^{-1}, \zeta^0, \zeta^3, \zeta^4)$ *which are either both* ζ*quadruples or both type C, and* $E = E_{\zeta^1,\zeta^3}$ *is defined to be* $E_{\zeta^{-1},\zeta^1} \setminus E_{\zeta^1,\zeta^3}$*. Also any component of* $(\sigma_{\alpha_\zeta} \circ s)^{-n}(E)$ *coincides with a component of* $(\sigma_{\zeta_k} \circ s)^{-n}(E)$*, where* ζ_k *is the longest common segment of* ζ^1 *and* ζ^3*.*

Proof. First we consider 4.7.2. Without loss of generality we have $E_{\zeta^1,\zeta^3} \subset E_{\zeta^{-1},\zeta^3}$, and hence any component of $s^{-n}(E_{\zeta^1,\zeta^3}$ is contained in the corresponding component of $s^{-n}(E_{\zeta^{-1},\zeta^3})$. By 4.7.1, any component of $s^{-n}(E_{\zeta^{-1},\zeta^3}$ is either contained in or disjoint from each of E_{ζ^{-1},ζ^3} and E_{ζ^{-1},ζ^1}. In the latter case we use the fact that since it is not possible for a component of $s^{-n}(E_{\zeta^{-1},\zeta^3})$ to contain E_{ζ^{-1},ζ^1}, for $n > 0$. So any component of $s^{-n}(E_{\zeta^{-1},\zeta^3}$ is either contained in or disjoint from $E_{\zeta^1,\zeta^3} = E_{\zeta^{-1},\zeta^3} \setminus E_{\zeta^1,\zeta^3}$. So the same is true for any component of $s^{-n}(E_{\zeta^1,\zeta^3}$

The proof of the analogue of 4.8.2 is essentially the same as the proof of 4.8.2. The proof that any component of $\sigma_{\alpha_\zeta} \circ s)^{-n}(E)$ coincides with a component of $\sigma_{\zeta_k} \circ s)^{-n}(E)$ is the same as before. For the remainder of the statement of 4.8.2, we only need to check that ζ_k does not cut through the interior of components of $s^{-n}(E)$ Since $E \subset E_{\zeta^{-1},\zeta^3}$, this is the same as before. □

THEOREM 4.10.2. *Let* $\beta \in \mathcal{Z}_\infty(3/7,+,+,0)$ *with endpoint* $p(x)$. *Suppose that* $\beta_j(\beta)$ *exists as in 2.8 for any* $j \leq 4i$. *For any* $i \geq 1$, *let* $\psi_{1,4i-1,\beta} = \xi_n \circ \cdots \circ \xi_1$, *where each* ξ_ℓ *is a disc exchange with support* $C(v_\ell \leftrightarrow v'_\ell, E_\ell)$, *where* $E_\ell = E_{4j-3,4j-1,\beta}$ *for some* $j < i$ *depending on* ℓ. *For any* $1 \leq \ell$, *let* $\psi = \xi_\ell \circ \cdots \xi_1$. *Each component* J_1 *of* $\psi(\overline{\tau_1} * \tau_2) \cap \{z : |z| \geq 1\}$ *over* x, *where* (τ_1, τ_2) *runs over all adjacent pairs* (τ_1, τ_2)*in* $R_{m,0}$, *is paired with some other arc* J_2 *which is similarly defined with respect to some adjacent pair* (τ_3, τ_4), *with the following properties.*

1. *Paired arcs* J_1 *and* J_2 *both cross* β, *with one inside the other.*
2. *Choosing the numbering so that arcs of* $\psi(\tau_1)$ *and* $\psi(\tau_3)$ *adjoin* J_1 *and* J_2 *at one end, there is no path in* $R_{m,0}$ *which cuts between* $\psi(\tau_1)$ *and* $\psi(\tau_3)$, *including cutting between* J_1 *and* J_2.
3. *Every pair of arcs* J_1 *and* J_2 *is of one of the following forms:*
   - *a) subarcs of* $\psi(\overline{\tau_1} * \tau_2)$ *and* $\psi(\overline{\tau_3} * \tau_4)$, *in either order, where* $(\tau_i : 1 \leq i \leq 4)$ *is a quadruple or lower quadruple or type C or AC;*
   - *b) subarcs of* $\psi(\overline{\tau_3} * \tau_4)$ *and* $\psi(\overline{\tau_1} * \tau_2)$ *where, for some adjacent pair* (τ_{-1}, τ_0), *both* $(\tau_i,\ -1 \leq i \leq 2)$ *and* $(\tau_{-1}, \tau_0, \tau_3, \tau_4)$ *are upper quadruples, or both type C or AC;*

Remark We shall refer to J_1 and J_2 as being *paired over* x. Of course a local S inverse of s^n often maps an upper or lower ζ quadruple $(\zeta_{4j+t}(x_1) : -3 \leq t \leq 0)$ to another upper or lower ζ quadruple $(\zeta_{4\ell+t}(x_2) : -3 \leq t \leq 0)$. Since S might map the upper unit circle into the lower unit circle, upper quadruples can be mapped to lower quadruples, and vice versa.

PROOF. The proof is by induction on i and ℓ. We assume inductively that $i = 1$ or that $i > 1$ and that the statement is already proved for $i - 1$ replacing i. By 4.9.1 to 4.12.3, we already know that $(\beta_t(\beta) : 1 \leq t \leq 4)$ is a quadruple of type A or C or AC.

So now we need to prove the lemma for i, and for each $\ell \geq 1$. We use induction on ℓ. So let ψ' be the identity if $\ell = 1$ and $\psi' = \xi_{\ell-1} \circ \cdots \circ \xi_1$ if $\ell > 1$. Writing $\xi = \xi_\ell$, we assume inductively that the lemma is true for ψ' replacing ψ — which is trivial if ψ' is the identity — and prove it for $\psi = \xi \circ \psi'$. Let the support of ξ be $C = C(v \leftrightarrow v', E)$, where $vE = E_{\delta_1,\delta_3}$ and $v'E = E_{\delta'_1,\delta'_3}$, where $(\delta_i : 1 \leq i \leq 4)$ and $(\delta'_i : 1 \leq i \leq 4$ are both quadruples of the inductive type. This means that either $(\delta_i : 1 \leq i \leq 4)$ is a ζ quadruple, or of type C or AC, or there are two quadruples $(\delta_i : -1 \leq i \leq 2)$ and $\{\delta_{-1}, \delta_0, \delta_3, \delta_4)$ which are both upper ζ quadruples or both type C, and $E_{\delta_1,\delta_3} = E_{\delta_{-1},\delta_1} \setminus E_{\delta_{-1},\delta_3}$, and similar properties hold for $(\delta'_i : 1 \leq i \leq 4)$. We know from 4.8.2 that the connecting arc of C between vE and $v'E$, apart from the central arc, consists of common arcs from each of the quadruples

$(\delta_i : 1 \leq i \leq 4)$ and $(\delta'_i : 1 \leq i \leq 4)$. Let J'_1 and J'_2 be paired sub-arcs of $\psi'(\overline{\tau'_1} * \tau'_2)$ and $\psi'(\overline{\tau'_3} * \tau'_4)$ where $(\tau'_i : 1 \leq i \leq 4)$ satisfies the inductive hypothesis.

If C does not intersect $J'_1 \cup J'_2$, then there is nothing to prove. If neither vE nor $v'E$ intersects $J'_1 \cup J'_2$, but C does intersect $J'_1 \cup J'_2$, then the intersection has to be of the arc of C between vE and $v'E$, with $J'_1 \cup J'_2$. But, using the inductive hypothesis, by property 2 of the lemma for $J'_1 \cup '_2$, no part of the connecting arc of C can separate J'_1 and J'_2.

If vE or $v'E$ contains $J'_1 \cup J'_2$, then $\xi(J'_1 \cup J'_2)$ is a single pair of arcs $J_1 \cup J_2$ with the required properties. In fact (J_1, J_2) is a paired arc of exactly the same type as (J'_1, J'_2) because ξ maps quadruples in vE to quadruples in $v'E$.

So it remains to consider what happens when one of vE or $v'E$ intersects $J'_1 \cup J'_2$ but does not contain it. Suppose without loss of generality that vE intersects, but does not contain, $J'_1 \cup J'_2$. Then, by the inductive hypothesis, there is a ζ quadruple $(\eta'_i : 1 \leq i \leq 4)$ such that J'_1 and J'_2 both intersect $E_{\eta'_1,\eta'_3}$ and $vE = E_{\delta_1,\delta_3}$ also intersects $E_{\eta'_1,\eta'_3}$. Also by the inductive hypothesis on J'_1 and J'_2, we have $J'_2 = I(\eta'_3, \eta'_4)$, and either $J'_1 = I(\eta'_1, \eta'_2)$, or J'_1 intersects the region between J'_2 and the nearest segment of one of η'_1 and η'_2 to $I(\eta'_1, \eta'_2)$. By the inductive hypothesis on vE we have $(\delta_1, \delta_2) = (\eta'_1, \eta'_2)$ or (δ_{-1}, δ_0) depending on whether or not $(\delta_i : 1 \leq i \leq 4)$ is itself a ζ quadruple or type C quadruple. So then there are various configurations to consider. The figure shows the configuration when $(\delta_i : 1 \leq i \leq 4)$ is a ζ quadruple and ψ' is the identity on $vE = E_{\delta_1,\delta_3}$ — which means that ψ' is also the identity on vE — and the new pairing is of (δ_3, δ_4) with $(\eta'_3, \eta'_4) = (\tau'_3, \tau'_4)$ – and also of (δ_3, δ_4) with $(\eta'_1, \eta'_2) = (\tau'_1, \tau'_2)$, but that is not shown in the picture. Now suppose that ψ' is not necessarily the identity on vE. If there is an intersection between vE and $J'_1 \cup J'_2$ then necessarily at least one of $I(\delta_3, \delta_4)$ and $I(\delta_1, \delta_2)$ is under J'_1, say $I(\delta_3, \delta_4)$. If $I(\delta_3, \delta_4) \subset \psi'(E_{\tau'_1,\tau'_3})$, then $I(\delta'_3, \delta'_4) = \psi'(I(\tau'_5 \tau'_6)$ for some $I(\tau'_5, \tau'_6) \subset E'$, where E' is in the support of ψ' and $E_{\tau'_1,\tau'_3} \subset E'$, where $(\tau'_5, \tau'_6, \tau'_3, \tau'_4)$ is another quadruple of the same type, and the pairing for $\xi \circ \psi' = \psi$ is of $\xi(J'_2)$ with an arc of $\xi \circ \psi'(I(\tau'_5, \tau'_6))$. If $I(\delta_3, \delta_4)$ is not in $\psi'(E')$ then it is separated from $\psi(E')$ by $\psi'(I(\tau'_5, \tau'_6))$ for some such (τ'_5, τ'_6), and the new pairing is still of (τ'_5, τ'_6) with each of (τ'_3, τ'_4) and (τ'_1, τ'_2). Then $(\tau'_i : 3 \leq i \leq 6)$ and $(\tau'_1, \tau'_2, \tau'_5, \tau'_6)$ are quadruples of the required types, and this completes the proof of the inductive hypothesis.

□

4.10.3. Satellite arcs. We have seen in 4.10.2 that there are a number of ways in which paired principal arcs for $[\psi]$ and $R_{m,0}$ can differ from a ζ quadruple, up to isotopy preserving $S^1 \cup Z_\infty$. We have seen that if



FIGURE 5. Pairing of type 3b)

$\psi = \psi_{1,4i-1,\beta}$ for any capture path $\beta \in \mathcal{Z}(3/7,+,+,0)$, the paired principal arcs for $\psi(R_{m,0})$ have no transversal intersection with principal arcs for $R_{m,0}$, and and pairs of principal arcs for $\psi(R_{m,0})$ are not separated by any other principal arcs. If both endpoints coincide, then we say that this is a pair of *satellite arcs (for ψ)*, or a *satellite pair.*Pairs of arcs which are not a satellite pair are called a *non-satellite pair.* If $I(\psi_{1,4i-5,\beta}(\beta_{4i-3}),\psi_{1,4i-5,\beta}(\beta_{4i-2}(\beta))$ and $I(\psi_{1,4i-1,\beta}(\beta_{4i-1}),\psi_{1,4i-1,\beta}(\beta_{4i}(\beta))$ are a satellite (or non-satellite) pair over $p(w(\beta))$, then we call $(\beta_{4i+t}(\beta):-3\leq t\leq 0)$ a *satellite quadruple* (or *non-satellite quadruple.*

COROLLARY 4.10.4. *Assume the existence of $\beta_j(\beta)$ with the properties of Theorem 2.8 for $j\leq 4i$. Then $\psi_{4i-3,4i-1,\beta}$ is the identity on $I(\beta_{4i-1}(\beta),\beta_{4i}(\beta))$ for all $i\geq 2$, and $(\beta_{4i-3}(\beta),\beta_{4i-2}(\beta))$ and $(\beta_{4i-1}(\beta),\beta_{4i}(\beta))$ are the adjacent pairs in $R_{m,0}$ whose principal arcs for $[\psi_{1,4i-5,\beta}]$ are the next two inside the principal arc of $(\beta_{4i-5}(\beta),\beta_{4i-4}(\beta))$ for $[\psi_{1,4i-5,\beta}]$ which is crossed by β.*

PROOF. By 4.7.2, 4.10.1 and 4.10.2, then any component of $s^{-n}(E_{4i-3,4i-1,\beta})$, for any $n>0$, is either strictly contained in $E_{4i-3,4i-1,\beta}$ or is disjoint from it. Hence it cannot intersect $I(\beta_{4i-1}(\beta),\beta_{4i}(\beta))$, which is in the boundary. So if $C(v\leftrightarrow v',E)$ is the support of any disc exchange in the composition for $\psi_{4i-3,4i-1,\beta}$ then vE and $v'E$ are disjoint from $I(\beta_{4i-1}(\beta), \beta_{4i}(\beta))$. There are also no transversal intersections between $I(\beta_{4i-1}, \beta_{4i})$ and the two segments of paths in$R_{m,0}$ in C and no intersections between $I(\beta_{4i-1}, \beta_{4i})$ and the central arc $\ell(C)$ of C . ☐

4.11. Satellite arcs

It will be clear from 4.10.2 that satellite arcs are formed by the action of disc exchanges. From the form given for the support of a disc exchange ξ in 4.8.2, and 4.10.1, we see that satellite arcs for $\psi_{1,4i-1,\beta}$ are always parallel to components of

$$(\omega_{1,n(C),m(C)} \cup \omega_{2,n(C),m(C)} \cup \ell(C)) \setminus S^1$$

for varying C, where C is the support of some disc exchange in the composition for $\psi_{1,4i-1,\beta}$. To see this , we need to use the fact that $w_1'(\beta_i)$ is constant for $i \geq 3$. So then there are never transverse intersections between $\ell(C)$ and $\ell(C')$, or between $\omega_{i,n(C),m(C)}$ and $\omega_{j,n(C'),m(C')}$, for different C and C', and for i, $j \in \{1, 2\}$. So satellite arcs are always parallel either to arcs $\ell(C)$ or to segments in the exterior of the unit disc of paths in $R_{m,0}$, not including the first segments of such paths.

4.12. Final proof of Theorem 2.8

We are now finally ready to prove Theorem 2.8 in full generality.We do this for the different cases of the quadruple $(\beta_i(\beta) : 1 \leq i \leq 4)$. Let $\beta \in |calZ_m(3/7, +, +, 0)$ In what follows, we use the notation $E_{4j-3,4j-1,\beta}$ for the set $E_{\beta_{4j-3},\beta_{4j-1},\beta}$ if $2j \leq i$ and the existence of the sequence $\beta_\ell(\beta)$ for $\ell \leq 2i$ as in Theorem 2.8 has been established. We write $E_{4j-3,4j-1}$ for $E_{4j-3,4j-1,\beta}$ if no confusion can arise.

THEOREM 4.12.1. *Let $(\beta_i : 1 \leq i \leq 4)$ be of type A. Then $(\beta_1, \beta_2), (\beta_3, \beta_4))$ extends to a sequence $(\beta_{2i-1}, \beta_{2i})$ of adjacent pairs in $R_{m,0}$ such that, if γ_i is the sequence of elements of $\pi_1(V_{3,m}, a_1)$ and $(\omega_{2i-1}, \omega_{2i})$ the sequence of adjacent pairs in Ω_m such that $\rho(\beta)$ successively crosses $\overline{\rho(\gamma_i * \omega_{2i-1})} * \rho(\gamma_i * \omega_{2i})$, then*

$$(\rho(\omega_{2i-1}), \rho(\omega_{2i})) = \begin{cases} (\beta_{2i-1}, \beta_{2i}) \text{ if } i \text{ is odd,} \\ (\beta'_{2i-1}, \beta'_{2i}) \text{ if } i \text{ is even,} \end{cases}$$

where $(\beta'_{2i-1}, \beta'_{2i})$ is the adjacent pair in $R'_{m,0}$ matched with $(\beta_{2i-1}, \beta_{2i})$. Then $I(\beta_{2j-1}, \beta_{2j})$ is contained in $D(\beta_1, \beta_2)$ for all $j \geq 2$.

PROOF. Naturally, this is proved by induction on j. It is true for $j = 2$ by the assumption that $(\beta_j : 1 \leq j \leq 4)$ is of type A. Using 4.8.2, this means that $I(\beta_3, \beta_4)$ is disjoint from $(\sigma\beta_1 \circ s)^{-n}(E_{1,3})$ for all $n > 0$, and $I(\beta_3, \beta_4) \subset D(\beta_1, \beta_2)$. Suppose that we have proved the Theorem for β_j for $j \leq 4i$. Then we consider $\psi_{1,4i-1,\beta}$. We claim that $\psi_{1,4i-1,\beta}$ does not map any principal arc under $I(\psi_{1,4i-1,\beta}(\beta_{4i-1}),\psi_{1,4i-1,\beta}(\beta_{4i})$ out of $D(\beta_1,\beta_2)$. To see this, as usual, we write

$$\psi_{1,4i-1,\beta}=\psi_{1,3,\beta}\circ\cdots\circ\psi_{4i-3,4i-1,\beta}$$

for $1\le j\le i$, and we write $\psi_{4j-3,4j-1,\beta}$ as a composition of disc exchanges with supports $C(u_{j,\ell}\leftrightarrow u'_{j,\ell},E_{4j-3,4j-1,\beta}$ where $E_{4j-3,4j-1,\beta}=E_{\beta_{4j-3}(\beta),\beta_{4j-1}(\beta)}$ and $u_{j,\ell}\leftrightarrow u'_{j,\ell}$ is an exchange for β_{4j-3}. This means $u_{j,\ell}\leftrightarrow u_{j,\ell'}$ is prefixed by a basic exchange $v_{j,\ell}a_{j,\ell}\leftrightarrow v_{j,\ell}b_{j,\ell}$, where the first letters of $a_{j,\ell}$ and $b_{j,\ell}$ are different, that is, each $a_{j,\ell}\leftrightarrow b_{j,\ell}$ is one of the basic exchanges listed in 3.9.1. From what we now know about the quadruple $(\beta_{4j+t}(\beta):-3\le t\le 0)$ from 4.10.2, and from the inductive hypothesis, we have

$$E_{4j_1-3,4j_1-1}\cap(L_3L_2R_3)^nE_{4j_2-3,4j_2-1,\beta}=\emptyset$$

for all $n>0$ and all j_1 and j_2. It follows that we can only have a nontrivial intersection between $u_{j_1,\ell_1}E_{4j_1-3,4j_1-1,\beta}$ and $u_{j_2,\ell_2}E_{4j_2-3,4j_2-1,\beta}$ if one of the following holds.

- $v_{j_1,\ell_1}=v_{j_2,\ell_2}$ and $\{a_{j_1,\ell_1},b_{j_1,\ell_1}\}=\{a_{j_2,\ell_2},b_{j_2,\ell_2}\}$;
- One of $v_{j_1,\ell_1}a_{j_1,\ell_1}$ or $v_{j_1,\ell_1}b_{j_1,\ell_1}$ is a prefix of v_{j_2,ℓ_2} or vice versa.

We also know from 4.9.1 that there is at most one basic exchange $a\leftrightarrow b$ and prefix v such that $C(a\leftrightarrow vb,D(X))$ which intersects both $D(\beta_1,\beta_2)$ and its complement. If $a\leftrightarrow b$ and v do not exist, then $\psi_{1,4i-1,\beta}$ preserves $D(\beta_1,\beta_2)$. So suppose that $a\leftrightarrow b$ and v do exist. Then there is an exchange $u\leftrightarrow u'$ for β_1 which is prefixed by $va\leftrightarrow vb$ and $u'E_{1,3}\cap D(\beta_1,\beta_2)\ne\emptyset$ (choosing u' without loss of generality). Then by hypothesis, since we are considering type A quadruples $I(\beta_3,\beta_4)\cap u'E_{1,3}=\emptyset$, and then we also have

$$D(\beta_3,\beta_4)\cap u'(E_{1,3}\cup D(\beta_1,\beta_2)=\emptyset.$$

Hence if (η_1,η_2) is an adjacent pair such that $I(\eta_1,\eta_2)\subset D(\beta_3,\beta_4)$ and $\psi_{1,4j-1}^{-1}(I(\eta_1,\eta_2))$ is not contained in $D(\beta_3,\beta_4)$, then the first disc exchange in the composition for $\psi_{1,4i-1,\beta}^{-1}$ which moves $I(\eta_1,\eta_2)$ out of $D(\beta_3,\beta_4)$ must have support $C(u_{j,\ell}\leftrightarrow u'_{j,\ell}E_{4j-3,4j-1})$ where $v_{j,\ell}a_{j,\ell}$ and $v_{,\ell}b_{j,\ell}$ do not have va or vb as a prefix, and similarly with the roles reversed. So then further disc exchanges cannot move out of $D(\beta_1,\beta_2)$. It follows that for any $j\le i$ we have

$$\psi_{1,4i-1}^{-1}(I(\eta_1,\eta_2)\subset D(\beta_3,\beta_4)\cup D(v_{j,\ell}a_{j,\ell})\cup D(v_{j,\ell}b_{j,\ell})$$

for some (j,ℓ) and $\{a_{j,\ell},b_{j,\ell}\}\ne\{a,b\}$. If

$$I(\eta_1,\eta_2)\subset D(v_{j,\ell}a_{j,\ell})\cup D(v_{j,\ell}b_{j,\ell})$$

for $\{a_{j,\ell},b_{j,\ell}\}\ne\{a,b\}$ then it cannot be moved into the domain of the disc exchange with support $C(va\leftrightarrow vbD(X))$ by $\psi_{1,4i-1,\beta}^{-1}$ It follows that we have a quadruple $(\beta_{4i+t} : 1 \leq t \leq 4)$ such that

$$(\rho(\omega_{4i+1}), \rho(\omega_{4i+2})) = (\beta_{4i+1}, \beta_{4i+2}),$$

$$(\rho(\omega_{4i+3}), \rho(\omega_{4i+4})) = (\beta'_{4i+3}, \beta'_{4i+4})$$

and $(\beta_{4i+t} : 1 \leq t \leq 4)$ has all the required properties.

$\square$

Now we deal with the cases of $(\beta_i : 1 \leq i \leq 4)$ being a quadruple of type C or AC

Theorem 4.12.2. *Let $\beta \in \mathcal{Z}_m(3/7, +, +, 0)$ and let $p(\beta)$ denote the endpoint of β. Let $(\beta_i(\beta) : 1 \leq i \leq 4) = (\beta_i : 1 \leq i \leq 4)$ be of AC. Let $va \leftrightarrow vb$ be the basic exchange as in 4.9.1, prefixing the exchange $u \leftrightarrow u'$ for $\beta 1$, also as in 4.9.1. Then βj exists as in Theorem 2.8*

$$D(\beta_{2j-1}, \beta_{2j}) \subset (D(\beta_1, \beta_2) \cap E_{1,3}) \cup_{1 \leq i} u^i(E_{1,3}$$

for all j.

Remark We automatically have $n \geq 1$, as $(\beta_i : 1 \leq i \leq 4)$ is of type AC.

Proof. The proof follows the same lines as the proof for type A quadruples. By the definition of type AC we have

$$D(\beta_3, \beta_4) \subset u(E_{1,3} \cap D(L_3 L_2)).$$

We also have

$$u' E_{1,3} \subset D(\beta_1, \beta_2) \cap E_{1,3}$$

but

$$u E_{1,3} \cap E_{1,3} = \emptyset$$

This might seem strange because the first common unit disc crossing of β_3 and β_4 is on one side of $uE_{1,3}$, and the second common crossing is on the other side of $uE_{1,3}$. But the paths of $R_{m,0}$ with first two unit disc crossings in common with β_3 and β_4, and the third crossing strictly between this and the second and the first, are not in $E_{1,3}$. The paths in $uE_{1,3}$ are of this type. Related to this, provided v is nontrivial, or $u \neq va$ we have

$$L_3 L_2 R_3 E_{1,3} \cap E_{1,3} = \emptyset$$

and , further,

$$L_3 L_2 R_3 (\cup_{i \geq 0} u^i E_{1,3}) \cap (\cup_{i \geq 0} u^i E_{1,3}) = \emptyset$$

If v is trivial and $u = a$ then we still have

$$L_3 L_2 R_3 (E_{1,3} \cap D(BC)) \cap (E_{1,3} \cap D(BC)) = \emptyset$$

and

$$L_3 L_2 R_3 (\cup_{i \geq 1} u^i E_{1,3}) \cap (\cup_{i \geq 1} u^i E_{1,3}) = \emptyset.$$

In all cases, we have

$$D(\beta_1,\beta_2) \cap L_3L_2R_3(E_{1,3} \cup D(\beta_1,\beta_2)) = \emptyset.$$

In view of this we define

$$E = \cup_{i\geq 1} u^i E_{1,3} \cup D(\beta_1,\beta_2).$$

Then in all cases we have

$$E \cap L_3L_2R_3E = \emptyset$$

The only exchange $u_2 \leftrightarrow u_2'$ for β_3 such that $C(u_2 \leftrightarrow u_2', E)$ intersects $uE_{1,3} \cup D(\beta_1,\beta_2)$ without being contained in it is $u \leftrightarrow u'$. Similarly the only exchange $u_2 \leftrightarrow u_2'$ such that $C(u_2 \leftrightarrow u_2', E)$ intersects $\cup_{1\leq i\leq n} u^i E_{1,3} \cup D(\beta_1,\beta_2)$ without being contained in it is $u^{n+1} \leftrightarrow u^n u'$. It follows by induction on i, using 4.9.1, 4.8.2, 4.10.1 and 4.10.2 as in 4.12.1, that β_ℓ exists as in the statement of 2.8 for all $\ell \leq 2i$, for all i, with $I(\beta_{2j-1},\beta_{2j}) \subset E$ for all $j \leq i$. These properties imply that $w_1'(\beta_j)- = w_1'(\beta_3)$ for all $j \geq 3$ and hence the exchanges for β_j are closely related to the exchanges for β_3. In particular it is still true that $C(u_2 \leftrightarrow u_2', E)$ is contained in E whenever it intersects it, where $u_2 \leftrightarrow u_2'$ is an exchange for β_j, for any $j \geq 3$, since $\overline{\beta_j} * \beta_3 \subset E$. □

THEOREM 4.12.3. *Let $\beta \in \mathcal{Z}_m(3/7,+,+,0)$ and let $p(\beta)$ denote the endpoint of β. Let $(\beta_i(x) : 1 \leq i \leq 4)$ be of type C, with associated basic exchange $va \leftrightarrow vb$ prefixing the exchange $u \leftrightarrow u'$ for β_3. Then β_i exists as in Theorem 2.8 with*

$$\psi_{1,3}(I(\zeta_{2i-1},\zeta_{2i}(x)) \subset D(\beta_1,\beta_2) \tag{4.12.3.1}$$

for all $i \geq 2$ and

$$D(\beta_{2i-1},\beta_{2i}) \subset (D(\beta_1,\beta_2) \cap E_{1,3}) \cup_{n\geq 1} (u)^n E_{\beta_1,\beta_3}. \tag{4.12.3.2}$$

for all $i \geq 3$.

PROOF. By hypothesis, $(\beta_i : 1 \leq i \leq 4)$ is of type C. We can write $E_{1,3} = E_1 \cup E_2$ where

$$E_1 = \{\zeta \in E_{1,3} : w_1'(\zeta_3) < w_1'(\zeta) <\leq w_1'(\zeta_1)\}$$

$$E_2 = \{\zeta \in E : w_1'(\zeta) = w_1'(\zeta_3)\}$$

Then the conditions of type C are such that

$$uE_{1,3} \cap E_{1,3} = uE_1 \cap E_2.$$

It is also the case that $u'D(L_3L_2) \cap E_{1,3} = \emptyset$ – which was not the case in 4.12.1. As before we have

$$E_{1,3} \cap L_3L_2R_3E_{1,3} = \emptyset$$

and the only exchanges $u_2 \leftrightarrow u_2'$ for which $C(u_2 \leftrightarrow u_2')$ intersects $E_{1,3}$ without being contained in it are $u \leftrightarrow u'$ and $uu \leftrightarrow uu'$. in fact, we now have

$$uuE_{1,3} \cup uu'E_{1,3}) \cap E_{1,3} = \emptyset,$$

but $C(uu \leftrightarrow uu', E_{1,3}$, which is the support of a disc exchange in the composition for $\psi_{1,3}$, has transversal intersection with $I(\beta_3, \beta_4)$. It follows that (β_5, β_6) exists and either $I(\beta_5, \beta_6) \subset D(\beta_3, \beta_4) \cap uE_{1,3}$ or $I(\beta_5, \beta_6) \subset u^2E_{1,3}$. In both cases (β_7, β_8) exists with all the required properties and $I(\beta_7, \beta_8) \subset D(\beta_3, \beta_4) \cap uE_{1,3}$ or $I(\beta_7, \beta_8) \subset u^2E_{1,3}$ respectively. Then similarly to 4.12.2, we define

$$E = D(\beta_1, \beta_2) \cup \cup_{n \geq 1} u^i E_{1,3}$$

Once again if $u_2 \leftrightarrow u_2'$ is any exchange for β_3, then $C(u_2 \leftrightarrow u_2')$ is contained in E whenever it intersects it. So by induction $\beta_\ell$exists as i Theorem 2.8 for $\ell \leq 2i$ for all $i \geq 1$, with $I(\beta_{2i-1}, \beta_{2i}) \subset E$, and $C(u_2 \leftrightarrow u_2')$ is contained in E whenever it intersects it for any exchange $u_2 \leftrightarrow$ for β_j, for any $j \geq 3$.

□

CHAPTER 5

Specific examples

In this chapter we expand on the examples in [**23**]. We show, somewhat surprisingly, that the Thurston equivalence classes found in [**23**] are considerably larger than was demonstrated there. We will use this class of examples in Theorem 6.2.

5.1. Some old and new notation

For any word x which ends in C, we write $p(x)$ for the point in $Z \cap D(x)$ of lowest possible preperiod under s.

It is convenient to modify and extend the notation of [**23**]. As in [**23**] we define

$$a = L_3(L_2R_3)^2, \quad b = L_3^5,$$

$$c = L_3^3L_2C,$$

$$d = L_3^2a,$$

$$v_0 = L_3L_2R_3a, \quad w_0 = L_3L_2R_3b,$$

$$u_0 = L_3L_2R_3L_3L_2C,$$

$$t_0 = L_3L_2R_3d,$$

and inductively, for $k \geq 0$, we define

$$v_{k+1} = v_kt_ka, \quad t_{k+1} = v_kt_kd, \quad u_{k+1} = v_kt_kc.$$

Now we generalise this construction. Let $\alpha = (\alpha(k)) \in \{a, \underline{\}}^{\mathbb{N}}$. We define words $v_{k,\alpha}$, $t_{k,\alpha}$ and $u_{k,\alpha}$ for all $k \geq 0$. To start the induction we define

$$v_{0,\alpha} = \begin{cases} v_0 \text{ if } \alpha(0) = a, \\ w_0 \text{ if } \alpha(0) = b, \end{cases} \qquad t_{0,\alpha} = t_0.$$

Then we define, for $k \geq 0$,

$$v_{k+1,\alpha} = v_{k,\alpha}t_{k,\alpha}\alpha(k), \quad t_{k+1,\alpha} = v_{k,\alpha}t_{k,\alpha}d, \quad u_{k+1,\alpha} = v_{k,\alpha}t_{k,\alpha}c.$$

Thus, the sequences v_k, t_k and u_k are $v_{k,\alpha}$, $t_{k,\alpha}$, and $u_{k,\alpha}$ in the special case where $\alpha = \alpha^\infty$, where $\alpha^\infty(k) = a$ for all a. We also define α^ℓ for $\ell \geq 0$ to be the sequence such that

$$\alpha^\ell(k)=\begin{cases}a \text{ if } k<\ell,\\ b \text{ if } \ell\geq k.\end{cases}$$

Then the word which was called $w_{\ell,r}$ in [**23**] for $0\leq\ell\leq r$ is v_{r,α^ℓ} in our current notation. It will sometimes be convenient to write

$$v_{k,\alpha^\ell}=v_{k,\ell},\quad t_{k,\alpha^\ell}=t_{k,\ell},\quad u_{k,\alpha^\ell}=u_{k,\ell}$$

which is the opposite order of indices to that used in [**23**], but since different letters are used, hopefully this will not cause confusion We define, for $r\geq 0$

$$x_r=v_ru_r$$

and, more generally,

$$x_{r,\alpha}=v_{r,\alpha}u_{r,\alpha},\quad x_{r,\alpha^\ell}=x_{r,\ell}.$$

We define $\gamma^{r,\alpha}$ to be the path in $\mathcal{Z}(a_1,+,+,0)$ with endpoint $p(x_{r,\alpha})$ and

$$\gamma^{r,\ell}=\gamma^{r,\alpha^\ell},\quad 0\leq\ell\leq r+1.$$

Since $x_{r,\alpha}$ depends on $\alpha(\ell)$ only for $\ell\leq r$, there are only finitely many choices for $x_{r,\alpha}$: in fact, 2^{r+1} choices. In [**23**], it was proved that the captures $\sigma_{\gamma^{r,\ell}}\circ s$, for $0\leq\ell\leq r+1$ are all Thurston equivalent. (The notation here has been changed from that in [**23**], as $\beta$with various indices is rather heavily used in this paper). In [**23**] the proof was direct. We shall now give a more indirect proof in keeping with the methods of this paper. Note that the preperiod of the points $x_{r,\alpha}$ is the same for all α. We fix m so that $p(x_{r,\alpha})$ is of pre-period $\leq m$ under s. Write $n_{r,\ell}=n(\gamma^{r,\ell})$, and $n(r,\alpha)=n(\gamma^{r,\alpha})$. We will show the following.

THEOREM 5.2. *Fix $r\geq 0$. Let α_1 and $\alpha_2\in\{a,b\}^{r+1}$ be such that, for some $p<r$,*

$$\alpha_1(i)=\alpha_2(i) \text{ for } i\leq p$$
$$\alpha_1(i)a,\alpha_2(i)=b \text{ for all } p<i.$$

Then for $\gamma_i=\gamma^{r,\alpha_i}$, the captures $\sigma_{\gamma_1}\circ s$ and $\sigma_{\gamma_2}\circ s$ are Thurston equivalent. Hence the captures $\sigma_{\gamma^{r,\alpha}}\circ s$ are Thurston equivalent for all $\alpha\in\{a,b\}^{r+1}$.

The final claim of Theorem 5.2 follows easily from the first part, because we see inductively that $\sigma_{\gamma^{r,\alpha}}$ is Thurston equivalent to $\sigma_{\gamma^{r,\infty}}$ for all $\alpha\in\{a,b\}^{\mathbb{N}}$, by moving from α^∞ to α in a series of steps, α_i for $0\leq i\leq n$. To do this, we define $\alpha_0=\alpha^\infty$, and let p_i be the increasing sequence of non-negative integers such that $\alpha(j)$ is constant for $p_i<j\leq p_{i+1}$ and $\alpha(i)=a$ for $i<p_1$. Then we define α_i by induction on i by

$$\alpha_{i+1}(j)=\alpha_i(j) \text{ if } j\leq p_i,$$

$$\alpha_{i+1}(j) = b \neq a = \alpha_i(j) \text{ if } p_i < j \text{ and if } i \text{ is even,}$$

$$\alpha_{i+1}(j) = a \neq b = \alpha_i(j) \text{ if } p_i < j \text{ and if } i \text{ is odd.}$$

Then $\alpha_n = \alpha$. We shall use Theorem 5.2 in Chapter 6 to show that the set of capture paths γ in $\mathcal{Z}_m$ such that $\sigma_\gamma \circ s$ is equivalent to $\sigma_{\gamma_2} \circ s$ for precisely 2^{r+1} capture paths γ_2 in $\mathcal{Z}_m$, is of positive density, as $m \to \infty$, for each $r \geq 0$ (in fact, also for $r = -1$, but that case does not use this construction). The idea of that proof will be to show that, for $\gamma = \gamma^{r,\alpha}$, for each $\alpha \in \{a,b\}^{\mathbb{N}}$, the density of paths $\omega \in R_{m,0}$ in a sufficiently small neighbourhood U of β such that $\sigma_\omega \circ s$ is equivalent to $\sigma_{\gamma_2} \circ s$ for exactly one path γ_2 in a given neighbourhood U' of γ, approaches 1 as U is made arbitrarily small, and that any capture path $\gamma_3 \in \mathcal{Z}$ such that $\sigma_\omega \circ s$ is equivalent to $\sigma_{\gamma_3} \in \mathcal{Z}$, has to be in a given neighbourhood U'_ℓ of $\gamma^{r,\ell}$ or $U'_{\ell,-}$ of $\gamma^{r,\ell,-}$, for U sufficiently small.

We now prove 5.2. This proof follows the lines of the proof of the corresponding result in [**23**] pretty exactly, with some minor differences of notation, to accommodate the notation of this paper. I had actually hoped to provide a proof which gave an explicit calculation of the path $\beta \in R_{m,0}$ such that $\sigma_\beta \circ s$ is equivalent to $\sigma_\gamma \circ s$ for $\gamma = \gamma^{r,\alpha}$, for all α. But the calculation is simply too complicated.

PROOF. Write $x = x_{r,\alpha_2}$ and $y = x_{r,\alpha_1}$. Exactly as in [**23**], $D(y)$ is to the right of $D(x)$. Now we imitate the proof of Lemma 2.6 of [**23**]. We claim that the suffixes u of y such that $D(u)$ is between $D(x)$ and $D(y)$ are precisely those which start with $v_{\alpha_1,p+1}$ or $t_{\alpha_1,p+1} = v_{\alpha_2,p+1}$ or $u_{\alpha_1,p+1} = u_{\alpha_2,p+1}$. We see this as follows. The largest common prefix of x and y is $v_{\alpha_1,p}t_{\alpha_1,p}L_3 = v_{\alpha_2,p}t_{\alpha_2,p}L_3$. So this must also be a prefix of u, if $D(u)$ is between $D(x)$ and $D(y)$. Since u is also a suffix of y, it must start with either $v_{\alpha_1,p+1}$ or $t_{\alpha_1,p+1}$ or $u_{\alpha_1,p+1}$. It remains to show that this is a sufficient condition for $D(u)$ to be between $D(x)$ and $D(y)$. If the prefix of u is $t_{\alpha_1,p+1}$ or $u_{\alpha_1,p+1}$, then this is clear. So now suppose that the prefix of u is $v_{\alpha_1,p+1}$. It is clear that $D(u)$ is to the right of $D(x)$. So we need to show that $D(u)$ is to the left of $D(y)$. For some k with $p+1 < k < r$, it must be the case that t_{k,α_1} is a prefix of u. Since $D(t_{k,\alpha_1})$ is to the left of $D(v_{k,\alpha_1})$, which contains $D(x)$, the claim is proved.

Now the proof continues exactly as in [**23**]. We choose z with $p(z) \in Z_\infty$ such that all the letters of z apart from the last letter C are in $\{L_3, L_2, R_3\}$, such that $D(z)$ is to the right of $D(y)$, and such that the forward orbit of $p(z)$ does not intersect the subset D' of the unit disc between $D(x)$ and $D(y)$, that is

$$p(z) \notin \cup_{n\geq 0} s^{-n}(D(x) \cup D' \cup D(y)).$$



FIGURE 1. ω

We let $\gamma(z)$ denote the capture path in $\mathcal{Z}(3/7,+,+,0)$ which crosses the upper unit circle into $D(z)$. Now we define paths η_1 from $p(z)$ to $p(y)$, and a closed path ω based at $p(z)$. The path η_1 first crosses the unit circle out of the gap of $L_{3/7}$ containing $p(z)$, and then crosses the second time into the gap containing $p(y)$. The closed loop ω also crosses the upper unit circle out of the gap containing $p(z)$. It then crosses the upper unit circle again at the top edge of the right-most leaf of $L_{3/7}$ in the boundary of the gap containing $p(x)$. It traces an anticlockwise path round the boundary of the subset of the unit disc bounded by $D(x)$ on the left and $D(y)$ on the right, as far as the top of the left-most leaf in the boundary of the gap containing $p(y)$. It then re-crosses the upper unit circle into the gap of $L_{3/7}$ containing $p(z)$, and ends at $p(z)$.

It is then immediate that

$$\sigma_{\gamma_1} \circ s \simeq \sigma_{\eta_1} \circ \sigma_{\gamma_z} \circ s.$$

It should be noted that the notation here is different from that in [**23**], where the paths which are here called η_1 and ω were called γ_1 and α. For most of this paper, paths β, with various indexes, denote paths in $R_{m,0}$, and, from now on, paths γ, with various indexes, denote capture paths, usually in $\mathcal{Z}(3/7,+,+,0)$. In particular, at present, the paths γ_1 and γ_2 denote the capture paths γ^{r,α_1} and γ^{r,α_2}.

Since the disc enclosed by ω is disjoint from the forward orbit of z, we have

$$(\sigma_{\gamma(z)} \circ s, Y_m) \simeq_{\psi_m} (\sigma_\omega \circ \sigma_{\gamma(z)} \circ s, Y_m)$$

for some homeomorphism ψ_m, for any m such that $p(x)$ and $p(y) \in Z_m$. Hence, if

$$\eta_2 = \omega * \psi_m(\eta_1),$$

we have

$$(\sigma_{\eta_1} \circ \sigma_{\gamma(z)} \circ s, Y_m) \simeq_{\psi_m} (\sigma_{\eta_2} \circ \sigma_{\gamma(z)} \circ s, Y_m).$$

The aim then, as in [**23**], is to compute ψ_m and to show that

$$(\gamma(z) * \eta_2, O^+(p(x))) \simeq (\gamma_2, O^+(p(x))), \tag{5.2.0.3}$$

where, here, $\simeq$ denotes homotopy constant on $O^+(p(x))$, the forward orbit of $p(x)$. We then have

$$\sigma_{\eta_2} \circ \sigma_{\gamma(z)} \circ s \simeq \sigma_{\gamma_2} \circ s$$

and hence

$$\sigma_{\gamma_2} \circ s \simeq \sigma_{\gamma_1} \circ s,$$

as required. As usual, ψ_m is a composition of disc exchanges with supports $(\sigma_{\gamma(z)} \circ s)^{-n}(D')$ for varying $n > 0$. We only need to consider those disc exchanges which have an effect on $p(y)$. These are the disc exchanges with supports $C(v'a \leftrightarrow v'b, D')$ where $v'b$ is a prefix of x_{r,α_2} of the same length as a prefix v of x_{r,α_1} such that $vau = x_{r,\alpha_1}$. It follows that $\psi_m(\eta_1)$ has endpoint $p(x)$, as required, because the successive changes from a to b are made preceding suffixes u of x_{r,α_1} with $D(u) \subset D'$, that is, precisely the changes that are needed to change $y = x_{r,\alpha_1}$ to $x = x_{r,\alpha_2}$. These disc exchanges also introduce hooks of $\psi_m(\eta_1)$ round the sets $D(v'a)$. But these do not matter, because such a $v'a$ never occurs as a subword of x, that is, none of these sets intersects the forward orbit $O^+(p(x))$ of $p(x)$. We see this as foilows. Each $v'b$ is a prefix of x_{r,α_2} which ends in x_{r,α_2} before a subword v_{p+1,α_2} or t_{p+1,α_2}. So $v'b$ must end in a subword v_{k,α_2} for some $k \geq p+1$. It follows that the only occurrences of v' in x_{r,α_2} must be followed by b, since $\alpha_2(i) = b$ for all $i \geq p+1$. So (5.2.0.3) follows, as required.

□

For the remainder of this section, we obtain information as possible about the paths $\beta_i(\gamma^{r,\alpha})$, in preparation for studying perturbations of the capture paths $\gamma^{r,\alpha}$ in Chapter 6.. We denote by $n(\alpha)$ the largest integer such that $(\beta_{2n(\alpha)}(\gamma^{r,\alpha})$ is defined. Even though, by Theorem 5.2,. we have

$$\beta_{2n(\alpha_1)-1}(\gamma^{r,\alpha_1}) = \beta_{2n(\alpha_2)-1}(\gamma^{r,\alpha_2}),$$

for all $\alpha_1, \alpha_2 \in \{a,b\}^{r+1}$, it does not follow, and is certainly not true, that $\beta_i(\gamma^{r,\alpha_1}) = \beta_i(\gamma^{r,\alpha_2})$ for all i.

5.3. Lemmas about the ζ sequence for $x_{r,\alpha}$

LEMMA 5.3.1. 1.

$$w_1'(v_0u_0) = L_3, \quad w_1'(w_0u_0) = L_3L_2R_3L_3^3,$$

$$w_2'(v_0u_0) = v_0L_3L_2R_3L_3^3, \quad w_2'(w_0u_0) = w_0L_3L_2R_3L_3^3.$$

More generally, if $\alpha(0) = a$ then, for all $r > 0$,

$$w_1'(x_{r,\alpha}) = L_3, \quad w_2'(x_{r,\alpha}) = v_0t_0aL_3,$$

and if $\alpha(0) = b$ *then for all* $r > 0$,

$$w'_1(x_{r,\alpha}) = L_3L_2R_3L_3^3, \quad w'_2(x_{r,\alpha}) = w_0L_3L_2R_3L_3^3L_2y$$

for some y.

2. *For any* $r > 0$ *and any* α, *and any prefixes* u_1 *and* u_1u_2 *of* $x_{r,\alpha}$ *such that each of* u_1 *and* u_2 *ends in* a *or* b *(or* u_1 *can be empty), the number of* j *such that*

$$|u_1| < |w'_j(x_{r,\alpha})| \leq |u_1| + |u_2|$$

is equal to the number of occurrences of da *or* db *or* dd *in* u_2. *If* u_2 *ends in* c, *that is, if* $x_{r,\alpha} = u_1c$ *then we add two for the last occurrence of* dc.

3. *In particular, for any* $\ell > 0$, *the number of* j *for which* $|w'_j(x_{r,\alpha})| \leq |v_\ell|+1$ *is* 2^ℓ *for all* $0 < \ell \leq r$, *the number for which* $|w'_j(x_{r,\alpha})| \leq |v_\ell|$ *is* $2^\ell - 1$ *if* $\alpha(\ell) = a$ *and* 2^ℓ *if* $\alpha(\ell) = b$, *and the number of* j *for which* $w'_j(x_{r,\alpha})$ *is defined is* 2^{r+1}.

PROOF. 1. This is a direct calculation.

2. Apart from the prefix v_0 or w_0 of $x_{r,\alpha}$ (depending on whether $\alpha(0) = a$ or b) every other occurrence of v_0 (or w_0) occurs in a subword dav_0t_0 or dbv_0t_0 (or daw_0t_0 or bw_0t_0). Also $x_{r,\alpha}$ starts with v_0t_0 or w_0t_0, and then is made up of subwords u with some overlapping:

$$dd, \quad dav_0t_0, \quad dbv_0t_0, \quad daw_0t_0, \quad dbw_0t_0, \quad dc.$$

For each such occurrence u, if z is the prefix of $x_{r,\alpha}$ which precedes the occurrence of u in $x_{r,\alpha}$, then, apart from dc, by the rules described in 3.4 to 3.7, the corresponding word $w'_j(x_{r,\alpha})$ is:

$$zdL_2R_3L_3, \quad zdaL_3, \quad zdL_2R_3L_3, \quad zdaL_3L_2R_3L_3^3, \quad zdbL_3L_2R_3L_3^3.$$

As for dc, if $x_{r,\alpha} = zdc$ then the two words $w'_j(x_{r,\alpha})$ corresponding to dc are:

$$zdL_2R_3L_3, \quad zdL_3^5.$$

So, as claimed, we simply need to count occurrences of dd, da and db, and two for the last occurrence of dc. Since the occurrences of db in $x_{r,\alpha}$ are simply obtained by replacing occurrences of da in x_r, it suffices to count occurrences of da and dd in x_r.

3. A simple calculation shows that the number $N(\ell)$ of occurrences of dd or da or db in t_k is the same as the number in v_k, for all $k \geq 1$, and $N(1) = 1$. Also, $N(\ell+1) = 2N(\ell)+1$. So, by induction, $N(\ell) = 2^\ell - 1$, and the number of j for which $|w'_j(x_r)| \leq |v_\ell|$ is $2^\ell - 1$, adding in $w'_1(x_r) = L_3$, but discounting $w'_{2^\ell}(x_r) = v_\ell L_3$, since this corresponds to the occurrence of da at the end of v_ℓ. The number of j for which $|w'_j(x_r)| \leq |v_\ell|+1$ is 2^ℓ. The

number of occurrences of dd or dc in u_r is the same as the number of da or dd in v_r. So the number of words $w'_j(x_r)$ is $2\cdot(2^r-1)+1=2^{r+1}-1$.

Now we consider $x_{r,\alpha}$ in general. The number of occurrences of dd, da and db in $v_{\ell,\alpha}$ is the same as the number of occurrences of dd or da in v_ℓ, and the number of occurrences of dd, da, db or dc in $x_{r,\alpha}$ is the same as the number of dd, da or dc in x_r. The only difference is that $w'_{2^\ell}(x_{r,\alpha})| < |v_{\ell,\alpha}|$ if and only if $\alpha(\ell)=b$. So the number of j such that $|w'_j(x_{r,\alpha})| < |v_\ell|$ is the same as the number with $|w'_j(x_{r,\alpha})| \le |v_\ell|+1$ if $\alpha(\ell)=b$.

□

Now we consider the sequence $\zeta_i(x)$ of 4.4 for $x=x_{r,\alpha}$. From 5.3 and the definitions in 4.5 and 4.6, it follows that if $\alpha(0)=a$, then $\zeta_i(x)$ is defined for all i with $2i\le n(x)$ and $(\zeta_i(x_{r,\alpha}):1\le i\le 4)$ is of type A. If $\alpha(1)=b$ then $\zeta_i(x)$ is only defined for $i=1$, 2, but $w'_2(x)=w_1(x)u$ where $L_3u=t_0u'$ for some u and u', and since $v_0<t_0<w_0$, the conditions of 4.7.3 are satisfied for the existence of a type C quadruple $(\zeta_j:1\le i\le 4)$ such that $\zeta_j(x)=\zeta_j$ for $j=1$, 2, and $\zeta_j=\zeta_j(x_{\alpha^\ell})$ for $j=3$, 4, for any $\ell\ge 1$. For most of the time, we will consider $\zeta_i(x_{r,\alpha})$ for α with $\alpha(0)=a$. We will see later that, if $\alpha(0)=b$, then the sequence to consider is $\zeta_i(x_{r,\alpha'})$, where $\alpha'(0)=a$ and $\alpha'(\ell)=\alpha(\ell)$ for all $\ell>0$.

For the moment, suppose that $x=x_{r,\alpha}$ with $\alpha(0)=a$. It is reasonable to expect that all but the last few disc crossings by $\zeta_{2i-1}(x)$ and $\zeta_{2i}(x)$ are in the sequence of crossings determined by $w'_j(x)$. Recall that $k=k(\zeta_{2i-1},\zeta_{2i})$ is the first integer for which the k'th unit-disc-crossings of ζ_{2i-1} and ζ_{2i} do not coincide. For the moment, we write $k(i,x)$ for this integer, so that $w'_j(\zeta_{2i-1}(x))=w'_j(\zeta_{2i}(x))$ for $j<k(i,x)$. The integer $k(i,x)$ is always odd, and $k(i,x)\le k(i+1,x)\le k(i,x)+2$ for all i.

LEMMA 5.3.2. *Let $x=x_{r,\alpha}$ for any $r\ge 0$ and any α with $\alpha(0)=a$. We write $k(i,x)=k(\zeta_{2i-1}(x),\zeta_{2i}(x))$.*

1. *Apart from (ζ_1,ζ_2), there are two pairs $(\zeta_{4i-1}(x),\zeta_{4i}(x))$ and $(\zeta_{4i+1}(x),\zeta_{4i+2}(x))$, with i even with*

$$w'_{k-1}(\zeta_{2i+s}(x))=w'_{k-1}(x),\quad -1\le s\le 2$$

for each odd $k\ge 5$. There are two extra pairs for each prefix $zv_0L_3L_2R_3L_3^3$ of $x_{r,\alpha}$ such that z has an even number of letters L_3 and L_2.

2. *$(\zeta_{4i+t}(x):-3\le t\le 0)$ is of type A if $w'_{k(2i-1,x)-1}(\zeta_{2i-1}(x))=w'_{k(2i-1,x)-1}(x)$ and of type B otherwise, in which case*

$$w'_{k(2i-1,x)-1}(\zeta_{2i-1}(x))=zv_0L_3L_2R_3L_3^3$$

is a prefix of x_r *such that* z *has an even number of letters* L_3 *and* L_2.

3. *For any* α, *and any* $r \geq 1$, *the pair* $(\zeta_{2i-1}(x), \zeta_{2i}(x))$ *is defined for*

$$i \leq 2^{r+1} + 2^{r-1} - 1.$$

Moreover, we have

$$|w'_{k(i,x)-1}(\zeta_{2i-1}(x))| < |v_1|$$

for $i \leq 3$, *and for any* $2 \leq \ell \leq r$, *we have*

$$|w'_{k(i,x)-1}(\zeta_{2i-1}(x))| < |v_\ell|$$

for $i \leq 2^\ell + 2^{\ell-1} - 1$ *if* $\alpha(\ell) = a$, *and for* $i \leq 2^\ell + 2^{\ell-1}$ *if* $\alpha(\ell) = b$. *If* $\alpha(\ell) = a$ *then for the next pair* $(\zeta_{2i-1}(x), \zeta_{2i}(x))$, *we have*

$$|w'_{k(i,x)-1}(\zeta_{2i-1}(x))| = |w'_{k-1}(\zeta_{2i}(x)| = |v_\ell| + 1.$$

4. *For* $1 \leq \ell \leq r$, *we have*

$$|w'_{k(i,x)-1}(\zeta_{2i-1}(x))| \leq |v_\ell| + 1 \quad \Leftrightarrow \quad k(i,x) \leq 2^\ell,$$

and

$$|w'_{k(i,x)-1}(\zeta_{2i-1}(x))| = 2^{r+1} - 1 \quad \Leftrightarrow \quad k(i,x) = 2^{r+1} - 1.$$

PROOF. First we consider the case of $x = x_r$. We write $\zeta_i(v_r u_r) = \zeta_i$ and $k_i = k(i, v_r u_r)$. Recall that, apart from the occurrence at the beginning of the word, each occurrence of $v_0 t_0$ in $v_r u_r$ is preceded by $da = L_3^2 a^2$. Now we consider the possible words $w'_j(v)$ for any word v for prefixes v of $v_r u_r$ ending in $L_3 L_2$, for any even value of j. These are the possible words for $w'_{k_i-1}(\zeta_{2i-1}) = w'_{k_i-1}(\zeta_{2i})$, for any i. Of course the words $w'_j(x)$ are possibilities, for even $j < 2^{r+1}$. We have seen in 5.3.1 that each word $w'_j(x_r)$ — without the restriction to even j — is determined by an occurrence of da or of dd, apart from $j = 1$ (which corresponds to the first occurrence of v_0) and the three largest possible values of j. More precisely, if $z_1 = zdd$ is a prefix of $v_r u_r$, then the largest j with $|w'_j(v_r u_r)| < |z_1|$ gives $w'_j(v_r u_r) = zdL_2 R_3 L_3$. If $z_2 = zdav_0 t_0$, then the largest j with $|w'_j(v_r u_r)| < |z_2|$ gives $w'_j(v_r u_r) = zdaL_3$. In both cases, if $z = z'd$, then $zL_2 R_3 L_3 = w'_{j-1}(v_r u_r)$, but this is associated to the previous occurrence of dd in $v_r u_r$. The other alternative is that $z = z' v_0 L_3 L_2 R_3$, and in this case $w'_{j-1}(v_r u_r)| < |z|$. By induction, there are $2^r - 1$ occurrences of da in x_r, and $2^r - 2$ occurrences of dd. Of these, $2^{\ell-1}$ and $2^{\ell-1} - 1$ respectively occur in the prefix v_ℓ of x_r if $\ell \geq 1$. We have seen that $j = 2^\ell$ for the occurrence of $dav_0 t_0$ preceded in $v_r u_r$ by z, for which the $zda = v_\ell$ for any $\ell \geq 2$. In particular, this j is even. It follows that if $zv_0 t_0$ is a prefix of v_ℓ, then the integers j_1 and j_2 such that $w'_{j_1}(v_r u_r) = zL_3$ and $w'_{j_2}(v_r u_r) = v_\ell z L_3$ have opposite parities. Hence, by induction, for each $r\geq\ell\geq 2$, the number of prefixes zda of v_ℓ with z nontrivial and for which $w'_j(zdav_0t_0)=zdaL_3$ and for which j is even, is the same as the number of such prefixes with j odd. So both these numbers are $2^{\ell-2}$. Similarly, the numbers for such prefixes zv_0 of v_ru_r, are 2^{r-1} for both j even and j odd, for all $r\geq 1$.

We have $k(1,x)=1$ and $k(i,x)=3$ for $2\leq i\leq 5$. We have

$$w'_2(\zeta_j)=v_0L_3L_2R_3L_3^3 \text{ for } 3\leq j\leq 6,$$

and

$$w'_2(\zeta_j)=v_0t_0aL_3 \text{ for } 7\leq j\leq 10.$$

The reason for the extra pairs is that

$$w'_3(v_0L_3L_2R_3L_3L_2BC)=w'_3(v_0L_3L_2R_3L_3^6L_2)=v_2L_3L_2R_3L_3^3,$$

and $v_0L_3L_3R_3L_3L_2BCL_1R_2R_3$ and $v_0L_3L_2R_3L_3^6L_2$ are prefixes of $w'_3(\zeta_3)$ and $w'_3(\zeta_4)$ respectively. It is therefore clear that $I(\zeta_3,\zeta_4)$ is over $D(x_r)$. Similarly we have extra pairs for each prefix $zv_0L_3L_2R_3L_3^3$ of x_r such that z has an even number of letters L_3 and L_2. We do not have any other extra pairs, because although there are other prefixes $z_1L_3^3$ of x_r, these are prefixes of the form $z_2dL_3^3=z_2L_3^3(L_2R_3)^2L_3^3$ rather than $z_2v_0L_3L_2R_3L_3^3$. It follows from the detail of the rules in 3.4 to 3.7 that there are no extra pairs in this case.

So the pairs $(\zeta_{2i-1}(x),\zeta_{2i}(x)$ correspond two-to-one with $w'_k(x_r)$ for odd values of $k\geq 5$ and with $k\leq 2^{r+1}-1$, and two extra pairs for each prefix $zv_0L_3L_2R_3L_3^3$ of x_r such that z has an even number of letters L_3 and L_2. We then have

$$w'_{k-2}(zv_0L_3L_2R_3L_3^3L_2)=zL_3=w'_{k-2}(x_r),$$

but $w'_{k-1}(x_r)\neq w'_{k-1}(zv_0L_3L_2R_3L_3^3L_2)$. If k is odd, and $k-1$ is even, there are pairs $(\zeta_{4i-1},\zeta_{4i})$ and $(\zeta_{4i+1},\zeta_{4i+2})$, for some i, such that $k(2i,x)=k(2i+1,x)$ and $w'_{k-1}(\zeta_{4i+s})=zL_3^3$ for $-1\leq s\leq 2$, and $w'_q(\zeta_{2i+s})=w'_q(v_ru_r)$ for $q\leq k-2$. The exceptional feature of these pairs is that $(\zeta_{4i+s}:1\leq s\leq 4)$ is of type B, with

$$k=k(\zeta_{4i+s-1},\zeta_{4i+s})=k(\zeta_{4i-1},\zeta_{4i})$$

for $0\leq s\leq 3$, and

$$w'_{k-1}(\zeta_{4i+t})\neq w'_{k-1}(\zeta_{4i+u})=w'_{k-1}(x),\quad -1\leq t\leq 2,\quad 3\leq y\leq 6.$$

So there are four pairs corresponding to each such value of $w'_{k-1}(x)$ with $k-1$ even. These are pairs $(\zeta_{4i+2s-1}(x),\zeta_{4i+2s}(x))$ for $0\leq s\leq 3$ for some i.

The number of occurrences of v_0t_0 in $x_r=v_ru_r$ is 2^{r+1} and the number of occurrences in $v_\ell=v_{\ell-1}t_{\ell-1}a$ is 2^ℓ, for all $\ell\geq 1$. For all $r\geq 1$, if uv_0t_0

is a prefix of v_r, then $v_ruv_0t_0$ is a prefix of x_r, and vice versa. Moreover, exactly one of v_r and u_r has an even number of letters L_3 and L_2, and hence the number of prefixes zv_0t_0 of x_r for which z has an even number of letters L_3 and L_2 is 2^r. Similarly for all $\ell \geq 2$ — which we need so that v_0t_0 is a prefix of $v_{\ell-1}$ and $t_{\ell-1}$ — the number of prefixes zv_0t_0 of v_ℓ for which z has an even number of letters L_3 and L_2 is $2^{\ell-1}$.

It follows that the number of pairs $(\zeta_{2i-1}(x), \zeta_{2i}(x))$ is

$$1 + 4 \cdot 2^r = 1 + 2^{r+1} + 2^r,$$

and the number for which $|w'_{k-1}(\zeta_{2i-1})| < |v_1|$ is 3, and, for $\ell \geq 2$, the number for which $|w'_{k-1}(\zeta_{2i-1})| < |v_\ell|$ is 1 — plus twice times the number of odd k (even $k-1$) such that $|w'_{k-1}(x)| < |v_\ell|$, which is $2^\ell - 2$ – plus twice for the even number $k-1$ such that $|w'_{k-1}(x)| = |v_\ell| + 1$ – which is $2^{\ell-1} - 1$ — which gives $2^{\ell+1} - 1$, as required. The final pair for which $k(i,x) \leq 2^\ell + 1$ is $i = 2^\ell + 2^{\ell-2}$.

$$i = 2^\ell + 2^{\ell-1}.$$

The possible values of $k(i,x)$ are precisely the odd k for which $w'_k(x_{r,\alpha})$ is defined. Thus, 4 for x_r follows from 5.3 for x_r, and 1 to 4 are proved for x_r. The proof for $x_{r,\alpha}$ is exactly the same, apart from minor differences because some occurrences of dav_0t_0 are replaced by dbv_0t_0.

□

Let $x = x_{r,\alpha}$ for any $r \geq 0$ and any α and let $\gamma(x)$ be the associated capture path in $\mathcal{Z}_m(3/7, +, +, 0)$ and write $\beta_i(x)$ for the associated sequence $\beta_i(\gamma(x))$ as in 3.3. We write $E_{4j-3,4j-1,x}$ for the sequence $E_{4j-3,4j-1,\beta(x)} = E_{\beta_{4j-3}(x),\beta_{4j-1}(x),\beta(x)}$ as in 4.12, following the earlier notation of 3.8 and 4.5. If $\alpha(0) = a$, we also write

$$E'_{4i-3,4i-1,\alpha} = E'_{4i-3,4i-1,x} = E_{\zeta_{4i-3}(x),\zeta_{4i-1}(x)}$$

Since $\zeta_i(x)$ depends only on $\alpha(j)$ for $j \leq t$, we see that $E'_{4i-3,4i-1,x} = E'_{4i-3,4i-1,\alpha}$ depends only on $\alpha(j)$ for $j \leq t$, so long as $|w'_{k(2i,x)-1}(x)| \leq 2 \cdot |v_t| - 7$. In particular, $E'_{1,3,x}$ is depends only on $\alpha(0)$. We also define

$$\psi'_{4i-3,4i-1,x} = \psi_{\zeta_{4i-3}(x),\zeta_{4i-1}(x)}.$$

We know from 4.8.2 that $\zeta_i(x) = \beta_i(x)$ for $i \leq 4$ if $\alpha(0) = a$. Thus, $E_{1,3,x} = E'_{1,3,x}$ if $\alpha(0) = a$. We also know that $(\beta_t(x)) : 1 \leq t \leq 4)$ is of type C if $\alpha(0) = b$, with $\beta_t(x) = \zeta_t(x)$ for $t \leq 2$ and $\beta_t(x) = \zeta(x_{\alpha_1,r})$ for $t = 3$, 4,where $\alpha_1(0) = a$ and $\alpha_1(k) = \alpha(k)$ for $k \geq 1$.

Lemma 5.3.3. *Let $x = x_{\alpha,r}$ for any α with $\alpha(0) = a$, and any $r \geq 1$. For all i,*

(5.3.3.1) $$k(2i, x) = k(2i+1, x) \leq k(2i+2, x),$$

$$\zeta_{4i}(x) < \zeta_{4i+s}(x) \text{ for all } s \geq 1. \tag{5.3.3.2}$$

$$k(2i+2,x) > k(2i+1,x) \Rightarrow k(2i+2,x) = k(2i,x)+2 \text{ for all } s \geq 4 \tag{5.3.3.3}$$

$$k(2i+2,x) = k(2i+1,x) \Leftrightarrow \zeta_{4i+2}(x) < \zeta_{4i+s}(x) \text{ for all } s \geq 3. \tag{5.3.3.4}$$

$$k(2i+2,x) > k(2i,x) \Leftrightarrow E'_{4j+1,4j+3,x} \subset E'_{4i+1,4i+3,x} \text{ for all } j > i \tag{5.3.3.5}$$

$$k(2i+2,x) = k(2i,x) \Leftrightarrow E'_{4j+1,4j+3,x} \cap E'_{4i+1,4i+3,x} = \emptyset \text{ for all } j > i \tag{5.3.3.6}$$

Proof. It is easily checked that (5.3.3.1) and (5.3.3.2) hold for the paths $\zeta_{4i+s}(x)$ which were found in 5.3.2. Also we have $k = k(2i+2,x) = k(2i,x)+2$ for exactly all the pairs $(\zeta_{4i+3}(x), \zeta_{4i+1}(x))$ for which $w'_{k-1}(\zeta_{4i+3}(x) = w'_{k-1}(x)$, as we saw in 5.3.2. We saw in 5.3.2 that (5.3.3.3) holds for these pairs and (5.3.3.4) holds for the others. Then (5.3.3.5) and (5.3.3.6) follow immediately from (5.3.3.1) to (5.3.3.4).

□

Lemma 5.3.4. *Once again, let $x = x_{r,\alpha}$ with $\alpha(0) = a$. The finite sequence of sets $\{E'_{4i_j-3,4i_j-1,x} : j \leq n\}$ is strictly decreasing, that is, satisfies*

$$E'_{4i_{j+1}-3,4i_{j+1}-1,x} \subset E'_{4i_j-3,4i_j-1,x}, \quad E'_{4i_{j+1}-3,4i_{j+1}-1,x} \neq E'_{4i_j-3,4i_j-1,x}$$

for all $1 \leq j < n$, if and only if $k(2i_j,x) < k(2i_j+2,x)$ for all $j < n$, that is, if and only if $w'_{k-1}(\zeta_{4i_j-3}(x)) = w'_{k-1}(x)$ for $k = (2i_j,x)$ for all $1 \leq j < n$.

Now let $\ell \geq 1$, and let $n_1(\ell) = n_1(\ell,\alpha)$ be the largest even integer such that

$$|w'_{k(j,x)-1}(\zeta_{2j-1}(x))| = |w'_{k(j,x)-1}(\zeta_{2j}(x))| < |v_\ell|$$

for $2j \leq n_1 - 1$. (By 5.3.2, $n_1(1) = 4$ and $n_1(\ell) = 2^\ell + 2^{\ell-1}$ for $\ell \geq 2$ is independent of α). Then the following hold.

1. *Let $u = dd$ or dv_0t_0 or av_0t_0 or bv_0t_0. Then*

$$D(t_{\ell,\alpha}u) \subset E'_{2n_1-3,2n_1-1,x} \Leftrightarrow \alpha(\ell) = \begin{cases} a \text{ if } \ell = 1, \\ b \text{ if } \ell > 1. \end{cases}$$

   Exactly similar results hold for $u_{\ell,\alpha}$.

2. *Let $\ell > 1$ and let α_1 and α_2 be such that $\alpha_1(\ell') = \alpha_2(\ell')$ for $\ell' < \ell$, but $\alpha_1(\ell) = b$ and $\alpha_2(\ell) = a$, so that $t_{\ell,\alpha_1} = t_{\ell,\alpha_2}$. Write $x_1 = x_{r,\alpha_1}$ and $x_2 = x_{r,\alpha_2}$. Write $n_1(\ell) = n_1$. Then*

$$D(v_{\ell,\alpha_1}) \subset E'_{2n_1-3,2n_1-1,x_1},$$

$$D(v_{\ell,\alpha_2}) \subset E'_{2n_1-3,2n_1-1,x_2} \subset E'_{2n_1-3,2n_1-1,x_1},$$

but

$$D(v_{\ell,\alpha_1}) \cap E'_{2n_1-3,2n_1-1,x_2} = \emptyset.$$

Hence, the number of sets $E'_{4i-3,4i-1,x_1}$ *containing* $D = D(t_{\ell,\alpha_1}L_3L_2) = D(t_{\ell,\alpha_2}L_3L_2)$ *is one more than the number of sets* $E'_{4i-3,4i-1,x_2}$ *containing* D. *These numbers are* $2^{\ell-1}$ *and* $2^{\ell-1}-1$ *respectively. In contrast, the number of sets* $E'_{4i-3,4i-1,x_1}$ *containing* $D' = D(v_{\ell,\alpha_1})$ *is the same as the number of sets* $E'_{4i-3,4i-1,x_2}$ *containing* $D(v_{\ell,\alpha_2})$. *These numbers are both* $2^{\ell-1}$.

PROOF. The first statement follows immediately from 5.3.3, in particular, from (5.3.3.5) and (5.3.3.6).

Now we consider the proof of 1. First consider the case $\ell = 1$. Since we are taking $\alpha(0) = a$, the set $E_{1,3} = E'_{1,3}$ is the same whether $\alpha(1) = a$ or b. In both cases, we have $w'_2(\zeta_i) = v_0L_3L_2R_3L_3^3$ for $3 \leq i \leq 6$ and $k(\zeta_{2i-1}, \zeta_{2i}) = 3$ for $2 \leq i \leq 4$. If $\alpha(1) = a$ then

$$w'_2(\zeta_7) = w'_2(\zeta_8) = v_0t_0aL_3 = v_0L_3L_2R_3L_3^2a^2L_3.$$

In this case, for any path $\zeta \in R_{m,0}$ such that ζ intersects $D(t_1u)$ for any u as in 1, all unit-disc crossings from the second onwards (determined by $w'_2(\zeta)$) are in $D(t_1) = D(t_{1,\alpha})$, and this second unit-disc crossing is between those of ζ_6 and ζ_7 (determined by $w'_2(\zeta_6)$ and $w'_2(\zeta_7)$). We therefore have $\zeta_6 < \zeta < \zeta_7$, and all of ζ, from the second unit-disc crossing onwards, is in $E'_{5,7}$ and we therefore have $D(t_1u) \subset E'_{5,7}$ for all such u.

If $\alpha(1) = b$, then

$$w'_2(\zeta_7) = w'_2(\zeta_8) = v_0t_0L_2R_3L_3 = v_0L_3L_2R_3L_3^2aL_2R_3L_3.$$

We still have $\zeta_6 < \zeta_7$. Now for any other path ζ such that ζ intersects $D(t_1) = D(t_{1,\alpha})$,we have $w'_2(\zeta) = w'_2(\zeta_7) = w'_2(\zeta_8)$ and, since the third unit-disc crossing ζ_8 is the furthest right of all such paths, the third unit disc crossing of ζ is to the left of the third unit-disc-crossing of ζ_8, but to the right of the second unit-disc-crossing of ζ. We therefore have $\zeta_8 < \zeta$ and hence ζ does not intersect $E'_{5,7}$ in this case, and we deduce that $D(t_1u) \cap E'_{5,7} = \emptyset$ in this case.

Now let $\ell \geq 2$. Then, writing $n_1(\ell) = n_1$,

$$k(n_1-1, x) = k(n_1, x) - 2 = k(\zeta_{2n_1,-3}, \zeta_{2n_1-1}),$$

and we have

$$\zeta_{2n_1-1} < \zeta_{2n_1} < \zeta_{2n_1-3} < \zeta_{2n_1-2},$$

which is different from the case $\ell = 1$. Write $k = k(n_1, x)$. Write $t_{\ell,\alpha} = zdd$. For all the choices considered of u, we have

$$w'_{k-1}(t_{\ell,\alpha}u) = zdL_2R_3L_3u_1,$$

for a word u_1 depending on u. We also have

$$w'_{k-1}(\zeta_{2n_1-1}) = w'_{k-1}(\zeta_{2n_1}) = \begin{cases} zdaL_3 \text{ if } \alpha(\ell) = a, \\ zdL_2R_3L_3 \text{ if } \alpha(\ell) = b. \end{cases}$$

For any path $\zeta \in R_{m,0})$ which intersects $D(t_{\ell,\alpha}u)$, we have

$$w'_{k-2}(\zeta) = w'_{k-2}(\zeta_{2n_1-1}),$$

and hence, for the same reasons as when $\ell = 1$ we have $\zeta_{2n_1} < \zeta$ if $\alpha(\ell) = b$ and $\zeta < \zeta_{2n_1-1}$ if $\alpha(\ell) = a$. But because of the difference between the ordering of ζ_{2n_1-3} and ζ_{2n_1-1} from the case of $\ell = 1$, we have

$$D(t_{\ell,\alpha}u) \subset E'_{2n_1-3,2n_1-1,\alpha} \text{ if } \alpha(\ell) = b, \tag{5.3.4.1}$$

$$D(t_{\ell,\alpha}u) \cap E'_{2n_1-3,2n_1,\alpha} = \emptyset \text{ if } \alpha(\ell) = a. \tag{5.3.4.2}$$

Now we consider the proof of 2. If $x_1 = x_{r,\alpha_1}$ and $x_2 = x_{r,\alpha_2}$ where $\alpha_1(\ell') = \alpha_2(\ell')$ for $\ell' < \ell$ but $\alpha_1(\ell) = b$ and $\alpha_2(\ell) = a$, then for the same reasons as before, we have

$$E'_{2n_1-3,2n_1-1,x_2} \subset E'_{2n_1-3,2n_1-1,x_1}.$$

and

$$D(v_{\ell,\alpha_j}) \subset E'_{2n_1-3,2n_1-1,x_j},$$

for $j = 1$, 2, and

$$D(v_{\ell,\alpha_1}) \cap E'_{2n_1-3,2n_1-1,\alpha_2} = \emptyset.$$

From (5.3.4.1) and (5.3.4.2), the number of sets $E'_{4i-3,4i-1,x_1}$ which contain $D(t_{\ell,\alpha_1})$, which is a decreasing sequence of sets ending in $E'_{2n_1-3,2n_1-1,x_1}$, is one more than the number of sets $E'_{4i-3,4i-1,x_2}$ which contain $D(t_{\ell,\alpha_2})$ (which is again a decreasing sequence of sets ending in $E'_{2n_1-3,2n_1-1,x_2}$). We have seen that the first number is the number of i with $2i \leq n_1$ such that $\zeta_{4i-3} < \zeta_{4i-5}$, that is the number of i such that $E'_{4i-3,4i-1,x_1}$ is of type A, that is, by 5.3.2, $2^{\ell-1}$ sets if $\ell \geq 2$ and just one set if $\ell - 1$. The number of sets $E'_{4i-3,4i-1,x_2}$ which contain $D(t_{\ell,\alpha_2})$ is one less if $\ell \geq 2$ and one more if $\ell = 1$, that is, $2^{\ell-1} - 1$ sets if $\ell \geq 2$ and two sets if $\ell = 1$.

□

The following lemma follows from 4.7.1.

Lemma 5.3.5. *Let $x = x_{r,\alpha}$ for any α and r. If $uE'_{4j-3,4j-1,x}$ intersects $I(\zeta_{4i-1-2s}(x), \zeta_{4i-2s}(x))$ for at least one of $s = 0$ or 1 then*

(5.3.5.1)

$$I(\zeta_{4i-1}(x), \zeta_{4i}(x)) \cup I(\zeta_{4i-3}(x), \zeta_{4i-2}(x)) \cup E'_{4i-3,4i-1,x} \subset uE'_{4j-3,4j-1,x}$$

5.4. The β sequence for $x_{r,\alpha}$

We now interpret 4.10.2 for $\gamma = \gamma^{r,\alpha}$, for any $\alpha \in \{a, b\}^{\mathbb{N}}$, and obtain more precise information about the paths $(\beta_{2i-1}(\gamma), \beta_{2i}(\gamma))$. We write $x = x_{r,\alpha}$ when no confusion can arise. In particular, we give some information about how the sequences $\{(\beta_{2i-1}(\gamma), \beta_{2i}(\gamma))\}$ and

$$\{(\psi_{1,4\lfloor(i-1)/2\rfloor-1,\gamma}(\beta_{2i-1}(\gamma)), \psi_{1,4\lfloor(i-1)/2\rfloor-1,\gamma}(\beta_{2i}(\gamma)))\}$$

differ from the sequence $\{(\zeta_{2j-1}(x), \zeta_{2j}(x))\}$. We sometimes write $\beta_i(x_{r,\alpha})$ or $\beta_i(x)$ for $\beta_i(\gamma) = \beta_i(\gamma^{r,\alpha})$, or even β_i, if no confusion can arise.

THEOREM 5.4.1. *Let $x = x_{r,\alpha}$ for any r and α and et $\gamma = \gamma^{r,\alpha}$. Then for any $i \geq 1$ with $2i \leq n(\gamma)$, paired satellite arcs for $\psi_{1,4i-1,\gamma}$ of type 3b) as in 4.10.2 do not occur. Consequently every quadruple $(\beta_{4i+t}(\gamma) : -3 \leq t \leq 0)$ is either an upper or lower ζ quadruple, or a type C quadruple. Type C quadruples only occur if $\alpha(0) = b$. Moreover, the following hold. Cases 2 and 3 only occur for satellite quadruples over x.*

1. *$\beta_{4i+t}(x) = \zeta_{4j+t}(x')$ for $-3 \leq t \leq 0$ and for x' and j depending on i, where $(\zeta_{4j+t}(x') : -3 \leq t \leq 0)$ is an upper ζ quadruple for x'. If $\alpha(0) = a$, then every subword v_0t_0 in x is a subword in the same position in x'. If $\alpha(0) = b$, then every subword w_0 in x is replaced by v_0 in the same position in x'. Once that is done, some occurrences of a or b preceding occurrences of v_0t_0 may be replaced by b or a*
2. *$I(\beta_{4i+t-1}(x), \beta_{4i+t}(x)) = uI(\zeta_{4j+t-1}(x'), \zeta_{4j+t}(x'))$ for $t = -2$ and $t = 0$, and for x' and j depending on i, where all letters of u are in $\{L_3, L_2, R_3\}$, and an odd number of letters are in $\{L_3, L_2\}$, and with the same restrictions on ux' as in 1.*
3. *For some $k \leq k(2i-1, x)$, the value of $w'_k(\beta_{4i+t}(x))$ (for $-3 \leq t \leq 0$) is maximal or minimal subject to the value of $w'_{k-1}(\beta_{4i+t}(x))$*

For any $1 \leq \ell \leq i$, the support of $\psi_{4\ell-3,4i-1,x}$ can only intersect $D(x')$ for any x' as above, if $(\beta_{4i+t}(x) : -3 \leq t \leq 0)$ is of form 1 or 2. Moreover, in each of these cases, the pair of principal arcs of $\psi_{1,4i-5,x}\cdot(\beta_{4i+t-1}(x), \beta_{4i+t}(x))$ (for $t = -2$, 0) is of one of the following forms, up to isotopy preserving $S^1 \cup Z_\infty$:

a) $I(\zeta_{4\ell+t-1}(x), \zeta_{4\ell+t}(x))$ for some ℓ, and $t = -2$, 0, or is in the image of $I(\zeta_{4\ell+t-1}(x'), \zeta_{4\ell+t}(x'))$ under a single disc exchange corresponding to a basic exchange $a \leftrightarrow b$, where x' is the image of x under that exchange;

b) both arcs are parallel to $I(\zeta_{4\ell-3}(x), \zeta_{4\ell-2}(x))$ for some ℓ;

c) both arcs are parallel to a common arc of $\zeta_{4\ell+s}(x)$ for $-3 \leq s \leq 0$ for some ℓ where the arc is between S^1 crossings, up to or before the $k(4\ell-3, x) - 1$'th crossing.

Similar properties hold if $\psi_{4\ell-3,4i-1}$ *is replaced by* ψ' *for any decomposition* $\psi_{4\ell-3,4i-1} = \psi'' \circ \psi'$*, where each of* ψ' *and* ψ'' *is a composition of disc exchanges.*

Remark The difference between 1 and 2 is the difference between the supports of disc exchanges spanning the upper and lower circle. The extra possibility at the end of a) arises because of the differences in the sequences $\zeta_j(x_{r,\alpha})$ and $\zeta_j(x_{r,\alpha'})$ if $\alpha(k) \neq \alpha'(k)$ for some k.

PROOF. Naturally, this is proved by induction. We can start the induction because the results are true for $i = 1$. Now suppose the results are true for $(\beta_{4j+t}(\gamma) : -3 \leq t \leq 0)$, and for $\psi_{1,4j-1,\gamma}$, for all $j < i$. As in 4.10.2, we aim to obtain the results for $(\beta_{4i+t}(\gamma) : 1 \leq t \leq 4)$ and for $\psi_{1,4i+3,\gamma}$, by using the information already obtained about $\psi^{-1}_{1,4i-5,\gamma}$, and about the disc exchanges of which it is a composition. We need to consider the action of $\psi^{-1}_{1,4i-5,\gamma}$ on $D(x)$.

We define x' to be *of permissible form* if x' is obtained from x_r by replacing some occurrences of a or b preceding occurrences of v_0t_0 by b or a. This is equivalent to the description of x' in terms of $x_{r,\alpha}$ in statement 1 of the theorem. By the inductive hypothesis, the support of each disc exchange in the composition which can have an effect on $D(x)$ is of the form $C(ya \leftrightarrow yb, E'_{4j-3,4j-1,z})$ where $E'_{4j-3,4j-1,z} = E_{\zeta_{4j-3}(z),\zeta_{4j-1}(z)}$ and y is some admissible word. Then we see that in order to have an effect on $D(x)$, the first disc exchange ξ_1 in the composition whose support intersects $D(x)$ must have ya, yb and z of permissible form, and then $\xi_1(D(x)) = D(x^1)$ where x^1 is also of permissible form. The exchanges for $\beta_j(\gamma)$ are all basic exchanges, because $w'_1(\beta_j) = w'_1(\beta_3) = L_3$ for all $j \geq 3$. Write $\psi^{-1}_{1,4i-5}\xi_n \circ \cdots \xi_1$ on $D(x)$. and write $\xi_{\ell,1} = \xi_\ell \circ \cdots \xi_1$. Let the support of ξ_ℓ be $C(y_\ell a \leftrightarrow y_\ell b, E_j)$, where $E_j = E_{4j_1(\ell)-3,4j_1(\ell)-1,z_\ell}$ for $e_\ell \subset \{a, b\}$, so that $\xi_{\ell,1}(D(x)) = D(x^\ell)$ and $x^\ell = y_{\ell+1}e'_{\ell+1}z_{\ell+1}$ for $e'_{\ell+1} \in \{a, b\}$, as it is in the support of the next disc exchange $\xi_{\ell+1}$. We do not have E_j of the form $E_{4n-3,4n-1,\gamma}$ for $(\beta_{4n+t} : -3 \leq t \leq 0)$ as in 3 in the statement of the lemma, because in that case, by the form of $y_\ell e_\ell z_\ell$, $uE_{4n-3,4n-1,x}$ does not intersect $D(y_\ell e_\ell z_\ell)$ for any ℓ. Because of the form of the words $y_\ell e_\ell b_\ell$, the set $y_\ell e_\ell E_\ell$ contains any set $E'_{4j_2-3,4j_2-1,x^{\ell+1}}$ that it intersects, and if the two sets intersect and are on the same level, so that the topmost principal pairs of the two coincide, then they coincide. It follows that principal arcs of type 3b) of 4.10.2 do not occur for $\xi_{\ell,1}$ for any ℓ, and hence they do not occur for $\psi_{1,4i-5,\gamma}$.

The types 2 and 3 of quadruples $(\beta_{4i+t}(\gamma) : -3 \leq t \leq 0)$ only occur because they are in the boundaries of sets $C(v \leftrightarrow v', E)$, and hence give satellite arcs. □

5.4.2. Satellites without further effect. From Theorem 5.4.1 there are two kinds of satellite quadruples for $\gamma = \gamma^{r,\alpha}$: those as in 2 or 3 of the lemma. If $(\beta_{4i+t}(\gamma) : -3 \leq t \leq 0)$ is as in 2 of the theorem, then some of the disc exchanges in the composition for $\psi_{4i-3,4i-1}^{-1}$ might be in the composition for $\psi_{1,4j-1}^{-1}$ which is not the identity on $D(x_{r,\alpha})$. But for quadruples $(\beta_{4i+t}(\gamma) : -3 \leq t \leq 0)$ as in 3 of the theorem this is impossible. We therefore refer to these as *satellite (quadruples) without further effect.*

The result of 5.4.1 can be improved. We have the following. In particular, this implies that x' in the statement of 5.4.1 can almost always be taken of the form $x_{r,\alpha'}$ for some $\alpha' \in \{a,b\}^{\mathbb{N}}$. For an $\alpha \in \{a,b\}^{\mathbb{N}}$, we write $n(\ell,\alpha)$ for the largest integer such that

$$D(v_{\ell-1,\alpha}t_{\ell-1,\alpha}) \subset \psi_{1,4\lfloor n(\ell)-1)/2\rfloor-1}(D(\beta_{2n(\ell)-1},\beta_{2n(\ell)})).$$

THEOREM 5.4.3. *Fix* $\alpha \in \{a,b\}^{\mathbb{N}}$. *Then, for* $2p \leq n(\ell)$,

$$\psi_{1,4p-1,\alpha} \mid \psi_{1,2n(\ell)-1,\alpha}^{-1}(D(v_{\ell-1,\alpha}t_{\ell-1,\alpha}))$$

can be written in the form

$$\xi'_{q_{p,\ell},p\ell} \circ \cdots \xi'_{1,p,\ell},$$

where each $\xi'_{j,p,\ell}$ *is a composition of disc exchanges* $\xi_{i,j,p,\ell}$ *with supports* $C(z_{j,\ell}a \leftrightarrow z_{j,\ell}b, E'_{i,j,p,\ell})$ *where the following hold.*

1. $|z_{j,\ell}|$ *is increasing with* j;
2. $z_{j,\ell}e$ *is a prefix of* x *for one of* $e = a$ *or* b;
3. *Each* $E_{i,j,p,\ell}$ *is of the form* $E'_{4j_1-3,4j_1-1,x_{i,j,p,\ell}}$ *for some* j_1 *depending on* i, *where* $x_{i,j,p,\ell} = x_{\theta_{i,j,p,\ell}}$ *for* $\theta_{i,j,p,\ell} \in \{a,b\}^{\mathbb{N}}$ *which can be taken to be* α *except when* $|z_{j,\ell}a| = |v_k|$ *for some* k, *when we can take* $\theta_{i,j,p,\ell}(s) = \alpha(s)$ *for* $s \neq k$.
4. $$\begin{aligned}\psi_{1,4p-1,\alpha}^{-1}&(D(v_{\ell-1,\alpha}t_{\ell-1,\alpha}a) \cup D(v_{\ell-1,\alpha}t_{\ell-1,\alpha}b))\\ &= D(v_{\ell-1,\theta}t_{\ell-1,\theta}a) \cup D(v_{\ell-1,\theta}t_{\ell-1,\theta}b)\end{aligned}$$
   for some $\theta = \theta_{\ell,p,\alpha} \in \{a,b\}^{\mathbb{N}}$ *such that* $\theta_{\ell_1,p_1,\alpha}(i) = \theta_{\ell_2,p_2,\alpha}(i)$ *for* $i \leq \ell_1 \leq \ell_2$ *and* $n(\ell_1) \leq 2\min(p_1,p_2)$.

- 5\. *A similar result to 4 holds for* $D(v_{\ell-1,\alpha}t_{\ell-1,\alpha}eu)$ *where* $e \in \{a,b\}$ *and* u *is any prefix of* $t_{\ell,\alpha}$. *This time, the image of* $D(v_{\ell-1,\alpha}t_{\ell-1,\alpha}eu)$ *under* $\psi_{1,4s-1,\alpha}^{-1}$ *is of the form* $D(v_{\ell-1,\theta}t_{\ell-1,\theta}e'u'$ *where* $e' \in \{a,b\}$ *and* u' *is a prefix of* $t_{\ell,\theta}$.

1. The proof involves rearranging the disc exchanges in the composition

$$\psi_{1,4p-1,\alpha} = \psi_{1,3,\alpha} \circ \cdots \circ \psi_{4r-3,4r-1,\alpha}.$$

Since each $\psi_{4i-3,4i-1,\alpha}$ is a composition of disc exchanges, we already have a representation of $\psi_{1,4p-1,\alpha}$ as a composition of disc exchanges $\xi'_q \circ \cdots \xi'_1$.

For such a composition we write $\xi'_{i,1} = \xi'_i \circ \cdots \circ \xi'_1$. First, we summarise what we know from 5.4, given that we are looking at the restriction of $\psi_{1,4r-1,\alpha}$ to $\psi^{-1}_{1,4r-1}(D(v_{\ell,\alpha}t_{\ell,\alpha})$. It follows from this that each disc exchange in the composition for $\psi_{4j-3,4j-1}$, for each j, has support of the form $C(za \leftrightarrow zb, E_{4j_1-3,4j_1-1,y})$ for some j_1, z and y, where each of z and y is obtained from a prefix of $x_{r,\alpha}$ (or, equivalently, of x_r) by replacing occurrences of a or b before some subwords $v_{1,\alpha}$ or $t_{1,\alpha}$ by b or a. So the same is true for the exchanges in the full decomposition for $\psi_{1,4n(\ell)-1,\alpha}$. So the support of ξ'_j is of the form $C(z'_ja \leftrightarrow z'_jb, E_{4j_1-3,4j_1-1,y'_j})$, where each z'_j and y'_j have the properties just stated for z and y respectively. However, we do not have from this that $|z'_j|$ is non-decreasing with j. In order to get this, we need to be able to commute a disc exchange ξ_{j_1,i_1} in the composition for $\psi_{4i_1-3,4i_1-1,x}$ with ξ_{j_2,i_2} in the composition for $\psi_{4i_2-3,4i_3-1,x}$, if $i_1 < i_2$ and the supports are $C_1 = C(z'_{j_1,i_1}a \leftrightarrow z'_{j_1,i_1}b, E_{4i_1-3,4i_1-1,x})$ and $C_2 = C(z'_{j_2,i_2}a \leftrightarrow z'_{j_2,i_2}b, E_{4i_2-3,4i_2-1,x})$ with $|z'_{j_1,i_1}| < |z'_{j_2,i_2}|$. We only have to do this restricted to $\xi'(D(v_{\ell,\alpha}t_{\ell,\alpha}))$, where ξ' is the composition of disc exchanges applied before ξ_{ℓ_2,i_2}. Commutativity holds if C_2 is contained in $z'_{j_1,i_1}eE_{4i_1-3,4i_1-1,x}$ (for $e = a$ or b) whenever it intersects it. This is true if

$$D(z'_{j_2,i_2}) \cap z'_{j_1,i_1}eE_{4i_1-3,4i_1-1,x} \neq \emptyset \Rightarrow D(z'_{j_2,i_2}) \subset z'_{j_1,i_1}eE_{4i_1-3,4i_1-1,x}.$$

Write y''_s and y'_s for the longest and shortest words respectively with

$$D(y''_s) \subset E_{4i_s-3,4i_s-1,x} \subset D(y'_s).$$

Then it suffices to show that if $i_1 < i_2$ and $|z_{j_1,i_1}| < |z'_{j_2,i_2}|$ and $z'_{j_1,i_1}ey'_1$ is a prefix of z'_{j_2,i_2} or z_{j_2,i_2} is a prefix of $z'_{j_1,i_1}ey''_1$ then $z'_{j_1,i_1}ey''_1$ is a prefix of z'_{j_2,i_2}. To do this we use properties of the set A of words y which are obtained from x_r by replacing some occurrences of a preceding subwords v_1 or t_1 by b. For each α, $x_{r,\alpha}$ is such a word, but the set of words y is considerably larger than the set of words $x_{r,\alpha}$. Nevertheless, any such word has strings of d's in certain positions, and this means that that the set A still has strong non-recurrence properties. In fact, the following holds. Suppose z and y are both prefixes of words in A. Then zy can only be a prefix of a word in A if z has the same length as a word of x_r which ends in v_i for some $i \geq k$, where k is such that $|v_{k-1}| < |y| \leq |v_k|$. The reason is simply that, for any $y \in A$, a string of at least k d's only occurs in the same position as in x_r, and any occurrence in x_r is at the end of a subword v_i for some $i \geq k$.

So now we apply this with $z = z'_{\ell_1,i_s}e_s$ and $y = y'_s$ for $s = 1$, 2 under the assumption that $|z_{j_1,i_1}| < |z'_{j_2,i_2}|$ and $i_1 < i_2$. We let k_s be such that $|v_{k_s-1}| \leq |y'_s| < |v_{k_s}|$ Then the assumption that $i_1 < i_2$ means that $k_1 \leq k_2$. In fact if $(\beta_{4i_1+t} - 3 \leq t \leq 0)$ is non-satellite then $|y'_1| < |y'_2|$,and if $(\beta_{4i_1+t} : -3 \leq t \leq 0)$ is satellite then from the way that satellites are produced we see that $|v_{k-1}| < |y_1'| \le |v_k|$ whenever $|v_{k-1}| < |y_2'| \le |v_k|$, that is $k_1 \le k_2$. It follows that if $|z_{j_1,i_1}| < |z'_{j_2,i_2}|$, then

$$|z_{j_1,i_1}e_1| + |v_{k_1}| \le |z_{j_1,i_1}e_1| + |v_{k_2}| < |z_{j_2,i_2}e_2|$$

and hence

$$|z_{j_1,i_1}e_1y_1''| < |z_{j_2,i_2}e_2|$$

as required.

The proof of 2, 3, 4 and 5 follow by induction. Once we know $\psi^{-1}_{1,4n(\ell)-1,x}$, then we know the prefixes $z_{j,\ell+1}$ for all j with $|z_{j,\ell+1}a| < |v_{\ell+1}|$. We have $z_{j,\ell} = z_{j,\ell+1}$ for $|z_{j,\ell+1}a| < |v_{\ell+1}|$. Moreover the disc exchanges in the decomposition for $\psi_{1,4r-1,\alpha}$ with supports of the form $C(z_{j,\ell}a \leftrightarrow z_{j,\ell}b, E)$ are the same for any $|z_{j,\ell}| \le |v_\ell|$ if $2n(\ell) \le r$. Now the disc exchanges in the composition for $\psi^{-1}_{1,4r-1}$ which affect the choice of $\theta = \theta_{\ell,r,\alpha}$ are those corresponding to prefixes $|z_{j,\ell}| \le |v_\ell|$. It follows that $\theta_{\ell,r,\alpha}(i) = \theta_{\ell,s,\alpha}(i)$ if $i \le \ell$ and $2n(\ell) \le \min(r,s)$, and also $\theta_{\ell_1,r,\alpha}(i) = \theta_{\ell_2,r,\alpha}(i)$ if $i \le \ell_1 \le \ell_2$ and $2n(\ell_1) \le r$. The result follows. □

From **??**, and in particular from 4 and 5 of **??**, we have the following corollary. The proof is immediate.

Corollary 5.4.4. *Let* $\alpha \in \{a,b\}^{\mathbb{N}}$ *and let* $x = x_{r,\alpha}$. *Let* $(\beta_{4i+t}(x) : -3 \le t \le 0)$ *be any quadruple which is not a satellite quadruple without further effect, such that with* $n(\alpha,\ell) < 2i$ *for some* $\ell \ge 1$. *Then there is* $\theta = \theta_{i,\alpha} \in \{a,b\}^{\mathbb{N}}$ *and* $j = j(i)$ *such that* $(\beta_{4i+t}(x) : -3 \le t \le 0)$ *is an* ζ *quadruple* $(\zeta_{4j+t}(x_{r,\theta_i}) : -3 \le t \le 0)$*, which is an upper quadruple if* $(\beta_{4i+t}(x) : -3 \le t \le 0)$ *is non-satellite. Moreover* $\theta_{i_1,\alpha}(s) = \theta_{i_2,\alpha}(s)$ *for all* $s \le \ell - 1$ *if* $n(\alpha,\ell) \le \min(2i_1, 2i_2)$.

Moreover

$$(\beta_{2n(r+1,\alpha)-1}(\gamma), \beta_{2n(\gamma)}(\gamma)) = (\zeta_{2n_1(r+1,y)-1}(y), \zeta_{2j}(y)).$$

for $y = x_{r,\theta}$ *and* $\theta = \theta_{\lfloor n(\alpha,r+1)/2 \rfloor,\alpha}$.

It follows from Theorem 5.2 that $\theta_{\lfloor n(\alpha,r+1)/2 \rfloor,\alpha}$ is independent of α, or can be assumed so, that is, $\theta_{\lfloor n(\alpha,r+1)/2 \rfloor,\alpha}(s)$ is independent of α for $0 \le s \le r$. We write α^* for this common value of $\theta_{\lfloor n(\alpha,r+1)/2 \rfloor,\alpha}$. Then combining the above with Theorem 5.2, we have the following corollary.

Corollary 5.4.5. *There is* $\alpha^* \in \{a,b\}^{\mathbb{N}}$ *with the following property. For all* $\alpha \in \{a,b\}^{\mathbb{N}}$*, we have*$\theta_{i,\alpha}(s) = \alpha^*(s)$ *for* $s \le \ell - 1$*, if* $n(\ell,\alpha) \le 2i$*, and* $\theta_{i,\alpha}(s) = \alpha(s)$ *for all* $s \le r$ *if* $i = \lfloor n(r+1,\alpha)/2 \rfloor$.

I have not been able to compute α^*. This is mainly because of the difficulty of detemining accurately the satellites with effect. The only thing which seems certain is that $\alpha^*(0) = \alpha^*(1) = a$ and $\alpha^*(2) = b$. As the notation suggests, α^* is independent of r, that is, the values of $\alpha^*(s)$ for $s \leq r$ are the same for $x_{\alpha,q}$ for all $q \geq r$, and, of course, for all $\alpha \in \{a, b\}^{\mathbb{N}}$.

CHAPTER 6

The main theorems

The theorems from which our main Theorem 1.9 is deduced are the following.

Theorem 6.1. *The set of capture paths $\gamma \in \mathcal{Z}_m(3/7,+,+,0)$ for which there is no other inequivalent capture path ζ such that $\sigma_\zeta \circ s$ and $\sigma_\gamma \circ s$ are Thurston equivalent, is of positive density bounded from* 0.

More precisely, there are capture paths $\gamma \in \mathcal{Z}_m(3/7,+,+,0)$ such that the following hold. There is no other capture path $\zeta \in \mathcal{Z}(3/7,+,+,0)$ (up to equivalence) such that $\sigma_\gamma \circ s$ and $\sigma_\zeta \circ s$ are Thurston equivalent, and there is a decreasing sequence of neighbourhoods U of a point P in the boundary of the gap of $L_{3/7}$ containing the endpoint x of γ such that

$$\lim_{U\to P}\lim_{n\to\infty}\frac{\#(V(U)\cap Z_n)}{\#(U\cap Z_n)}=1$$

where $V(U)\subset U$, and $y\in V(U)\cap Z_\infty$ if and only if the following hold. The point y is the endpoint of a capture path $\gamma(y) \in\in \mathcal{Z}(3/7,+,+,0)$ such that there is no other capture path $\zeta \in \mathcal{Z}(3/7,+,0)$ (up to equivalence) such that $\sigma_{\gamma(y)} \circ s$ and $\sigma_\zeta \circ s$ are Thurston equivalent.

Theorem 6.2. *For each integer $r \geq 0$, and for all sufficiently large m given r, the set of capture paths $\gamma = \gamma_1 \in \mathcal{Z}_m(3/7,+,+,0)$ such that $\sigma_\gamma \circ s$ is equivalent to $\sigma_{\gamma_i} \circ s$ for exactly 2^{r+1} inequivalent capture paths γ_i for $1 \leq i \leq 2^{r+1}$, and with $\gamma_i \in \mathcal{Z}_m(3/7,+,+,0)$ for $1 \leq i \leq 2^{r+1}$, and to no path in $\mathcal{Z}_m(3/7,+,-,0)$ with S^1 crossing at $e^{2\pi it}$ for $t \in [\frac{39}{56},\frac{5}{7})$, is of positive density bounded from* 0.

More precisely there is a capture path $\gamma_1 \in \mathcal{Z}_m(3/7,+,+,0)$ such that $\sigma_\gamma \circ s$ is equivalent to $\sigma_{\gamma_i} \circ s$ for exactly $2r$ inequivalent capture paths γ_i for $1 \leq i \leq 2r$, with endpoints x_i, and $\gamma_i \in \mathcal{Z}(3/7,+,+,0)$ for all $i \leq 2^{r+1}$, and there are neighbourhoods U_i of points P_i in the boundaries of the gaps of $L_{3/7}$ containing the points x_i, and a sequence of neighbourhoods of P_i in U_i, converging to P_i such that, for U_i' in this sequence,

$$\lim_{U_i'\to P_i}\lim_{n\to\infty}\frac{\#(V(U_i':U_1,\cdots U_{2N})\cap Z_n)}{\#(U_i'\cap Z_n)}=1$$

where U_i' *is in this sequence of neighbourhoods, and* $V(U_i' : U_1, \cdots U_{2r})$ *is the set of points* $y_i \in U_i' \cap Z_\infty$ *with the following property. The point* y_i *is the endpoint of a capture path* $\gamma(y_1) \in \mathcal{Z}(a_1, +, +)$ *and there are exactly* 2^{r+1} *inequivalent capture paths* $\gamma(y_j)$ *such that* $y_j \in U_j$ *for* $1 \leq j \leq 2^{r+1}$ *and* $y_i \in U_i'$ *and* $\sigma_{\gamma(y_i)} \circ s$ *are all Thurston equivalent, but there is no other inequivalent capture path* $\zeta \in \mathcal{Z}(3/7, +, 0)$ *such that* $\sigma_\zeta \circ s$ *and the* $\sigma_{\gamma(y_i)} \circ s$ *are Thurston equivalent. Moreover there are neighbourhoods* $U_j''(U_i')$ *of* y_j *with* $U_i''(U_i') = U_i'$ *such that*

$$V(U_i' : U_1, \cdots U_{2r}) = V(U_i', U_1''(U_i'), \cdots U_{2r}''(U_i'))$$

and

$$\lim_{U_i' \to y_i} U_j''(U_i) = y_j.$$

6.3. Reductions

The more precise statements make it likely that these are really the same theorem. This is in fact the case, given Theorem 5.2, which gives us capture paths γ_i for $1 \leq i \leq 2^{r+1}$ to work with, as in the more precise statement of 6.2. We will make choice of the path γ, its endpoint x and the point P of 6.1, in 6.6. Similarly, we will make a precise choice of the γ_i, the endpoints x_i and the P_i of 6.2, in 6.7. We will choose P of the same preperiod as x, and similarly for the P_i.

The usual way to construct sets of positive density is to use a convergent product: for example, a product of the form

$$\prod_{n=1}^{\infty} (1 - \exp(-(\log n)^\alpha)$$

for any $\alpha > 1$. This is the type of convergent product that we will use.

Let $\gamma \in \mathcal{Z}_m(3/7, +, +)$. A sufficient condition for $\sigma_\gamma \circ s$ not to be Thurston equivalent to $\sigma_{\gamma'} \circ s$ for any inequivalent capture path γ' is that, for all i,

(6.3.0.1)

$$(\beta_{2i-1}(\gamma), \beta_{2i}(\gamma)) = (\beta_{2i'-1}(\gamma'), \beta_{2i'}(\gamma')) \Rightarrow i = i' \wedge \beta_j(\gamma) = \beta_j(\gamma') \forall j \leq 2i.$$

To see that this is sufficient, the pairs $(\beta_{2i-1}(\gamma), \beta_{2i}(\gamma))$ are defined so that $\beta = \beta_{2n(\gamma)-1}(\gamma)$ is the path in $R_{m,0}$ such that $\sigma_\beta \circ s$ is Thurston equivalent to $\sigma_\gamma \circ s$. Therefore, if $\sigma_\gamma \circ s$ and $\sigma_{\gamma'} \circ s$ are Thurston equivalent, we have $n(\beta) = n(\gamma) = n$ and

$$\beta = \beta_{2n-1}(\gamma) = \beta_{2n-1}(\gamma')$$

Since $(\beta_{2n-1}(\gamma), \beta_{2n}(\gamma))$ is an adjacent pair in $R_{m,0}$ with $\beta_{2n-1}(\gamma) < \beta_{2n}(\gamma)$, and similarly for γ', we then have $\beta_{2n(\gamma)}(\gamma) = \beta_{2n(\gamma')}(\gamma')$, and hence (6.3.0.1) implies that

$$\beta_i(\gamma) = \beta_i(\gamma') \text{ for all } i \leq 2n(\beta).$$

Now (6.3.0.1) also implies that

(6.3.0.2)

$$(\beta_{2i-1}(\gamma), \beta_{2i}(\gamma)) = (\beta_{2i'-1}(\gamma'), \beta_{2i'}(\gamma')) \Rightarrow \psi_{4j-3,4j-1,\gamma} = \psi_{4j-3,4j-1,\gamma'}$$

for all $2j \leq i$. Hence if the condition holds we also have

$$\psi_{1,2i-1,\gamma} = \psi_{1,2i-1,\gamma'}.$$

In particular if (6.3.0.1) holds for γ then for $n = n(\gamma) = n(\gamma')$ we have

$$\beta_{2n-1}(\gamma) = \beta_{2n-1}(\gamma') \Rightarrow \psi_{1,2n-1,\gamma} = \psi_{1,2n-1,\gamma'}. \tag{6.3.0.3}$$

But then, denoting the endpoints of $\beta = \beta_{2n-1}(\gamma)$, and γ and γ' by $p(\beta)$ and so on (as usual), we have

$$p(\gamma) = \psi_{1,2n-1}^{-1}(p(\beta)) = p(\gamma')$$

and hence $\gamma = \gamma'$. The method for proving Theorems 6.1 and 6.2 is essentially to prove that Condition (6.3.0.1) holds with positive density for all i. This will be done by showing that, if A_i is the set for which (6.3.0.1) holds, then, for a suitable sequence n_i converging to ∞, the density of $A_{n_{i+1}}$ in A_{n_i} approaches 1 as i tends to ∞, sufficiently fast to yield a set of positive density.

Naturally enough, the positive density condition that we seek will be obtained by proving that a set of conditions which imply the sought condition, hold with positive density. The theorem which we will prove, which will imply both theorems 6.1 and 6.2, is the following. Here, $D(\zeta_1, \zeta_2)$ is as in 4.2.

THEOREM 6.4. *Let $U \subset \{z : \text{Im}(z) > 0\}$ be an open set in Z_∞, and let $\mathcal{U}$ be the set of capture paths in $\mathcal{Z}(3/7, +, +, 0)$ with endpoints in U. Let $n_0 \geq 1$. Suppose that $\beta_i(\gamma)$ is constant for $\gamma \in \mathcal{U}$ and $i \leq 4n_0 + 2$, so that $\psi_{1,4i-1,\gamma}$ is also constant for $2i \leq n_0$. Write*

$$(\beta_{0,1}, \beta_{0,2}) = (\beta_{4n_0+1}(\gamma), \beta_{4n_0+2}(\gamma))$$

for any $\gamma \in \mathcal{U}$, and write

$$D_0 = D(\beta_{0,1}, \beta_{0,2}).$$

Then

$$\limsup_{n\to\infty} d_n = d > 0,$$

where:

- $d_n = \dfrac{\#(U_1 \cap Z_n)}{\#(U \cap Z_n)}$;

- *$\mathcal{U}_1$ is the set of capture paths in $\mathcal{Z}(3/7,+,+,0)$ with endpoints in U_1 and $\gamma \in \mathcal{U}_1$;*
- *U_1 is the subset of U on which the following hold for suitable constants.*

There are strictly increasing sequences of integers n_i and $i(r)$, depending on γ, but with n_0 as before and, such that the following hold. Write

$$(\beta_{i,1},\beta_{i,2}) = (\beta_{4n_i+1}(\gamma),\beta_{4n_i+2}(\gamma)),$$

$$D_i = D(\beta_{i,1},\beta_{i,2}),$$

$$E_i = \cup\{E_{\omega_1,\omega_3} : (\omega_t : 1 \le t \le 4) \text{ is a quadruple, } (\omega_1,\omega_2) = (\beta_{i,1}(\gamma),\beta_{i,2}(\gamma))\}.$$

Let u_j be the longest word such that

$$D(\beta_{2j-1},\beta_{2j}) \subset D(u_j).$$

For suitable α_1 and $\alpha_2 > 1$, we have

$$|u_{n_{j+1}+1}| - |u_{2n_j+1}| \le (\log|u_{2n_j+1}|)^{\alpha_1}$$

$$|u_{2n_{i(r+1)}+1}| \ge \exp((\log|u_{2n_{i(r)}+1}|)^{\alpha_2})$$

Also, the following hold:

a) For all i, if C_1 is a component of

$$\cup\{C : C \text{ is a component of } (\sigma_{\alpha_{1,4\ell-3}}\circ s)^{-n}(E_{4\ell-3,4\ell-1}) \text{ with } \ell \le n_i,\ n > 0, D_i \not\subset C\},$$

or if $(\beta_{2j-1},\beta_{2j})$ is an inactive satellite, and C_1 is a component of

$$\cup\{C : C \text{ is a component of } (\sigma_{\alpha_{1,4\ell-3}}\circ s)^{-n}(E_{4\ell-3,4\ell-1}) \text{ with } 2\ell < j,\ n > 0, D_i \not\subset C\}$$

then for any principal arc I for an adjacent pair in $R_{m,0}$ which is inside D_i and over D_{i_1} for some $i_1 > i$, we have

$$(C_1 \cap I = \emptyset) \vee (C_1 \subset D_i \wedge (C_1 \subset E_{2n_i-3,2n_i-1} \vee C_1 \cap E_{2n_i-3,2n_i-1} = \emptyset)).$$

b) For any r, ℓ and j with $2n_{i(r-1)} < 2\ell < j \le 2n_{i(r+1)}$, such that $(\beta_{2j-1},\beta_{2j})$ is not an inactive satellite, and any $n > 0$ and any component C of $(\sigma_{\beta_{4\ell-3}} \circ s)^{-n}(E_{4\ell-3,4\ell-1})$, or $\sigma_{\beta_{i(r-1),1}} \circ s)^{-n}(D_{i(r-1)})$, we have

$$C \cap I(\beta_{2j-1},\beta_{2j}) = \emptyset.$$

Moreover, given n_0 we can find n_j for $j \ge 1$ so that d is a function $d(d^0)$ of $d^0 = \limsup_{m\to\infty} d_{m,0}$, where

$$d_{m,0} = \sup_{D_0 \subset U} \frac{\#((D_0 \setminus \cup_{n>0} s^{-n}(E_0)) \cap Z_m)}{\#(D_0 \cap Z_m)},$$

and

$$\lim_{d^0 \to 1} d(d^0) = 1.$$

6.5. Why 6.4 implies 6.1 and 6.2

Of course, it is not immediately obvious that Theorem 6.4 implies 6.1 and 6.2. The main purpose of the conditions on U_1 and $\mathcal{U}_1$ in 6.4 is to bound the effect of the homeomorphisms $\psi_{1,4i-1,\gamma}$ on sets $I(\beta_{2j-1}(\gamma), \beta_{2j}(\gamma))$ for $\gamma \in \mathcal{U}_1$. The following lemma is the key to showing that 6.4 implies 6.1 and 6.2.

We say that a pair $(\beta_{2j-1}, \beta_{2j})$ is an *inactive satellite* for γ if

$$\psi^{-1}_{1,4\lfloor j/2\rfloor-1}(p(\gamma)) \notin D(\beta_{2j-1}, \beta_{2j}).$$

Lemma 6.5.1. *The conditions on $\mathcal{U}_1$ in 6.4 imply the following for $\gamma \in \mathcal{U}_1$, where n_i is the sequence, which depends on γ, which is given in 2 of 6.4.*

1. *For all $j \geq 2n_i$,*

$$D(\beta_{2j-1}(\gamma), \beta_{2j}(\gamma)) \subset D_i.$$

Moreover $\psi_{4\ell-3,4\ell-1,\gamma}$ is the identity on $I(\beta_{2j-1}(\gamma), \beta_{2j}(\gamma))$ for all $n_i \leq \ell \leq n_{i+1}$ and $2n_k \leq j \leq 2n_{k+1}$, whenever $i(r) \leq i < k \leq i(r+2)$ for some r, and provided that $(\beta_{2j-1}, \beta_{2j})$ is not an inactive satellite.

2. *If*

$$\beta_\ell(\zeta) = \beta_\ell(\gamma) \text{ for all } \ell \leq 4n_0 = 4n_0(\gamma), \tag{6.5.1.1}$$

and

$$I(\beta_{2i-1}(\zeta), \beta_{2i}(\zeta)) \subset D_k, \tag{6.5.1.2}$$

for some i and k, then

$$\beta_j(\zeta) = \beta_j(\gamma) \text{ for all } j \leq 2n_k. \tag{6.5.1.3}$$

3. *If $\zeta \in \mathcal{U}$ and $\beta_{2n(\gamma)}(\gamma) = \beta_{2n(\zeta)}(\zeta)$ then $\zeta = \gamma$.*

Proof. 1. The proof of the first statement is by induction on j. The pair $(\beta_{2j-1}, \beta_{2j})$ is the pair (ω_1, ω_2) with the property that $I(\omega_1, \omega_2)$ is mapped by $\psi_{1,4\lfloor (j-1)/2\rfloor-1}$ to cross γ immediately inside the principal arc of $(\psi_{1,4\lfloor (j-2)/2\rfloor-1}(\beta_{2j-3}), \psi_{1,4\lfloor (j-1)/2\rfloor-1}(\beta_{2j-2}))$. So we need to show that $I(\omega_1, \omega_2) \subset D_i$. So we consider the homeomorphism

$$\psi_{1,4\lfloor (j-1)/2\rfloor-1} = \psi_{1,4n_i-1} \circ \psi_{4n_i+1,4\lfloor (j-1)/2\rfloor-1}.$$

By the inductive hypothesis,

$$\psi_{4n_i+1,4\lfloor (j_1-1)/2\rfloor-1} = \text{ identity on } I(\beta_{2j_1-1}, \beta_{2j_1}) \text{ for all } 2n_i + 1 \leq j_1 < j.$$

So

$$\psi_{1,4\lfloor (j-1)/2\rfloor-1} = \psi_{1,4n_i-1} \text{ on } I(\beta_{2j_1-1}, \beta_{2j_1}) \text{ for all } 2n_i + 1 \leq j_1 < j.$$

By condition b), this will also be true for $j_1 = j$, provided $(\beta_{2j-1}, \beta_{2j})$ is not an inactive satellite. So now we need to consider the action of $\psi_{1,4n_i-1}^{-1}$ on the region bounded by

$$I(\psi_{1,4\lfloor (j-2)/2\rfloor -1}(\beta_{2j-3}), \psi_{1,4\lfloor (j-1)/2\rfloor -1}(\beta_{2j-2})).$$

We use 3.8. Note that

$$\psi_{1,4n_i-1}^{-1} = \psi_{4n_i-3,4n_i-1}^{-1} \circ \cdots \circ \psi_{1,3}^{-1}$$

Therefore, the homeomorphism $\psi_{1,4n_i-1}^{-1}$ is a composition of disc exchanges $\xi_1 \circ \cdots \circ \xi_n$, with supports of the form $C(v_i \leftrightarrow v_i', E_i')$. For some of these disc exchanges, one of the exchanged topological discs contains D_i. For the others, any component of the union which intersects D_i, and which also intersects some $I(\beta_{2j_1-1}, \beta_{2j_1})$ for some $j_1 < j$, is contained in D_i. This means that the first disc exchanges map $\psi_{1,2n_i-1}(D_i)$ to D_i and then after that the only relevant disc exchanges preserve D_i. By the condition on $I(\omega_1, \omega_2)$ in a), it follows that $I(\beta_{2j-1}, \beta_{2j}) \subset D_i$. To see that $I(\beta_{2j-1}, \beta_{2j}) \subset D_i$ if $(\beta_{2j-1}, \beta_{2j})$ is an inactive satellite, we use the other part of condition a). This time the first disc exchanges which are applied are all in the composition for $\psi_{1,2n_i-1}^{-1}$, and the second group are in the composition for $\psi_{1,4\lfloor (j-1)/2\rfloor -1}^{-1}$, and the second part of condition a) ensures that $I(\beta_{2j-1}, \beta_{2j}) \subset D_i$.

2. Suppose that (6.5.1.1) holds and (6.5.1.2) holds for some i and k. Then (6.5.1.2) also holds for k' replacing k, for any $k' \leq k$. Therefore we can proceed by induction, both on j and on k. We can assume that (6.5.1.1) and (6.5.1.2) hold, and that (6.5.1.3) holds for $k-1$ replacing k. Then for all j with $2n_{k-1} \leq j \leq i$ such that $(\beta_{2j-1}, \beta_{2j})$ is not an inactive satellite,

$$\psi_{1,4n_{k-1}-1,\gamma} I(\beta_{2j-1}(\gamma), \beta_{2j}(\gamma)) = \psi_{1,4\lfloor (j-1)/2\rfloor -1,\gamma}(I(\beta_{2j-1}(\gamma), \beta_{2j}(\gamma))$$

and these are mapped over

$$\psi_{1,4\lfloor (i-1)/2\rfloor -1,\zeta}(I(\beta_{2i-1}(\zeta), \beta_{2i}(\zeta)) \subset \psi_{1,4n_{k-1}-1,\gamma}(D_k).$$

It follows by induction on k that, for $4n_{k-1} \leq j \leq 4n_k$,

$$\beta_j(\zeta) = \beta_j(\gamma)$$

and for $n_{k-1} \leq j \leq n_k$,

$$\psi_{1,4j-1,\zeta} = \psi_{1,4j-1,\gamma},$$

which give (6.5.1.3) for j. The inductive step for inactive satellites relies on the fact that an inactive satellite $I(\beta_{2j-1}(\zeta), \beta_{2j}(\zeta))$ is obtained from the support of $\psi_{1,2\lfloor (j-1)/2\rfloor -1,\zeta}$ and similarly for γ replacing ζ. So once again, the proof is inductive.

3. By 2, (6.5.1.3) holds for the largest j with $2n_j \leq n(\gamma)$. So

$$\psi_{1,4n_j-1,\gamma} = \psi_{1,4n_j-1,\zeta} = \psi_{1,4i-1,\gamma} \text{ on } I \text{ for all } 2n_j \leq 2i \leq n(\beta).$$

For any $i > 2n_1$, and any $i' > n_1$, by induction on i, the set $I(\beta_{2i-1}(\zeta), \beta_{2i}(\zeta))$ and the support of $\psi_{4i'-3,4i'-1,\zeta}$, are both contained in

$$\cup_{n>0}(\sigma_{\beta^j} \circ s)^{-n}(E_j)$$

which is disjoint from

$$I = I(\beta_{2n(\gamma)-1}(\gamma), \beta_{2n(\gamma)}(\gamma)) = I(\beta_{2n(\zeta)-1}(\zeta), \beta_{2n(\zeta)}(\zeta)).$$

It follows that

$$\psi_{1,4\lfloor n(\gamma)/2\rfloor-1,\gamma}(I) = \psi_{4n_j-3,4n_j-1,\gamma}(I) = \psi_{1,4\lfloor n(\zeta)/2\rfloor-1,\gamma}(I)$$

and hence since one endpoint of $\psi_{1,4\lfloor n(\gamma)/2\rfloor-1}(I)$ is the endpoint of γ by the definition of $I(\beta_{2n(\gamma)-1}(\gamma), \beta_{2n(\gamma)}(\gamma))$ and the same endpoint is the endpoint of ζ by the corresponding definition for ζ, we have $\gamma = \zeta$. □

In order to use this lemma to prove Theorems 6.1 and 6.2, we need to know that the conditions of 6.4 can be satisfied. This is given by the following theorem.

THEOREM 6.5.2. 1. *It is possible to choose a set*

$$U \subset \{z : \mathrm{Im}(z) > 0\}$$

and an even $n_0 > 0$ such that the following hold for U and for the set $\mathcal{U}$ of capture paths in $\mathcal{Z}(3/7, +, +, 0)$ with endpoints in U.

(i) $\beta_j(\gamma) = \beta_j$ is constant for $\gamma \in \mathcal{Z}(3/7, +, +, 0)$ with $j \le 4n_0+2$, so that

$$D_0 = D(\beta_{4n_0+1}(\gamma), \beta_{4n_0+2}(\gamma)$$

is independent of $\gamma \in \mathcal{U}$.

(ii) If $\zeta \in \mathcal{Z}(3/7, +, 0)$ and $\sigma_\gamma \circ s$ and $\sigma_\zeta \circ s$ are Thurston equivalent for some $\gamma \in \mathcal{U}$ then $\zeta \in \mathcal{Z}(3/7, +, +, 0)$ or ζ crosses S^1 at $e^{2\pi it}$ for $t \in (\frac{19}{28}, \frac{39}{56})$. If $\zeta \in \mathcal{Z}(3/7, +, +, 0)$ and

$$D(\beta_{2n(\zeta)-1}(\zeta), \beta_{2n(\zeta)}(\zeta)) \subset D_0,$$

then $\zeta \in \mathcal{U}$, and consequently

$$\beta_i(\zeta) = \beta_i \text{ for } i \le 4n_0.$$

2. *Fix any integer $r \ge 1$. Then we can find sets $U_i \subset \{z : \mathrm{Im}(z) > 0\}$, for $1 \le i \le 2^{r+1}$, and even integers $n_{0,i}$ such that the following hold, given $d^0 > 1$, where $\mathcal{U}_i$ is the set of capture paths in $\mathcal{Z}(3/7, +, +, 0)$ which endpoints in U_i.*

   (i) For each $i \le 2^{r+1}$ and each $j \le 2n_{0,i}+2$, the map $\gamma \mapsto \beta_j(\gamma) = \beta_{j,i}$ is constant on $\mathcal{U}_i$.

(ii) *The map* $(i,\gamma)\mapsto(\beta_{2n_{0,i}+1}(\gamma),\beta_{2n_{0,i}+2}(\gamma))$ *is constant on* $\cup_{j=1}^{2^{r+1}}(\{j\}\times\mathcal{U}_j$ *Consequently*

$$(\beta_{0,1},\beta_{0,2})=(\beta_{4n_{0,i}+1}(\gamma),\beta_{4n_{0,i}+2}(\gamma))$$

is independent of i *and of* $\gamma\in\mathcal{U}_i$*, and therefore*

$$D_0=\psi^{-1}_{1,4n_{0,i}-1}(U_i)$$

is independent of i*.*

(iii) *If* $\zeta\in\mathcal{Z}(3/7,+,0)$ *is such that* $\sigma_\gamma\circ s$ *and* $\sigma_\zeta\circ s$ *are Thurston equivalent for some* $\gamma\in\cup_i\mathcal{U}_i$*, then* $\zeta\in\mathcal{Z}(3/7,+,+,0)$ *or* ζ *crosses* S^1 *at* $e^{2\pi it}$ *for* $t\in(\frac{19}{28},\frac{39}{56})$*. If* $\zeta\in\mathcal{Z}(3/7,+,+,0)$ *and* $\gamma\in\mathcal{U}_i$ *for some* $i\le 2^{r+1}$*, and*

$$D(\beta_{2n(\zeta)-1}(\zeta),\beta_{2n(\zeta)}(\zeta))\subset D_0,$$

then $\zeta\in\mathcal{U}_j$ *for some* $j\le 2^{r+1}$*, and consequently*

$$\beta_i(\zeta)=\beta_{i,j}\text{ for }i\le 4n_{0,i}.$$

In both cases we can choose U*, or the* U_i*, so that*

$$d^0\le\limsup_{m\to\infty}\frac{\#((D_0\setminus\cup_{n>0}s^{-n}(E_0))\cap Z_m}{\#(D_0\cap Z_m)},$$

where

$$E_0=\cup\{E_{\omega_1,\omega_3}:(\omega_1,\omega_2)=(\beta_{0,1},\beta_{0,2}).$$

If we can prove both this theorem and Theorem 6.4 then we have proved Theorems 6.1 and 6.2. Of course this theorem has two cases, the first of which is needed for 6.1, and the second for 6.2. Each of these two cases has two subcases, as we shall see. It seems likely that for $\gamma\in\mathcal{U}$ or $\gamma\in\mathcal{U}_i$, there is no capture path $\zeta\in\mathcal{Z}(3/7,+,-,0)$ such that $\sigma_\zeta\circ s$ and $\sigma_\gamma\circ s$ are Thurston equivalent. Indeed it seems likely that there is also no such $\zeta\in\mathcal{Z}(3/7,+,p)$ for any $p>0$. But it has not been possible to completely prove this because of incomplete knowledge of $R_{m,p}$ for $p\ge 1$: an issue which has also affected all our earlier analysis.

6.6. The choice of U for Theorem 6.1

We make a quite explicit example, with $n_0=1$. We choose β_1 and β_2 so that β_1 is the maximum path in $R_{m,0}$ with first unit-disc-crossing to the right of $D(u)$,where $u=(L_3L_2R_3)^2L_3^3L_2R_3L_3^4L_2$ and therefore β_2 is the minimum path in $R_{m,0}$ with first unit-disc-crossing to the left of $D(u)$. We then have $w_1'(\beta_3)=w_1'(\beta_4)=L_3$. Whatever the choice of $w_2'(\beta_3)$, there is a set $C(u_1\leftrightarrow u_2,E_{\beta_1,\beta_3})$ which intersects $D(\beta_1,\beta_2)$ without being contained in it, where

$$u_1=(L_3L_2R_3)^2L_3^3L_2R_3L_3^3,\ u_2=(L_3L_2R_3)^2L_3L_2BCL_1R_2R_3L_2R_3.$$

We choose

$$w_2'(\beta_3) = w_2'(\beta_4) = u_2 L_3 x L_3$$

for a suitable nontrivial x. Then E_{β_1,β_3} contains the subset of $D(L_3L_2R_3)$ which starts at the right-hand edge of $D(L_3)$ and ends at the first unit-disc-crossing of β_1 and β_2. It follows that the second unit-disc-crossing of β_3 and β_4 is in the interior of u_2E. Thus $(\beta_t : 1 \leq t \leq 4)$ is a type AC quadruple, in the language of section 4. In 4.9.1, it was shown that for any capture path γ with $\beta_i(\gamma) = \beta_i$ for $i \leq 4$, we have

$$I(\beta_{2i-1}(\gamma), \beta_{2i}(\gamma)) \subset \cup_{n \geq 0} u_2^n D(\beta_1, \beta_2).$$

In fact we will make conditions that ensure that, for all $i \geq 3$,

$$I(\beta_{2i-1}(\gamma), \beta_{2i}(\gamma)) \subset D(\beta_3, \beta_4)$$

To start the process, we choose

$$D(\beta_5, \beta_6) \subset E_{1,3} \cap D(\beta_3, \beta_4) \setminus \cap_{n>o} s^{-n}(E_{1,3} \cup D(\beta_1, \beta_2)),$$

so that

$$\psi_{1,3} \mid D(\beta_5, \beta_6) = \xi \mid D(\beta_5, \beta_6),$$

where ξ is the single disc exchange with support $C(u_1 \leftrightarrow u_2, E_{1,3})$.

Since $n_0 = 1$, we have $D_0 = D(\beta_5, \beta_6)$, and E_0 is the union of all quadruples $(\omega_t : 1 \leq t \leq 4)$ with $(\omega_1, \omega_2) = (\beta_5, \beta_6)$. It is clear that by choice of (β_5, β_6), we can ensure that

$$\psi_{1,3}(D(\beta_5, \beta_6)) \subset \{z : \text{Im}(z) > 0\},$$

for example by choosing

$$D(\beta_5, \beta_6) \subset (u_2 u_3 X),$$

where $X = BC$ or UC, and u_3 has an even number of letters in $\{L_3, L_2\}$ if $X = BC$, and an odd number if $X = UC$. This is because u_1 has an odd number of letters in $\{L_3, L_2\}$. Also, we can choose (β_5, β_6) so that

$$\limsup_{m \to \infty} \frac{\#(\cup_{n>0} s^{-n}(E_0) \cap Z_m)}{\#(D_0 \cap Z_m)}$$

is arbitrarily close to 1. Now we choose

$$U = \psi_{\beta_1,\beta_3}(D(\beta_5, \beta_6)),$$

and choose $\mathcal{U}$ to be the set of capture paths in $\mathcal{Z}(3/7, +, +, 0)$ with endpoints in U. Then $\beta_i = \beta_i(\gamma)$ for all $\gamma \in \mathcal{U}$, and $\psi_{1,3,\gamma} = \psi_{\beta_1,\beta_3}$ for all $\gamma \in \mathcal{U}$. Therefore the conditions at the beginning of 6.4 are satisfied. For such a (β_5, β_6), we can take d^0 arbitrarily close to 1. Therefore, the condition at the end of 6.4 is satisfied.

Now we need to prove Theorem 6.5.2. So we need to show that any $\zeta \in \mathcal{Z}(3/7,+,0)$ such that $D(\beta_{2n(\zeta)-1}(\zeta),\beta_{2n(\zeta)}(\zeta)) \subset D_0$, we have $\zeta \in \mathcal{U}$. We have

$$\mathcal{Z}(3/7,+,0) = \mathcal{Z}(3/7,+,+,0) \cup \mathcal{Z}(3/7,+,-,0).$$

First we deal with the case of $\mathcal{Z}(3/7,+,+,0)$. We will deal with the case of $\mathcal{Z}(3/7,+,-,0)$ in 6.8.

LEMMA 6.6.1. *If* $\zeta \in \mathcal{Z}(3/7,+,+,0)$ *and* $D(\beta_{2n(\zeta)-1}(\zeta),\beta_{2n(\zeta)}(\zeta)) \subset D_0$, *then* $\beta_i(\zeta) = \beta_i(\gamma)$ *for* $i \leq 4$ *for any* $\gamma \in \mathcal{U}$.

PROOF. By the conditions on $\beta_i(\gamma)$ for $i \leq 6$ and $\gamma \in \mathcal{U}$, we see that $(\beta_i(\gamma) : 1 \leq i \leq 4)$ is a quadruple of type AC. By 4.12.2, and the choice of $D_0 = D(\beta_5(\gamma),\beta_6(\gamma)$, we have $I(\beta_{2i-1}(\gamma),\beta_{2i}(\gamma)) \subset D(\beta_3(\gamma),\beta_4(\gamma)$ for all $i \geq 2$ and all $\gamma \in \mathcal{U}$. Also by the choice of γ and by 4.12.1, 4.12.2 and 4.12.3, $D(\beta_{2i-1}(\gamma),\eta_{2i}(\gamma)$ does not intersect any of the sets of the form

$$D(\beta_1(\zeta),\beta_2(\zeta)) \cap E_{1,3,\zeta} \cup \cup_{n\geq 1} u^n E_{1,3,\zeta}$$

for any $\zeta \in \mathcal{Z}(3/7,+,+,0)$ such that $(\beta_i(\zeta) : 1 \leq 4)$ is different from $(\beta_i(\gamma) : 1 \leq i \leq 4)$. So if $D(\beta_{2n(\zeta)-1}(\zeta),\beta_{2n(\zeta)}(\zeta)) \subset D_0$, we must have $\beta_i(\gamma) = \beta_i(\zeta)$ for $i \leq 4$. ☐

6.7. The choice of U for Theorem 6.2

We use the examples constructed in Chapter 5. So fix $r \geq 1$. We consider $\gamma^{r,\alpha}$ for $\alpha \in \{a,b\}^{\mathbb{N}}$. The definition of $\gamma^{r,\alpha}$ depends only on $\alpha(k)$ for $0 \leq k \leq r$. For any $\alpha \in \{a,b\}^{\mathbb{N}}$, let $n(k,\alpha)$ be as in 5.4.3, that is, $n(k,\alpha)$ is the largest (automatically even) integer $2i$ such that

$$I(\psi_{1,4i-1,\alpha} \cdot (\beta_{4i-1}(\alpha)),\beta_{4i}(\alpha)) \not\subset D(v_{k,\alpha})$$

if $1 \leq k \leq r$, and $n(r+1,\alpha)$ is the largest integer i such that $(\beta_{2i-1}(\alpha),\beta_{2i}(\alpha))$ is defined.

We have seen that the branched coverings $\sigma_{\gamma^{r,\alpha}} \circ s$ are all Thurston equivalent for all $\alpha \in \{a,b\}^{\mathbb{N}}$, that is,

$$\beta_{2n(r+1,\alpha)}(\gamma^{r,\alpha})) = \beta_{2n(r+1,\alpha^\infty)}(\gamma^{r,\alpha^\infty}) \text{ for all } \alpha,$$

where α^∞ is defined by $\alpha^\infty(k) = a$ for all $k \in \mathbb{N}$. But otherwise the sequences $(\beta_i(\gamma^{r,\alpha}))$ and $(\beta_i(\gamma^{r,\alpha^\infty}))$ are quite different. Define

$$\Delta_{i,\alpha} = D(\beta_{2n(i,\alpha)-1}(\gamma^{r,\alpha}),\beta_{2n(i,\alpha)}(\gamma^{r,\alpha}))$$

for $i \leq r$ and

$$\Delta_{r+1,\alpha} = D(\beta_{2n(r+1,\alpha)-3}(\gamma^{r,\alpha}),\beta_{2n(r+1,\alpha)=2}(\gamma^{r,\alpha}))$$

$$\Delta'_{i,\alpha} = \psi_{1,2n(i,\alpha)-5,\gamma^{r,\alpha}}(\Delta_{i,\alpha})$$

for $i \leq r$ and similarly for $i = r+1$. Although there is no reason why this should be true in general, we have seen that $\Delta'_{i,\alpha} = D(\omega_1, \omega_2)$, for some adjacent pair (ω_1, ω_2) in $R_{m,0}$ (depending on i and α).

Let $\zeta_i(x)$ be the sequence defined in 4.4 for some words x. In particular, $\zeta_i(x)$ defined for all i if $x = x_{r,\alpha}$ with $\alpha(0) = a$. Let α^* be as in 5.4.5. We define

$$n_1(i,*) = n_1(i,\alpha^*), \quad x^* = x_{r,\alpha^*}, \quad \Delta_{i,*} = \Delta_{i,\alpha^*},$$

where the first two of these definitions are as in 5.3.4 and 5.4.4 respectively. For $1 \leq k \leq r$, $n(k,\alpha)$, as given above, is also the largest even integer $2i$ such that

$$D(\beta_{4i-1}(\alpha), \beta_{4i})(\alpha)) \not\subset D(v_{k-1,*}t_{k-1,*}a) \cup D(v_{k-1,*}t_{k-1,*}b).$$

Define $\alpha^{*,i}$ by

$$\alpha^{*,i}(j) = \alpha^*(j) \text{ if } j \neq i, \quad \alpha^{*,i}(i) \neq \alpha^*(i),$$

and define

$$x^{*,i} = x_{r,\alpha^{*,i}}, \quad \Delta_{i,i,*} = \Delta_{i,\alpha^{*,i}}.$$

Then

$$n_1(i, \alpha^{*,i}) = n_1(i,*).$$

We saw in Chapter 5, in particular in 5.4.4 and 5.4.5, that for any $\alpha \in \{a,b\}^{\mathbb{N}}$,

$$\begin{aligned} &(\beta_{2n(i,\alpha)-1}(x_{r,\alpha}), \beta_{2n(i,\alpha)}(x_{r,\alpha})) = (\zeta_{2n_1(i,*)}(x^*), \zeta_{2n_1(i,*)}(x^*)) \\ &\text{ or } \text{ if } (\zeta_{2n_1(*)-1}(x^{*,i}), \zeta_{2n_1(i,*)}(x^{*,i}))), \end{aligned} \tag{6.7.0.1}$$

but it is not always clear which of these is true. We have

$$I(\beta_{2n(\gamma^{r,\alpha})-1}(\gamma^{r,\alpha}), \beta_{2n(\gamma^{r,\alpha})}(\gamma^{r,\alpha})) \subset D(v_0t_0a) = D(v_1)$$

for all $\alpha \in \{a,b\}^{\mathbb{N}}$. Now $x_{r,\alpha}$ consists only of the letters L_3, L_2 and R_3, apart from the last letter C. The number of letters of $x_{r,\alpha}$ in $\{L_3, L_2\}$ is odd. The same are true for x^*. Therefore, writing $x_{r,\alpha} = yC$, and $x^* = y^*C$, we have

$$D(yBC) \cup D(y^*BC) \subset \{z : \mathrm{Im}(z) > 0\}.$$

For any capture path $\gamma \in \mathcal{Z}(3/7,+,+,0)$ with endpoint in $D(yBC)$, it is implicit in the results of Section 5 that

$$\beta_i(\gamma) = \beta_i(\gamma^{r,\alpha}) = \beta_i(\alpha) \text{ for } i \leq 2n(\gamma^{r,\alpha}) - 2,$$

and hence, for $n = n(\gamma^{r,\alpha}) - 1$,

$$D(\beta_{2n-1}(\gamma), \beta_{2n}(\gamma) = Dr+1,*.$$

Write $x^* = y^*C$. Our candidate for the set U is a set of the form

$$D(\omega_1, \omega_2) = \psi_{1,2n(r+1,\alpha)-3}(D(\eta_1, \eta_2)),$$

where $D(\eta_1, \eta_2) \subset D(y^*BC)$ and $D(\omega_1, \omega_2) \subset D(yBC)$. We choose such a set U with

$$\psi^{-1}_{1,2n(r+1,\alpha)-1}(U) \subset \{z : \mathrm{Im}(z) > 0\}.$$

It is certainly possible to choose U so that this is true. We can also choose U so that $d^0 = d^0(U)$ arbitrarily close to 1, by choosing U so that

$$U \subset D(yBCL_1R_1^N)$$

for an arbitrarily large N. We will then have

$$D_0 = D(\beta_{4n_0+1}, \beta_{4n_0+1}) \subset D_{r+1,*} = D(\beta_{4n_0-1}, \beta_{4n_0}).$$

It remains to prove Theorem 6.5.2 for this choice of U. So as usual we let $\mathcal{U}$ be the set of capture paths in $\mathcal{Z}(3/7, +, +, 0)$ with endpoints in U. We write $4n_0 = 2n(r + 1, \alpha) - 2$. Suppose that $\zeta \in \mathcal{Z}(3/7, +, 0)$ and

$$D(\beta_{2n(\zeta)-1}(\zeta), \beta_{2n(\zeta)}(\zeta)) \subset D_0 = \Delta_{r+1,*}. \tag{6.7.0.2}$$

We want to show that there is some $\theta \in \{a, b\}^{\mathbb{N}}$ such that

$$\beta_i(\zeta) = \beta_i(\gamma^{r,\theta}) \text{ for } i \leq 2n(r + 1, \theta) - 2.$$

Once again, we consider the case of $\zeta \in \mathcal{Z}(3/7, +, +, 0)$ first, and we will deal with $\zeta \in \mathcal{Z}(3/7, +, -, 0)$ in 6.8.

Theorem 6.7.1. *Let $\mathcal{U}$ be as above. Let $\gamma \in \mathcal{U}$ and $\zeta \in \mathcal{Z}(3/7, +, +, 0)$ such that (6.7.0.2) holds. Then there is α and $2n_0 = n(r + 1, \theta)$ such that*

$$\beta_i(\zeta) = \beta_i(\gamma) = \beta_i(\gamma^{r,\theta}) \text{ for } i \leq 4n_0.$$

Proof. The proof, as usual, is an induction.We shall show the following.

A-1 If $r \geq 1$, then $p(\zeta) \in D(v_0t_0) \cup D(w_0t_0)$ where, as usual, $p(\zeta)$ denotes the endpoint of ζ and $\beta_i(\zeta) = \beta_i(\gamma^{r,\theta})$ for $1 \leq i \leq 4$, for some $\theta \in \{a, b\}^{\mathbb{N}}$. Moreover

$$D(\beta_{2j-1}(\zeta), \beta_{2j}(\zeta)) \subset D(\beta_3(\zeta), \beta_4(\zeta))$$

for all $j \geq 2$.

A-2 For each i with $1 \leq i \leq r$ there exists $\theta \in \{a, b\}^{\mathbb{N}}$ such that $p(\zeta) \in D(v_{i,\theta})$ and $\beta_j(\zeta) = \beta_j(\theta)$ for $j \leq n(i, \theta)$. Moreover

$$D(\beta_{2j-1}(\zeta), \beta_{2j}(\zeta)) \subset D(v_{i-1,*}t_{i-1,*}a) \cup D(v_{i-1,*}t_{i-1,*}b)$$

for all $j > n(i, \theta)$.

A-3 $p(\zeta) \in D(v_{r,\theta}v_{r-1,\theta}t_{r-1,\theta})$ and $\beta_j(\zeta) = \beta_j(\gamma^{r,\theta})$ for all $j \leq n(\gamma^{r,\theta}) - 2$.

To prove A-1, we use 4.9.1, and 4.12.1 to 4.12.3, which give information about the sequence $\beta_i(\zeta)$. By (6.7.0.2), we need

$$\Delta_{r+1,*} \subset (D(\beta_1,\beta_2) \cap E_{1,3}) \cup \cup_{n\geq 1} u^n E_{1,3}.$$

where $u \leftrightarrow u'$ is a basic exchange for β_1 of the type in (3.9.1.4). Since $\Delta_{r+1,*} \subset D(v_0t_0)$ if $r \geq 1$ and $\Delta_{r+1,*} \subset D(v_0L_3L_2R_3L_3^3)$ if $r = 0$, this gives just two choices for (β_1,β_2): namely, β_1 is maximal with $w_1'(\beta_1)$ to the right of $D(v_0L_3L_2R_3L_3^3L_2)$ or $D(w_0L_3L_2R_3L_3^3L_2)$, that is,

$$(\beta_1(\zeta),\beta_2(\zeta)) = (\beta_1(\theta),\beta_2(\theta))$$

for some $\theta \in \{a,b\}^{\mathbb{N}}$. Of course, as a result of this, $(\beta_1(\zeta),\beta_2(\zeta))$ only depends on $\theta(0)$, but then we can also choose θ so that

$$(\beta_3(\zeta),\beta_4(\zeta)) = (\beta_3(\gamma^{r,\theta}),\beta_4(\gamma^{r,\theta})).$$

By 4.12.1 and 4.12.3, and the choice of $\beta_i(\zeta)$ for $i \leq 4$, we then have

$$D(\beta_{2j-1}(\zeta),\beta_{2j}(\zeta)) \subset D(L_3L_2L_3L_3) \setminus D((L_3L_2R_3)^2L_3)$$

for all $j \geq 1$. We now claim that we have

$$D(\beta_{2j-1}(\zeta),\beta_{2j}(\zeta)) \subset D(\beta_3(\zeta),\beta_4(\zeta)) \subset D(v_0t_0)$$

for all $j \geq 2$. If this is not true for a least j, then $j = 2j_0+1$ is odd by 4.10.2, and there is an exchange $u_1 \leftrightarrow u_1'$ for β_3 (and for β_{j_1} for $3 \leq j_1 < 4j_0$) such that

$$u_1(E_{1,3}) \cup D(\beta_3,\beta_4)) \subset D(\beta_3,\beta_4)$$

Then u_1 does not exist if $r = 0$ and v_0t_0a or v_0t_0b must be a proper prefix of u_1 if $r \geq 1$. But then the same v_0t_0a or v_0t_0b is also a prefix of $u_1'$and from considering sets $C(u_2 \leftrightarrow u_2', E')$ which can intersect $C(u_1 \leftrightarrow u_1', E)$, for E and $E' \subset D(L_3L_2L_3)L_3) \setminus D(L_3L_2R_3)^2L_3)$, we see that the component of the support of $\psi_{5,4j_0-1,\zeta}$ which intersects $D(\beta_{4j_0+1}(\zeta),\beta_{4j_0+2}(\zeta))$ is contained in $D(u_1) \cup D(u_1')$. The same is true for any set $C(u_2 \leftrightarrow u_2', E_{1,3})$ which intersects $D(\beta_{4j_0+1}(\zeta),\beta_{4j_0+2}(\zeta))$, apart from $C(v_0 \leftrightarrow w_0, E_{1,3})$ which contains it. It follows that

$$D(\beta_{4j_0+1}(\zeta),\beta_{4j_0+2}(\zeta)) \subset D(u_1) \cup D(u_1') \subset D(\beta_3,\beta_4).$$

Hence we have

$$D(\beta_{2j-1}(\zeta),\beta_{2j}(\zeta)) \subset D(\beta_3(\zeta),\beta_4(\zeta)) \text{ for all } j \geq 2.$$

This completes the proof of [A-1]. So now we consider [A-2]. By a similar argument, if $C(u_1 \leftrightarrow u_1', E)$ is the first exchange to intersect $D(\beta_{2j-1}(\zeta),\beta_{2j}(\zeta))$ for a minimal j, where v_0t_0a or v_0t_0b is a proper prefix of u_1 then we have

$$D(\beta_{2j-1}(\zeta),\beta_{2j}(\zeta)) \subset D(u_1) \cup D(u_1').$$

Then by induction,

$$D(\beta_{2\ell-1}(\zeta), \beta_{2\ell}(\zeta)) \subset D(u_1) \cup D(u_1') \text{ for all } \ell \geq j. \tag{6.7.1.1}$$

By (6.7.0.2) this can only be true for u_1 a prefix of $x_{r,*}$. Also we are assuming that $v_0t_0a = v_1$ or v_0t_0b is a proper prefix of any such u_1. The first such prefix of $x_{r,*}$ is $u_1 = v_1t_1a$ or v_1t_1b (since $v_{1,*} = v_1$ and $t_{1,*} = t_1$), that is, $u_1 \leftrightarrow u_1'$ is $v_{2,*} \leftrightarrow v_1t_1b$ (interchanging u_1 and u_1' if necessary). It follows that for all sufficiently large ℓ,

$$D(\beta_{2\ell-1}(\zeta), \beta_{2\ell}(\zeta)) \subset D(v_1t_1a) \cup D(v_1t_b). \tag{6.7.1.2}$$

It follows by induction on $j \geq 4$ that that there is θ such that

$$\beta_j(\zeta) = \beta_j(\theta) \text{ for } j \leq n(1, \theta),$$

and (6.7.1.2) holds for all $\ell \geq n(1, \theta)$. This completes the proof of [A-2] for $i = 1$. The proof of [A-2] for a general i is similar. We assume inductively that [A-2] is proved for $i - 1$. Then, as before, we can prove that

$$D(\beta_{2j-1}(\zeta), \beta_{2j}(\zeta)) \subset D(v_{i-2,*}t_{i-2,*}a) \cup D(v_{i-2,*}t_{i-2,*}b)$$

for all $j \geq n(i-1, \theta)$. For if this is not true then as before we can find $2j_0 > n(i, \theta)$ an exchange $u_1 \leftrightarrow u_1'$ for β_j for all j with $2n(i-1, \theta) < j \leq 4j_0$ such that $v_{i-2,*}t_{i-2,*}e$ is a proper prefix of u_1 for one of $e = a$ or $e = b$, and such that

$$D(\beta_{2j-1}(\zeta), \beta_{2j}(\zeta)) \subset D(u_1) \cup D(u_1') \text{ for all } j > 2j_0.$$

This is only possible if $u_1 \leftrightarrow u_1'$ is one of the exchanges for which this happens for $(\beta_{2j-1}(\theta), \beta_{2j}(\theta))$, which means end in $v_{k-1,*}t_{k-1,*}e$ for some $k \leq i - 1$, or in $t_{i-1,*}e$, for $e \in \{a, b\}$. So then by induction on j we can define $\theta(i) \in \{a, b\}$ so that

$$\beta_j(\zeta) = \beta_j(\theta) \text{ for all } j \leq 2n(i, \theta),$$

and the rest of [A-2] for i is proved as before.

The proof of [A-3] is similar.

□

6.8. Paths in $\mathcal{Z}(3/7+, -, 0)$

In order to complete the proof of 6.5.1) and 6.5.2, it suffices to show that for any capture paths $\gamma \in \mathcal{U}$ and $\zeta \in \mathcal{Z}(3/7, +, -, 0)$, the maps $\sigma_\gamma \circ s$ and $\sigma_\zeta \circ s$ are not Thurston equivalent. In the process, we shall obtain some restrictions on the set of paths $\beta \in R_{m,0}$ such that $\sigma_\beta \circ s$ is equivalent to $\sigma_\zeta \circ s$, for $\zeta \in \mathcal{Z}(3/7, 0, +, -, 0)$. These restrictions appear to be of interest in their own right. More specifically, we shall prove the following.

THEOREM 6.8.1. *Let* $\zeta \in \mathcal{Z}(3/7,+,-,0)$ *which crosses* S^1 *at* $e^{2\pi it}$ *for* $t \in [\frac{39}{56},\frac{5}{7})$ *and* $\beta \in R_{m,0}$ *such that* $\sigma_\zeta \circ s$ *and* $\sigma_\beta \in s$ *are Thurston equivalent. Then*

$$w_1'(\beta) = L_3 \text{ or } L_3L_2R_3L_2u \text{ or } L_3L_2UCu \text{ or } L_3L_2R_3L_3^2u$$

for some u*, and if* $w_1'(\beta) = L_3$ *then*

$$w_2'(\beta) = L_3^4u \text{ or } BCu \text{ or } L_3(L_2R_3)^2u$$

for some u*.*

For the sets $\mathcal{U}$ that we have chosen, for theorem 6.1 (chosen in 6.6) we have

$$w_1'(\beta_{2n(\gamma)}(\gamma)) = (L_3L_2R_3)^2L_3^3$$

for all $\gamma \in \mathcal{U}$ and for the set chosen for Theorem 6.2 (chosen in 6.7) we have

$$w_1'(\beta_{2n(\gamma)}(\gamma)) = L_3, \quad w_2'(\beta_{2n(\gamma)}(\gamma)) = v_0t_0a = L_3L_2R_3L_3(L_2R_3)^2u$$

for some u. It follows that $\sigma_\gamma \circ s$ is not equivalent to $\sigma_\zeta \circ s$ for any $\zeta \in \mathcal{Z}(3/7+,-,0)$ which crosses S^1 at $e^{2\pi it}$ for $t \in [\frac{39}{56},\frac{5}{7})$.

PROOF. We shall choose a suitable ψ and α for which

$$(s, Y_m) \simeq_\psi (\sigma_\alpha \circ s, Y_m),$$

where $\simeq$ denotes Thurston equivalence. We shall then consider

$$\zeta^1 = \alpha * \psi(\zeta).$$

The basic reason for considering such a ζ^1 was explained in 2.4: $\sigma_\zeta \circ s$ and $\sigma_{\zeta^1} \circ s$ are Thurston equivalent, and this is realised by composing the above equivalence on the left by σ_ζ. The aim is to choose ψ and α so that the first two unit-disc crossings of ζ^1 coincide with those of a path in $R_{m,0}$. If that is so, then we shall show that the path $\beta \in R_{m,0}$ such that $\sigma_\beta \circ s$ is equivalent to $\sigma_\zeta \circ s$ has the same first unit disc crossing as ζ^1, and the same second unit -disc crossing if $w_1'(\beta) = w_1'(\zeta^1) = L_3$.

Let ζ cross the unit circle at $e^{2\pi it}$ for $t \in (\frac{19}{28},\frac{5}{7})$. Then $t \in (\frac{39}{56},\frac{5}{7})$ or $t \in (\frac{19}{28},\frac{39}{56})$. We can discount ζ with $t = \frac{39}{56}$ because this point is in the boundary of the gap containing the point of preperiod 2 labelled by L_3L_2C, and it is then clear that $\sigma_\zeta \circ s$ is not equivalent to $\sigma_\gamma \circ s$ for any $\gamma \in \mathcal{U}$, because the endpoints of paths in $\mathcal{U}$ are in U, and all points in U have preperiod higher than 2, in fact, at least 11.

First suppose that $t \in (\frac{39}{56},\frac{5}{7})$. Let $\psi_{m,2/7}$ and $\alpha_{m,2/7}$ be as in 3.12 (and as in 3.3 and 7.2 of [**22**]), that is, these homeomorphisms and closed loops arise from considering the equivalence between the type II captures $\sigma_{\zeta_{5/7}}^{-1} \circ \sigma_{\beta_{5/7}} \circ s$ and $\sigma_{\zeta_{2/7}}^{-1} \circ \sigma_{\beta_{2/7}} \circ s$. Our initial choices of ψ and α are:

$$\psi = \psi_{m,2/7}$$

and

$$\alpha = \alpha_{m,2/7}.$$

This is just a first choice, which will need to be modified later.

Now let $t \in (\frac{19}{28}, \frac{39}{56})$. We use the notation of 3.13 (and of 3.3 of [**23**] – see the last paragraph of that section). Let $q = \frac{17}{56}$. This time our initial choices of ψ and α are:

$$\psi = \psi_{m,q},$$

$$\alpha = \alpha_{m,q}.$$

Once again, this is just a first choice, which will need to be modified later.

First, we consider the case $t \in (\frac{39}{56}, \frac{5}{7})$ and $\psi = \psi_{m,2/7}$. Referring to the definitions in 3.12, we see that $\psi_{2,2/7}$ is the identity on $\beta_{5/7}$, and hence $\alpha_{2/2/7}$ is an arbitrarily small perturbation of $\beta_{2/7} * \overline{\beta_{5/7}}$. For any path ζ based at ∞ we also have

$$\alpha_{m,2/7} * \psi_{m,2/7}(\zeta) = \beta_{2/7} * \psi_{m,2/7}(\overline{\beta_{5/7}} * \zeta).$$

For $\zeta \in \mathcal{Z}(3/7, +, 0)$, the homeomorphism $\psi_{2,2/7}$ is the identity on $\overline{\beta_{5/7}} * \zeta$ and hence for all $m \geq 3$,

$$\alpha_{m,2/7} * \psi_{m,2/7}(\zeta) = \beta_{2/7} * \xi_{m-1,2/7} \circ \psi_{m-1,2/7}(\overline{\beta_{5/7}} * \zeta).$$

Referring again to 3.12, we see that the support of $\xi_{2,2/7}$ intersects $\psi_{2,2/7}(\overline{\beta_{5/7}} * \zeta)$ with intersection contained in $D(BC) \cap C_{2,2/7}$. Since $\xi_{2,2/7}$ is an anticlockwise twist in the annulus $C_{2,2/7}$, this implies that $\psi_{3,2/7}(\overline{\beta_{5/7}} * \zeta)$ intersects the unit circle in the boundary of the central gap at $e^{2\pi i(11/14)}$ rather than $e^{2\pi i(5/7)}$, and the intersection with the unit disc nearer to the endpoint of ζ is unchanged. The support of $\xi_{3,2/7}$ does not intersect $\psi_{3,2/7}(\overline{\beta_{5/7}} * \zeta)$. Therefore

$$\psi_{3,2/7}(\overline{\beta_{5/7}} * \zeta) = \psi_{4,2/7}(\overline{\beta_{5/7}} * \zeta)$$

Referring to the description of $\xi_{4,2/7}$ restricted to $C_{4,2/7}$ in 3.12, we see that there are no intersections between $C_{4,/7}$ and $\psi_{4,2/7}(\overline{\beta_{5/7}} * \zeta)$ because of the restriction that $t > \frac{39}{56}$. However, there is an intersection with $A_{4,2/7}$, if and only if ζ crosses the unit circle between $e^{2\pi i(79/112)}$ and $e^{2\pi i(5/7)}$, that is, if and only if the endpoint of ζ is in $D(L_3L_2R_3L_3)$. For the moment we assume that the endpoint of ζ is in $D(L_3L_2R_3L_3)$. Then, because $\xi_{4,2/7}(D(L_3L_2R_3L_3)) = D(L_3^4)$, with the movement being along an arc with endpoints in the lower unit circle, we see that $\psi_{5,2/7}(\overline{\beta_{5/7}} * \zeta)$ intersects the unit disc in $D(L_3^4)$ rather than in $D(L_3L_2R_3L_3)$, and crosses the unit circle between $e^{2\pi i(37/56)}$ and $e^{2\pi i(75/112)}$ instead of between $e^{2\pi i(79/112)}$ and $e^{2\pi i(5/7)}$. There are actually two components of intersection with the unit disc in $D(L_3^4)$, because the action of $\xi_{4,2/7}$ pushes $\psi_{4,2/7}(\overline{\beta_{5/7}} * \zeta)$ up through

$D(L_3^4L_2)$, so that the last component of intersection of $\psi_{5,2/7}(\overline{\beta_{5/7}} * \zeta)$ enters $D(L_3^4L_2)$ from the top of the unit circle. There is no intersection between $\psi_{5,2/7}(\overline{\beta_{5/7}} * \zeta)$ and $C_{5,2/7}$. In fact there is never any intersection between $A_{i,2/7}$ and $C_{i+1,2/7}$, because there is no intersection between $A_{1,2/7}$ and $C_{2,2/7}$. There is, however an intersection between $\psi_{5,2/7}(\overline{\beta_{5/7}} * \zeta)$ and $A_{5,2/7}$ if and only if the endpoint of ζ is in $D(L_3L_2R_3L_3^2)$, that is, if and only if the endpoint of $\psi_{5,2/7}(\zeta)$ is in $D(L_3^5)$. In this case the action of $\xi_{5,2/7}$, with $\xi_{5,2/7}(D(L_3^5)) = D(BCL_1R_2R_3L_3)$, gives that the endpoint of $\psi_{6,2/7}(\zeta)$ is in $D(BCL_1R_2R_3L_3)$. There is then no intersection between $\psi_{6,2/7}(\overline{\beta_{5/7}}*\zeta)$ and $C_{6,2/7}$, and there is an intersection between $\psi_{6,2/7}(\overline{\beta_{5/7}}*\zeta)$ and $A_{6,2/7}$ if and only if the endpoint of ζ is in $D(L_3L_2R_3L_3^3)$, that is, if and only if the endpoint of $\psi_{6,2/7}(\zeta)$ is in $D(BCL_1R_2R_3L_3^2)$. Using the move $\xi_{6,2/7}(D(BCL_1R_2R_3L_3^2)) = D(L_3(L_2R_3)^2L_3)$, the endpoint of $\psi_{7,2/7}(\zeta)$ is in $D(L_3(L_2R_3)^2L_3)$. Continuing in this way, the endpoint of $\psi_{m,2/7}(\zeta)$ is in $D(u)$, where

$$u = u_1L_2 = L_3^4L_2(R_3L_3L_2)^r, \text{ or } BCL_1R_2R_3L_3L_2(R_3L_3L_2)^r, \text{ or } L_3(L_2R_3)^2L_3L_2(R_3L_3L_2)^r, \tag{6.8.1.1}$$

if $m = 5 + 3r$ or $6 + 3r$ or $7 + 3r$ respectively, for any $r \geq 0$. The action of $\xi_{i,2/7}$ restricted to components of $C_{i,2/7}$ is relevant for $i \geq 8$. Note that, for the first two choices of u, for any extension uv of u, we have $w_1'(uv) = L_3$, and u_1 is a prefix of $w_2'(uv)$. This is not true for the third choice of u.

The path $\zeta^1 = \alpha_{m,2/7} * \psi_{m,2/7}(\zeta)$ obtained in this way has first unit-disc crossing along $D((L_3L_2R_3)^\infty)$, but then departs from all paths $\beta \in R_{m,0}$ with $w_1'(\beta) = L_3$, because of the action of $\xi_{2,2/7}$ restricted to $C_{2,2/7}$ on $\overline{\beta_{5/7}} * \zeta$. In order to determine the next iteration ζ^2 towards a path in $R_{m,0}$, we consider the word $w = w(\zeta^1) = w(\psi_{m,2/7}(\zeta))$ representing the endpoint of $\psi_{m,2/7}(\zeta)$. If L_3^4 or L_3^2BC or BC is a prefix of w, when we define

$$\zeta^2 = \alpha^1 * \psi^1(\zeta^1),$$

where

$$(s, Y_m) \simeq_{\psi^1} (\sigma_{\alpha^1} \circ s, Y_m),$$

and β^1 is the unique path in $R_{m,0}$ which shares an endpoint with ζ^1 and with $w_1'(\beta^1) = L_3$, and α^1 is an arbitrarily small perturbation of $\beta^1 * \overline{\zeta^1}$ such that the disc (or union of discs) bounded by α^1 does not contain the common endpoint of β^1 and ζ^1. Therefore,

$$\zeta^2 = \beta^1 * \overline{\zeta^1} * \psi^1(\zeta^1).$$

Since u_1 is a prefix of $w_2'(w(\zeta^1))$, all of β^1 beyond the first unit disc crossing is in $D(u)$. Therefore all of α^1, apart from the first and last unit-disc crossing along $D(L_3L_2R_3)^\infty)$, is in

$$D(u^4)\cup D(u^5)\cup D(u^6),$$

where the u^i are the different possibilities for u in (6.8.1.1) with $4\le m\le 6$, that is,

$$u^4=L_3^4L_2,\quad u^5=BCL_1R_2R_3L_3L_2,\quad u^3=L_3(L_2R_3)^2L_3L_2.$$

From this, we can estimate the size of the components of the support of ψ^1 which intersect ζ^1. The difference between ζ^1 and β^1 is contained in a small part of the supports of the homeomorphisms $\xi_{m,2/7}$ for $m\ge 4$, in fact, for each one, a single component of intersection with the unit disc together with a single adjacent component of intersection with the complement of the unit disc. The path α^1 has a first and last crossing of the unit disc along the leaf $D((L_3L_2R_3)^\infty)$ between $e^{\pm 2\pi i(2/7)}$. It follows that the only component of $\text{supp}(\psi^1)$ which is not contained in $D(u^m)$ (for $4\le m\le 6$) has one intersection with $D(u^m(R_3L_3L_2)^\infty)$. and then the later intersections are with $D(v^mv^nL_3)$ for $4\le n\le 6$, where v^i is obtained from u^i by deleting the last two letters L_3L_2. So this component of the support is disjoint from ζ^1. It follows that the path ζ^2 is in $D(u)$ from the second unit-disc crossing onwards.

The case when $u=L_3(L_2R_3)^2L_3L_2(R_3L_3L_2)^r$ is a bit more involved. In this case, because of the definition of w_1' (in 3.6), it is possible to have $w_1'(w)\ne L_3$. In order to see what happens, we need to look at the longest prefix u_1u_2 of w such that u_2 has all letters in $\{L_3,L_2,R_3\}$ and is of the form

$$u_2=(L_2R_3)^{r_1}L_2 \text{ for } r_1\ge 0,$$

or

$$u_2=(L_2R_3)^{r_1}L_3^{r_2} \text{ for } r_1\ne 1 \text{ and } r_2\ge 1.$$

In the first case, u_2 is followed in w by BC or UC, and in the second case it is followed in w by L_2. In the first case we do have $w_1'(w)=L_3$, we define α^1, ψ^1 and ζ^2 as above, and it does follow, much as in the previous cases, that ζ^2 is in $D(L_3(L_2R_3)^2L_3)\cup D(L_3L_2R_3L_2UC)$ from the second unit-disc crossing onwards. In the second case, we can also define α^1, ψ^1 and ζ^2 as above, in the cases when $w_1'(u_1u_2L_2)=L_3$, and we have the same bounds on ζ_2 from the second unit-disc crossing onwards. It remains to consider the cases when $w_1'(u_1u_2L_2)\ne L_3$. This happens precisely when r_1+r_2 is odd and ≤ 3. We have:

$$w_1'(u_1u_2L_2)=\begin{cases} L_3(L_2R_3)^2(L_3L_2R_3)^{r-1}L_3 \text{ if } r_1=0 \text{ and } r_2=1,\\ u_1L_2R_3L_3^{r_2-2} \text{ if } r_1=0 \text{ and } r_2>1,\\ u_1(L_2R_3)^{r_1-2} \text{ if } r_1\geq 2 \text{ and } r_2=1,\\ u_1(L_2R_3)^{r_1-1} \text{ if } r_1\geq 2 \text{ and } r_2=3,\\ u_1(L_2R_3)^{r_1}L_3^{r_2-3} \text{ if } r_1\geq 2 \text{ and } r_2\geq 4.\end{cases}$$

In all these cases, we have

$$w_1'(w)=w_1'(u_1u_2L_2)u_3$$

for a word u_3 which is often empty. In all these cases we consider the leaf $\ell=D(w_1'(u_1u_2L_2)(L_2R_3L_3)^\infty)$ of $L_{3/7}$. Exactly one endpoint $e^{2\pi ip}$ is the first intersection point on the unit circle of at least one path in $R_{m,0}$. We then define α^1 to be the anticlockwise closed loop based at $\infty=v_2$ which goes along ℓ and then back through the disc along the leaf from $e^{2\pi i(5/7)}$ to $e^{2\pi i(2/7)}$. We then define ψ^1 and ζ^2 as above. It is possible that $w_1'(\zeta^2)\neq w_1'(u_1u_2L_2)$, and if the first unit disc crossing of ζ^1 is in the support of ψ^1, in the support of a disc exchange in the composition for ψ^1 which has connecting arc along the top of the unit circle. But if that happens then both components of intersections with the unit disc are in $D(w_1'(u_1u_2L_2))$, and there is no movement out of this set by other disc exchanges in the composition. So the first unit disc-crossing of ζ^2 is in $D(w_1'(u_1u_2L_2))$, and the process can then be repeated.

Now we need to consider what happens for $t\in(\frac{39}{56},\frac{79}{112})$. The case of $t=\frac{79}{112}$ is not a problem because if ζ crosses the circle at $e^{2\pi i(79/112)}$ then the end of ζ is in the gap in $D(L_3L_2R_3L_2)$ which also intersects the unit circle at $e^{2\pi i(33/112)}$. There is a capture path $\beta\in R_{m,0}$ with $w(\beta)=L_3L_2R_3L_2C$. The intersection if β with the unit circle is of course with the upper unit circle. The intersection point is not $e^{2\pi i(33/112)}$ (as was noted in passing in Chapter 7 of [**22**]) but at the leftmost point $e^{2\pi i(67/224)}$. For this β we have $w_1'(\beta)=L_3(L_2R_3)^3L_3$, which satisfies the conditions of the theorem.

Otherwise for $t\in(\frac{39}{56},\frac{79}{112})$, in some contrast to what was done before, we define $w=w(\zeta)$ and $x=w_1'(w)$. Then $\ell=x(L_2R_3L_3)^\infty$ is a vertical leaf. Let $e^{2\pi ip}$ be the point of ℓ which is the starting point of at least one path in $R_{m,0}$. Our new choice of ψ and α — with $\zeta^1=\alpha*\psi(\zeta)$ — are

$$\psi=\psi_{m,p},$$

$$\alpha=\alpha_{m,p}.$$

The structure of $\psi_{m,p}$ is similar to that of $\psi_{m,2/7}$, as noted in 3.13 (and earlier, in 3.3 of [**22**]). In particular, $\psi_{i+1,p}=\xi_{i,p}\circ\psi_{i,p}$ for all $i\geq 1$, where the support of $\xi_{i,p}$ is $A_{i,p}\cup C_{i,p}$ and

$$A_{i,p}=(\sigma_{\beta_p}\circ s)^{1-i}(A_{1,p}),\quad C_{i,p}=(\sigma_{\beta_p}\circ s)^{1-i}(C_{1,p}),$$

using the notation of 3.13. The annuli of $A_{i,p}$ are thinner than the corresponding annuli of $A_{i,2/7}$ but the annuli of $C_{i,p}$ are thicker than the corresponding annuli of $C_{i,2/7}$. It is not accurate to say that the annuli "correspond, because it is not true (for example) that each annulus component of $A_{i,2/7}$ contains a unique annulus component of $A_{i,p}$. But this is true for components of $A_{i,2/7}\cap\{z:|z|\leq 1\}$ and components of $A_{i,p}\cap\{z:|z|\leq 1\}$ and even on some (or most) unions of components of intersection with, alternately, $\{z:|z|\leq 1\}$ and $\{z:|z|\geq 1\}$.

In many cases, $w(\zeta^1)=w(\psi_{m,p}(\zeta))\in U^x$. We consider this case first. Then as when $x=L_3$, ζ^1 is likely to diverge from all paths of $R_{m,0}$ after the first unit-disc-crossing. In this case, we define β^1 to be the unique path in $R_{m,0}$with $w_1'(\beta^1)=x$ and such that β^1 shares an end point of ζ^1 α^1 to be a perturbation of $\beta^1*\zeta^1$ which bounds a disc (or union of discs) disjoint from the common endpoint of β^1 and ζ^1. We then define

$$\zeta^2=\alpha^1*\psi^1(\zeta^1).$$

As in the case when $x=L_3$, the action of ψ^1 on ζ^1 does not cancel the first unit-disc crossing of α^1, and keeps the second unit-disc crossing in U^x. Therefore the first two unit-disc crossings of ζ^2 coincide with those of a path of $R_{m,0}$ with $w_1'=x$.

Now suppose that $w(\zeta^1)\notin U^x$. There are more possibilities than in the case $x=L_3$, when $w(\zeta^1)\notin U^{L_3}$ occurred in essentially just one way. The new possibilities come from the difference between the homeomorphisms $\psi_{i,2/7}$ and $\psi_{i,p}$, more precisely, in the difference between the annuli in $A_{i,2/7}$ and $C_{i,2/7}$, and the annuli in $A_{i,p}$ and $C_{i,p}$. For any $p>\frac{2}{7}$, the component of $C_{2,p}$ which intersects $D(C)$ also has intersection with interior with both $D(L_3)$ and $D(R_3)$. Write

$$x=x_1L_3^2(L_2R_3L_3)^n \text{ for some } n\geq 0,\quad x_2=x_1L_3,\quad x_2'=x_1L_2,$$

or

$$x=x_1R_3L_2R_3(L_3L_2R_3)^nL_3 \text{ for some } n\geq 0,$$

$$x_2=x_1(R_3L_2)^2R_3,\quad x_2'=x_1R_3L_2,$$

or

$$x=x_1R_3,\ x_2=xL_2R_3,\quad x_2'=x_1.$$

Note that

$$C_{2,2/7}\cap D(C)=C_{2,p}\cap D(C),$$

and

$$A_{4,2/7}\cap D(L_3)\subset A_{4,p}\cap D(L_3)\cup C_{2,p}\cap D(L_3))\cup L_3L_2(C_{2,p}\cap D(R_3)).$$

Hence the same is true if we prefix by x_2. There is an important difference between $C_{5,2/7}$ and $C_{5,p}$, which arises because $\beta_{2/7}$ crosses $C_{2,2/7}$, but β_p does not intersect $C_{2,p}$. This means that, if x_2' has length i, there is a component of $C_{i,p}$ which wraps round the unique point of $Z_i\cap D(x_2'C)$, and $\xi_{i,p}$ restricted to this component is simply a half twist round this annulus. Furthermore, for each $n>0$, there is a component of $C_{i+3n,p}$ within this component of $C_{i,p}$, such that $\xi_{i+3n,p}$ restricted to this component is a fractional $1/2^n$ twist. The sets $w(\zeta)$ for which $D(w(\zeta^1))\not\subset U^x$ are of three types:

1. $D(w(\zeta^1))\subset D(x_2'X)\subset x_2'C_{2,p}$, where $X=UC$ or BC, depending on whether x_2' has an even or odd number of letters in $\{L_3,L_2\}$. If this happens, we have $w(\zeta)\in D(x_2'X')$ where $X'=BC$ or UC respectively, so that $X'\neq X$.
2. $D(w(\zeta^1))\subset D(y_1u_2)$, where $y_1=x_2L_3(L_2R_3)^2L_3(L_2R_3L_3)^r$ for some maximal $r\geq 0$, and u_2 with all letters in $\{L_3,L_2,R_3\}$ ending in L_2 or L_3, of one of the two forms as in the case $x=L_3$, and $w_1'(y_1u_2X)\neq x$, where $X\in\{BC,UC\}$ or $X=L_2$, whichever is appropriate. In this case, it is always also true that $w_1'(y_1u_2X,x)\neq x$. Also, we have $D(w(\zeta))\subset D(x_2L_3L_2R_3L_3^{3r+3})$. But in that case, $x_2L_3L_2R_3L_3^3$ is a proper prefix of $x=w_1'(w(\zeta))$. This contradicts the forms given above for x_2 in terms of x. So, in fact, this case does not occur.
3. $D(w(\zeta^1))\subset x_2L_3^2(C_{2,p}\cap D(L_3))$. In this case, we have $D(w(\zeta))\in x_2L_3L_2(C_{2,p}\cap D(R_3))$, and $w_1'(w(\zeta^1))\neq w_1'(w(\zeta))$.

In cases 1 and 3, noting that case 2 does not occur, we define $v_1=w_1'(\zeta^1)=w_1'(\psi_{m,p}(\zeta))$ and let $\ell_1=D(v_1(L_2R_3L_3)^\infty)$. Let $e^{2\pi ip_1}$ be the unique endpoint of ℓ_1 which is the first S^1 intersection point of at least one path in $R_{m,0}$, and define

$$\psi^1=\psi_{m,p_1},\quad \alpha^1=\alpha_{m,p_1},\quad \zeta^2=\alpha^1*\psi^1(\zeta^1).$$

In case 1 above we have $w(\psi_{m,p_1}(\zeta^2))\subset D(x_2'X)$, because $D(x_2'X)\subset C_{2,p_1}$, and in case 3 we have $D(w(\zeta^2))\subset x_2L_3^2(C_{2,p_1}\cap D(L_3))$, because $e^{2\pi ip_1}$ is to the left of $e^{2\pi ip}$, and consequently $C_{2,p_1}\supset C_{2,p}$. If $w_1'(\zeta^2)=x^2$ then all of ζ^2, from the first intersection with S^1 onwards, is in U^{x^2}, but, as in the case $x^2=L_3$, the path ζ^2 diverges from all paths in $R_{m,0}$ after the first unit disc crossing, and we need to adjust by defining

$$\zeta^3=\alpha^2*\psi^2(\zeta^2),$$

where, similarly to before, α^2 is a perturbation of $\beta^2 * \overline{\zeta^2}$, where $\beta^2 \in R_{m,0}$ is a path with $w_1'(\beta^2) = x^2$ and the second unit disc crossing coincides with the second unit-disc-crossing of ζ^2. As before, all of ζ^3 also, from the first intersection with the unit disc onwards, is contained in U^{x^2}. If $w_1'(\zeta^2) \neq x^2$, then we define p_3 similarly to p_2, as the unique endpoint of $D(w_1'(\zeta^2)(L_2R_3L_3)^\infty)$ which is the first S^1 intersection point of at least one path in $R_{m,0}$, and then define α^3, ψ^3 and ζ^3 similarly to α^2, ψ^2 and ζ^2. Then we iterate this process.

We finish with a discussion of the case of $t \in (\frac{19}{28}, \frac{39}{56})$, where the proof cannot be completed simply because we have no comprehensive description of the paths in $R_{m,1}$. Recall that this time we are taking $\psi = \psi_{m,q}$ for $q = \frac{17}{56}$. As before, the support of $\xi_{m,q}$ is a union of annuli $A_{m,q} \cup C_{m,q}$. The annulus components of $C_{1,q}$ are larger than the annulus components of $C_{1,2/7}$ while the annulus $A_{1,q}$ is smaller than the annulus $A_{1,2/7}$. The first intersection between $\psi_{i,q}(\zeta)$ and $A_{i,q} \cup C_{i,q}$ is for $i = 4$, as ζ intersects the component of $C_{4,q}$ which surrounds the point in $s^{-2}(0)$ with code L_3L_2C. As the twist in components of $C_{4,q}$ is anticlockwise, and is a half-twist, we see that $\alpha_q * \psi_{5,q}(\zeta)$ has first intersection point with S^1 at a point $e^{2\pi i u}$ for $u \in (\frac{17}{56}, \frac{9}{28})$. As for $\zeta^1 = \alpha_{m,q} * \psi_{m,q}(\zeta)$, one of two things happens. Either

$$\zeta^1 \cap \{z : |z| \leq 1\} \subset D(L_3L_2UC), \tag{6.8.1.2}$$

or, in a few cases, the endpoint of $\psi_{5,q}(\zeta)$ is in $C_{7,q}$, and is subject to a further quarter twist, and possibly even further fractional twists. If this happens then the path $\zeta^1 = \alpha_{m,q} * \psi_{m,q}(\zeta)$ has first unit disc crossing either along the leaf joining $e^{\pm 2\pi i(9/28)}$, with the remainder of the path in $D(L_3L_2BCL_1R_2BC)$, or with all of the path beyond the first unit disc crossing in

$$D(L_3^2(L_3L_2R_3)^3).$$

In both cases, we have

$$\zeta^1 \cap \{z : |z| \leq 1\} \subset U^1 \tag{6.8.1.3}$$

We have not defined U^1 in this work, but it is mentioned in 3.4. In both cases, all further iterations give paths with the same properties. If (6.8.1.2) holds, the path $\beta \in R_{m,0}$ such that $\sigma_\beta \circ s$ and $\sigma_\zeta \circ s$ are Thurston equivalent satisfies $w_1'(\beta) \in D(L_3L_2UC)$. If (6.8.1.3) holds, it seems likely that there is no path $\beta \in R_{m,0}$ such that $\sigma_\beta \circ s$ and $\sigma_\gamma \circ s$ are Thurston equivalent, but that there is such a path in $R_{m,1}$, by continuing the iteration. But it is not possible to be certain of this, because the paths in $R_{m,1}$ have never been completely described. $\square$

6.9. Idea of the proof of 6.4

The proof of the main theorems 6.1 and 6.2 is now reduced to proving 6.4. We start with a set $\mathcal{U}$ of capture paths γ such that $\beta_i(\gamma)$ is constant for $\gamma \in \mathcal{U}$ and $i \leq 4n_0 + 2$. We have

$$D_0 = D(\beta_{4n_0+1}(\gamma), \beta_{4n_0+2}(\gamma)) = D(\beta_{0,1}(\gamma), \beta_{0,2}(\gamma))$$

for any $\gamma \in \mathcal{U}$.

Our set U_1 will be of the form

$$U_1 = \cap_{i=0}^{\infty} U_{1,i}$$

where $U_{1,0} = U = D_0$ and $U_{1,i}$, where $U_{1,i}$ is a union of sets $\psi_{1,4n_i-1}^{-1}(D_i)$, where $\beta_j(\gamma)$ is constant on D_i for $j \leq 4n_i$, and hence $\psi_{1,4n_i-1}^{-1}$ is also constant on D_i and $\psi_{4n_{i'}+1,4n_i-1}^{-1}$ is constant on D_i for all $0 \leq i' < i$, and hence $D_{i'}$ is constant on D_i for all $0 \leq i' < i$, the numbers $n_{i'}$ are constant on D_i for all $i' \leq i$, and the sets $E_{4j-3,4j-1}$ are constant on D_i for all $j \leq 4n_i$. Then $U_{1,i}$ will be the union of sets $\psi_{1,4n_i-1}(D_i)$ such that the conditions of 2a), 2b), 2c) of 6.4 which concern β_j for $j \leq 4n_i$, $D_{i'}$ for $i' \leq i$, and $E_{4j-3,4j-1}$ for $j \leq n_i$, all hold. We then need to show that

$$\prod_{i=0}^{N} \frac{\#(U_{1,i+1} \cap Z_n)}{\#(U_{1,i} \cap Z_n)} \cdot \#(U \cap Z_n)$$

is bounded from 0 for all sufficiently large n. So we need to show that this product is convergent.

Obviously we are going to use induction. Given $U_{1,i(r)}$, such that conditions 1 to 3 hold for sets D_i for $i \leq i(r)$ such that $U_{1,i}$ is a union of sets $\psi_{1,4n_i-1}(D_i)$, we need to choose $U_{1,i(r+1)}$ so that the conditions are satisfied. So for each fixed $D_i(r)$ in $\psi_{1,4n_{i(r)-1}}^{-1}(U_i)$, we need to determine the possible D_i for $i < i(r+1)$ with $D_{i+1} \subset D_i \subset D_{i(r)}$, and for this set of $D_{i(r+1)}$, and $i(r) \leq i < i(r+1)$, it suffices to have

$$\#(D_{i+1}) \cap Z_n) \geq (1 - \delta_i) \cdot \#(D_i \cap Z_n),$$

where $\sum_i \delta_i < \infty$, because

$$\#(\psi_{1,4n_{i(r)}-1}(D_{i(r)} \cap Z_n)) = \#(D_{i(r)} \cap Z_n),$$

and similarly for $i(r+1)$.

To give the idea of the proof, it is easiest to describe what could be done if the situation were somewhat simpler.

(i) Suppose it were true that $\psi_{\omega_1,\omega_3} =$ identity on all principal arcs $I(\zeta_1, \zeta_2) \subset D_{i(r)}$ for which $I(\zeta_1, \zeta_2)$ is not contained in $D(u)$ for some word u with $|u| > \exp(|u|)^{\alpha_1}$ and for all $(\omega_t : 1 \leq t \leq 4)$ with $E_{\omega_1,\omega_3} \subset D_{i(r)}$, again with E_{ω_1,ω_3} not contained in any such $D(u)$.

(ii) Suppose also that it were true that if C were a component of $(\sigma_{\beta_{4\ell-3}} \circ s)^{-n}(E_{4\ell-3,4\ell-1})$ for $\ell < n_i$ then either $C \cap D_{i(r)} = \emptyset$ or $D_{i(r)} \subset C$.

Then it would be easy to choose D_{i+1} and D_j for $i(r-1) \leq i < j \leq i(r+1)$. But none of these is true. There is always at least one component C of $(\sigma_{\beta_1} \circ s)^{-n}(E_{1,3})$ for some n, which is not contained in $D_{i(r)}$, and corresponding components of $(\sigma_{\beta_{4\ell-3}} \circ s)^{-n}(E_{4\ell-3,4\ell-1})$, for each $\ell \leq n_i$. Indeed $C \cap Z_n$ is a positive proportion of $D_i \cap Z_n$, bounded from 0. So we need to choose u_{n_i} and D_i carefully to ensure that $I(\beta_{2j-1}, \beta_{2j})$ is disjoint from this C, for $j \leq 2n_i$. The size of $|u_{2n_{i+1}+1}| - |u_{2n_i+1}|$ is chosen so that such a choice will be possible, with a set of positive density.

First we consider (i). Every type 2 arc in D_i is crossed by some component C of $(\sigma_{\omega_1} \circ s)^{-n}(E_{\omega_1,\omega_3})$ for any quadruple (ω_i). But the crossing is usually on the central arc $\ell(C)$ of $C$and if E_{ω_1,ω_3} is already small then whenever $(\omega_t : \leq t \leq 4) = (\beta_{-4+4l+t} : 1 \leq t \leq 4)$ and some ℓ, it is possible to bound the size of the component of

$$\cup_{0<m,n_i<j}(\sigma_{\beta_{4j-3}} \circ s)^{-m}(E_{4j-3,4j-1})$$

containing C. In fact, we shall do that as follows. We define $k(C) = k(\omega_1, \omega_2)$, where ω_1 and ω_2 are the two paths in $R_{m,0}$ which bound C, as in 4.8.2. Then $k(C)-1$ is constant on each of the two components E_1 and E_2 of $\text{int}(C)$, and if $\{D(v) : w'_{k(C)+2}(v) = u\}$ intersects C, it is contained in C. We shall ensure that if $k(C) \geq 3$ then any component of $(\sigma_{\beta_{4\ell-3}} \circ s)^{-n}(E_{4\ell-3,4\ell-1})$ which intersects C is contained in

$$C_1 \cup \bigcup \{D(v) : k(v) \geq k(C) + 2,\ w'_{k(C)+2}(v) = u \text{ for some } D(u) \subset C_1\}$$

where C_1 is the corresponding component of $(\sigma_{\beta_1} \circ s)^{-n}(E_{1,3})$, or

$$C_1 \cup C_2 \cup \bigcup \{D(v) : k(v) \geq k(C) + 2,\ w'_{k(C)+2}(v) = u \text{ for some } D(u) \subset C_1 \cup C_2\}$$

where C_2 is a component of $(\sigma_{\beta_1} \circ s)^{-p}(E_{1,3})$ which intersects $E_{1,3}$. This helps us to obtain the required bound on satellites. We aim to show that, for $n_{i(r-1)} \leq \ell$ any component of $(\sigma_{\beta_{4\ell-3}} \circ s)^{-n}(E_{4\ell-3,4\ell-1})$ is disjoint from $I(\beta_{2j-1}, \beta_{2j})$ for $n(i(r-1)) \leq 2j \leq n(i(r+1))$, except when $(\beta_{2j-1}, \beta_{2j})$ is purely satellite.

Now we consider (ii). There is always at least one such C such that $C \cap D_i \neq \emptyset$ and $C \setminus D_i \neq \emptyset$. In fact, if C is a component of $(\sigma_{\beta_1} \circ s)^{-n}(E_{1,3})$, then the proportion of points in $C \cap D_i \cap Z_k$ is a positive proportion of points in $D_i \cap Z_k$ for all sufficiently large k, independent of i and D_i. But the proportion of sets D_i for which it will be impossible to choose n_{i+1} and D_{i+1} such that $C \cap D(\beta_{2j-1}, \beta_{2j}) \cap C = \emptyset$ for $j \geq 2n_{i+1}$ is of proportion $< \lambda^{(\log n_i)^\alpha}$ for some fixed $\lambda < 1$, and we can choose $\alpha > 1$. In this case, this is $< n_i^{d \log \lambda}$ for any $d > 0$ for sufficiently large i, and since $n_i > i$, we have

$$\sum_i \lambda^{(\log n_i)^\alpha} < +\infty.$$

In all cases, we will make our initial choice so that every component C' of $(\sigma_{\beta_{4\ell-3}} \circ s)^{-n'}(E_{4\ell-3,4\ell-1})$ which intersects this C is either contained in it or contains D_i. So then if $C' \subset C$ we can ensure that $D_{i+1} \cap C = \emptyset$, and hence $D_{i+1} \cap C' = \emptyset$.

The method which is used is loosely, but quite closely, related to standard techniques of proving non-uniform hyperbolicity properties — as in [**17**], for example, but the idea probably occurs in every single paper on the subject. We need to exclude certain behaviours of the critical orbit. Early and long recurrence needs to be excluded, which, in general terms, is what is needed to prove non-uniform hyperbolicity – in the cases which can be handled by methods currently available. Something more needs to be excluded because of the group action, but everything that has to be excluded is related to early long recurrence — what were known as "followers" in [**17**].

6.10. Proof of 6.4

One basic idea of the proof is to translate everything into symbolic dynamics. This means that we use the admissible words w in our standard alphabet

$$\{BC, UC, L_i, R_j : 1 \leq i, j \leq 3\}$$

and the sets $D(w)$. The number of admissible words of length n is boundedly proportional to 2^n, and if w_0 and z_0 are admissible and of lengths m and p respectively, then the number of admissible words of length n and starting with w_0 and ending with z_0 is boundedly proportional to 2^{n-m-p}.

Fix $\gamma \in \mathcal{U}$ and write $\beta_j = \beta_j(\gamma)$. As in 6.4, we define u_j to be the longest word such that

$$D(\beta_{2j-1}, \beta_{2j}) \subset D(u_j).$$

One thing which we need to determine is how u_j differs from a prefix of $w(\gamma)$. Putting $j = 2n_i + 1$, we have

$$D_i \subset D(u_{2n_i+1}).$$

The conditions of 6.4 ensure that u_{2n_i+1} is a prefix of u_j for all $j \geq 2n_i + 1$. In particular, u_{2n_i+1} is a prefix of $u_{n(\gamma)}$ for all $i \geq 0$ and u_{2n_j+1} is a prefix of u_{2n_i+1} for all $j \leq i$. It is not the case, however, that u_{j_1} is a prefix of u_{i_1} for all $j_1 \leq i_1$, and not even if $i_1 = 2n_i + 1$ for some i. Still, if j and i are the largest integers with $2n_j + 1 \leq j_1$ and $2n_i + 1 \leq i_1$, then u_{2n_j+1} is a prefix of u_{2n_i+1}, and hence is also a prefix of u_{i_1}. Since $|u_{j_1}| - |u_{2n_j}| = o(|u_{j_1}|)$, this is useful information.

We also define $u_{j,1}$ to be the longest word such that

$$\psi_{1,4\lfloor j/2\rfloor-1}(D(\beta_{2j-1},\beta_{2j}))\subset D(u_{j,1}),$$

and $u_{2j-1,-}$ is the longest word with

$$E_{4j-3,4j-1}\subset D(u_{2j-1,-}).$$

For the choice of $(\beta_i : 1\le i\le 4)$ for proving Theorem 6.1, there is no single nonempty word $u_{1,-}$ with $E_{1,3}\subset D(u_{1,-})$. So in this case, we define $u_{1,-}$ to be a pair of words $(u_{1,1,-},u_{1,2,-})$ with $D(u_{1,1,-})\subset D(L_3)$ and $D(u_{1,2,-})\subset D(BC)$, where $u_{1,1,-}$ and $u_{1,2-}$ are the longest possible with

$$E_{1,3}\subset D(u_{1,1,-})\cup D(u_{1,2,-}).$$

We also define

$$D(u_{1,-})=D(u_{1,1,-})\cup D(u_{1,2,-}).$$

Our set U_1, which we want to construct to prove Theorem 6.4, will be a union of sets $u_{2n(\gamma),1}$. By the inductive construction there is a one-to-one correspondence between the words $u_{j,1}$ and u_j. We will be choosing the set U_1 to be a union of sets $D(u_{2n(\gamma),1})\cap Z_\infty$ by choosing the union of sets $D(u_{2n(\gamma)})\cap Z_\infty)$ and we want the union of sets $D(u_{2n(\gamma)}\cap Z_m)$ to be of positive density for all sufficiently large m. So, of course, we need to control the difference between u_j and $u_{j,1}$.

We want to ensure that any changes between u_j and $u_{j,1}$ can be controlled by looking either at the different prefixes u_{2n_i+1} in u_j, or beyond the largest such prefix. In order to do this, we shall ensure the following.

(i) For $j>2n_i+1$, for any suffix a_1 of a such that $a\leftrightarrow b$ is a basic exchange with corresponding letters of a and b all different, any occurrence of $a_1u_{1,-}$ which is a subword of u_j, and which starts before the end of the prefix u_{2n_i+1} in u_j, is a subword of u_{2n_i+1}.

(ii) No non-prefix subword of u_{2n_i+1} is equal to $u_{2n_{i(r-1)}+1}$, for $i\le i(r+1)$.

We claim that these two conditions are enough to ensure the conditions a) and b) of 6.4. The reason is as follows. The only way that a) or b) can fail is if $D(v)$ is moved outside $D(u_{2n_i+1})$ by $\psi_{4n_i+1,4\ell-1}$ for some $\ell>n_i$, and for some word v which is an extension of u_{2n_i+1}, but a prefix of $u_{2n_{i+1}+1}$. Now $\psi_{4\ell+1,4\lfloor (j-1)/2\rfloor-1}$ is a composition of disc exchanges for each $2\ell<j-1$. If we look just at the disc exchanges in the composition which have an effect on $D(u_j)$, we see that the support of each one is either contained in $D(u_{2n_i+1}$ or contains it, for each i with $2n_i+1\le j$. So the disc exchanges which have an effect on $D(u_j)$ can be permuted, and we can group together those whose supports contain $D(u_{2n_i+1})$ together, for each i. Then we can permute the order in which these occur. We can apply this corresponding to the largest i first, and then apply them in decreasing order of i. This makes sense,

because then all the disc exchanges, in the composition of $\psi_{4\ell+1,4\lfloor (j-1)/2\rfloor -1}$ which have an effect on $D(u_j)$, have support contained in $D(u_{2n_i+1})$ for any $n_i \leq 2\ell$. So $\psi_{4\ell+1,4\lfloor (j-1)/2\rfloor -1}(D(u_j))$ is of the form $D(u_{j,\ell+1}$ for some word $u_{j,\ell+1}$ where $u_{j,\ell+1}$ has u_{2n_i+1} as a prefix for any $2n_i \leq \ell+1$. For those disc exchanges which have support between $D(u_{2n_i+1})$ and $D(u_{2n_{i+1}+1})$, it is not possible to commute in any order, but the first exchange which is applied can only give a change before an occurrence of $u_{1,-}$, or before a word arrived at by a rather special sequence of disc exchanges preceding some $u_{2\ell-1,-}$.

Clearly, in order to satisfy these conditions, we need to choose the n_i and u_{2n_i+1} suitably, given u_j. In fact, we will choose n_i and u_{2n_i+1}, for $i \leq i(r)$, given an extension u of $u_{2n_{i(r)}+1}$. From the statement of 6.4, we want the conditions

$$|u_{2n_{j+1}+1}| - |u_{2n_j+1}| \leq (\log |u_{2n_j+1}|)^{\alpha_1} = o(|u_{2n_j+1}|)$$

and

$$|u_{2n_{2i(r+1)}+1}| \geq \exp(\log |u_{2n_{2i(r)}+1}|)^{\alpha_2})$$

From this, and the corresponding lower bound on $|u_{2n_{i(r)}+1}|$ in terms of $|u_{2n_{i(r-1)}+1}|$, we have

$$|u_{2n_{i(r+1)}+1}| \geq \exp(\log |u_{2n_{i(r-1)}+1}|^{\alpha_2^2})$$

Recall that we are aiming to construct U_1 as an intersection of sets $U_{1,r}$. We shall construct $U_{1,r}$ as a union of sets $D(u)$ for prefixes $u_{2n_i+1,1}$ have been defined for $i \leq i(r)$ with

$$|u| - |u_{2n_{i(r)}+1}| \leq (\log |u_{2n_{i(r)}+1}|)^{\alpha_1/4})$$

We shall do this by constructing $U'_{1,r}$ with

$$U_{1,r} = \psi_{4n_{i(r)}+1}(U'_{1,r}).$$

The method is inductive. For a sequence of integers m_r, we will choose $U'_{1,r}$ to be a union of sets $D(u)$ with $|u| = m_r$. We start by taking $m_0 = |u_{2n_0+1}|$ but for $r > 0$, we will have $m_r > |u_{2n_{i(r)}+1}|$ for each choice of $u_{2n_{i(r)}+1}$. Since we can take $|u_{2n_0+1}|$ arbitrarily large, we can also take m_0 arbitrarily large. We shall construct $U'_{1,r+1}$ as a union of sets $D(v)$ where each $D(v)$ is a subset of one of the sets $D(u)$ in the union $U'_{1,r}$, that is, v has some u as a prefix, where $D(u)$ is part of the union $U'_{1,r}$. We shall ensure that

$$\#(U'_{1,r+1} \cap Z_{m_{r+1}}) \geq (1 - \delta'_r)\#(U'_{1,r} \cap Z_{m_{r+1}}).$$

where $\sum \delta'_r$ converges, and, in fact, can be taken arbitrarily small.

So it remains to show that, given u such that $D(u)$ is one of the sets in the union $U'_{1,r}$, we can find a high density set of extensions v for which the properties are satisfied. Note that $i(r)$, $n_{i(r)}$ and $u_{2n_{i(r)}+1}$ all depend on u. The numbers $i(r+1)$, n_i for $i(r) < i \leq i(r+1)$, and the prefixes u_j

for $2n_i(r)+1 < j \le 2n_{i(r+1)}+1$, will all depend on the extension v of u. But we need to find a good proportion of extensions v of u for which these numbers and prefixes can be defined with the appropriate properties. We will choose m_{r+1} so that

$$\exp((\log m_r)^{\alpha_2/2} \le m_{r+1} \le \exp\log(m_r^{2\alpha_2}).$$

Note that this also implies that

$$\exp((\log m_{r-1})^{\alpha_2^2/4} \le m_{r+1} \le \exp\log(m_{r-1}^{\alpha_2^2}),$$

or otherwise put,

$$(\log m_{r+1})^{\alpha_2^{-2}} \le m_{r-1} \le (\log m_{r+1})^{4\alpha_2^{-2}}.$$

So fix u. Then $i(r)$ is defined depending on u and n_i and u_{2n_i+1} are defined for each $i \le i(r)$, depending on u, such that u_{2n_i+1} is a prefix of u. To satisfy conditions (i) and (ii), we need to obtain a lower bound on the number of v extending u such that

- every subword of length $(1/10)(\log m_r)^{\alpha_1}$ in the extension of u in v contains a word L_3x_0 such that $D(L_3x_0) \cap D(u_{1,-}) = \emptyset$ and $u_{1,-}$ is not a suffix of L_3x_0;
- $u_{2n_{i(r-1)}+1,-}$ is not a subword of the extension of u in v.

If $r = 0$ then in the second condition we simply replace $u_{2n_{i(r-1)}+1,-}$ by $u_{2n_{i(r)}+1,-}$.

These conditions are sufficient to ensure (i) and (ii) respectively. In order to ensure (i) we can take a particular choice of L_3x_0. A choice of L_3x_0 which will work for all our different choices of $u_{1,-}$ is $(L_3(L_2R_3)^3)^2$. So now we need to estimate the proportion of extensions v of u of length m_{r+1} for which these conditions are not satisfied. The proportion of words of length N which do not contain an occurrence of L_3x_0 is $\le e^{-N\lambda}$ for some $\lambda > 0$ which depends only on L_3x_0 – which we have fixed. We want to avoid such words, with $N \ge (1/10)(\log m_r)^{\alpha_1}$, over length m_{r+1}. So the proportion of words that we want to avoid for (i) is at most

$$m_{r+1}\exp(-(\lambda/10)(\log m_r)^{\alpha_1} \le \exp((\log m_r)^{2\alpha_2} - (\lambda/10)(\log m_r)^{\alpha_1}).$$

This proportion is at most

$$\exp(-(\log m_r)^{\alpha_1/2}$$

provided that $\alpha_2 < \alpha_1/3$, that is, provided that α_1 is sufficiently large given α_2. The only requirement on α_2 is that $\alpha_2 > 1$. Similarly for (ii), for which it suffices to avoid $u_{2n_{i(r-1)}+1,-}$, the proportion of words which we want to avoid is, for a suitable $\lambda > 0$,

$$\le m_{r+1}\exp(-\lambda m_{r-1}) \le \exp((\log m_{r-1})^{4\alpha_2^2} - \lambda m_{r-1}).$$

Assuming, as we may do, that m_{r-1} is sufficiently large given λ and α_2, this is at most

$$\exp(-\lambda m_{r-1}/2).$$

So altogether the proportion that has to be avoided is

$$\sum_{r\geq 0}\exp(-(\log m_r)^{\alpha_1/2})+\sum_{r\geq 0}\exp(-\lambda m_{r-1}/2)$$

where in the second sum we define $m_{-1}=m_0$. This sum can be taken arbitrarily small by taking m_0 arbitrarily large for fixed $\lambda > 0$ and $\alpha_1 > 2$. This completes the construction of $U_{1,r+1}$ from $U_{1,r}$, and hence completes the proof.

Bibliography

1. M. Aspenberg and M. Yampolsky, *Mating of nonrenormalisable polynomials*, Comm. Math. Phys. **287** (2009), 1–40.
2. J. Avila A., Kahn and M. Lyubich, *Rigidity of unicritical polynomials*, Ann. of Math. **170** (2009), 783–797.
3. V.P. Chistyakov and B.A. Sevast'yanov, *Asymptotic normality in the classical ball problem*, Theory of probability and its applications **9** (1964), 198–211.
4. A. Douady and J.H. Hubbard, *Etudes dynamiques des polynômes complexes, avec la collaboration de P. Lavaurs, Tan Lei, P. Sentenac. Parts I and II*, Publications Mathématiques d'Orsay, 1985.
5. ———, *A proof of Thurston's topological characterization of rational functions*, Acta Math. **171** (1993), 263–297.
6. J.H. Hubbard, *Local connectivity of Julia sets and bifurcation loci: three theorems of J.C. Yoccoz*, Topological methods in modern mathematics, Publish or Perish, 1993, volume issued in association with conference in honour of John Milnor's sixtieth birthday, SUNY at Stony Brook, 1991, pp. 467–511.
7. J. Kahn, *A priori bounds for some infinitely renormalizable quadratics: I. bounded primitive combinatorics*, http://front.math.ucdavis.edu/math.DS/0609045.
8. J. Kahn and M. Lyubich, *A priori bounds for some infinitely renormalizable quadratics: II. Decorations*, Ann. Sci. Ec. Norm. Sup. **41** (2008), 57–84.
9. ———, *Local connectivity of julia sets for unicritical polynomials*, Ann. of Math. **170** (2009), 413–426.
10. ———, *A priori bounds for some infinitely renormalizable quadratics: III. Molecules*, Complex Dynamics: Families and Friends (D. Schleicher and N. Selinger, eds.), A.K. Peters, 2009, the proceedings of the conference in honour of J.H. Hubbard's 60th birthday, pp. 229–254.
11. ———, *The quasi-additivity law in conformal geometry*, Ann. of Math. **170** (2009), 561–593.
12. J. Kiwi and M. Rees, *Counting hyperbolic components*, See http://arxiv.org/pdf/1003.6104v1.
13. M. Lyubich, *Dynamics of quadratic polynomials III: Parapuzzles and SBR measures*, Geometrie complexe et systemes dynamiques, Asterisque, vol. 261, Société Mathématique de France, 2000, Volume arising from conference in honour of sixtieth birthday of Adrien Douady, Orsay 1995, pp. 173–200.
14. D. Meyer, *Expanding Thurston maps as quotients*, Preprint.
15. ———, *Unmating of rational maps, sufficient criteria and examples*, http://arxiv.org/pdf/1110.6784.
16. ———, *Invariant Peano curves of expanding Thurston maps*, To appear in Acta Math, 2011.

17. M. Rees, *Positive measure sets of ergodic rational maps*, Ann. Scient. Ec. Norm. Sup. **19** (1986), 383–407.
18. ______, *Components of degree two hyperbolic rational maps*, Invent. Math. **100** (1990), 357–382.
19. ______, *A partial description of the parameter space of rational maps of degree two: Part 1*, Acta Math. **168** (1992), 11–87.
20. ______, *A partial description of the parameter space of rational maps of degree two: Part 2*, Proc. Lond. Math. Soc. **70** (1995), 644–690.
21. ______, *Views of parameter space: topographer and resident*, Astérisque, vol. 288, Société Mathématique de France, 2003.
22. ______, *A fundamental domain for* V_3, Mem. Soc. Math. de France **121** (2010), 1–139.
23. ______, *Multiple equivalent matings with the aeroplane polynomial*, Ergod. Th. and Dynam. Syst. **30** (2010), 1239–1257, with erratum, p.1259.
24. P. Roesch, *Holomorphic motions and puzzles (following m. shishikura)*, The Mandelbrot Set, Theme and Variations, LMS Lecture Note Series, vol. 274, Cambridge U. P., 2000, pp. 117–132.
25. Lei Tan, *Matings of quadratic polynomials*, Ergod. Th. and Dynam. Sys. **12** (1992), 589–620.
26. W.P. Thurston, *On the combinatorics of iterated rational maps*, Complex dynamics: families and friends (D. Schleicher and N. Selinger, eds.), AK Peters, 2009, Article orginally appeared as a Princeton University and I.A.S. preprint in 1985. Volume dedicated to 60th birthday of J.H. Hubbard. Editorial notes by D. Schleicher and N. Selinger, pp. 3–109.
27. V. Timorin, *External boundary of* M_2, Proceedings of the Fields Institute dedicated to the 75th birthday of J. Milnor, Fields Institute Communications, vol. 53, AMS, 2006, pp. 225–267.
28. B. Wittner, *On the bifurcation loci of rational maps of degree two*, Ph.D. thesis, Cornell University, 1988.